

Woods Hole Oceanographic Institution  
ATLAS - GAZETTEER COLLECTION

NOAA Technical Memorandum NMFS



JANUARY 1988

**ICHTHYOPLANKTON AND STATION DATA FOR  
CALIFORNIA COOPERATIVE OCEANIC FISHERIES  
INVESTIGATIONS SURVEY CRUISES IN 1965**

Elizabeth G. Stevens  
Richard L. Charter  
H. Geoffrey Moser  
Larry R. Zins

PLEASE RETURN  
TO  
INSTITUTION DATA LIBRARY  
**McLEAN**

NOAA-TM-NMFS-SWFC- 96

U.S. DEPARTMENT OF COMMERCE  
National Oceanic and Atmospheric Administration  
National Marine Fisheries Service  
Southwest Fisheries Center

QL  
639.25  
JZ8  
No. 96

1255-AB  
Atlas  
Shef  
[series]  
1965

## NOAA Technical Memorandum NMFS

The National Oceanic and Atmospheric Administration (NOAA), organized in 1970, has evolved into an agency which establishes national policies and manages and conserves our oceanic, coastal, and atmospheric resources. An organizational element within NOAA, the Office of Fisheries is responsible for fisheries policy and the direction of the National Marine Fisheries Service (NMFS).

In addition to its formal publications, the NMFS uses the NOAA Technical Memorandum series to issue informal scientific and technical publications when complete formal review and editorial processing are not appropriate or feasible. Documents within this series, however, reflect sound professional work and may be included in scientific and technical literature.

OUT

1987-. Ichthy  
Oceanic Fisher  
SOUTHWEST FISH  
SHF(series) /c

Woods Hole Oceanographic Institution  
ATLAS - GAZETTEER COLLECTION

1765-AB  
Atlas Sheet  
[Series]  
1965

**NOAA Technical Memorandum NMFS**

This TM series is used for documentation and timely communication of preliminary results, interim reports, or special purpose information, and have not received complete formal review, editorial control, or detailed editing.



**JANUARY 1988**

**ICHTHYOPLANKTON AND STATION DATA FOR  
CALIFORNIA COOPERATIVE OCEANIC FISHERIES  
INVESTIGATIONS SURVEY CRUISES IN 1965**

Elizabeth G. Stevens  
Richard L. Charter  
H. Geoffrey Moser  
Larry R. Zins

Southwest Fisheries Center  
National Marine Fisheries Service  
La Jolla, CA 92038

**NOAA-TM-NMFS-SWFC-96**

U.S. DEPARTMENT OF COMMERCE  
C. William Verity, Jr., Secretary  
National Oceanic and Atmospheric Administration  
Anthony J. Calio, Administrator  
National Marine Fisheries Service  
William E. Evans, Assistant Administrator for Fisheries

MBI WHOI Library  
00301100002985



## CONTENTS

	Page
List of Figures .....	iii
List of Tables .....	iv
Abstract .....	1
Introduction .....	1
Sampling Area and Pattern .....	2
Sampling Gear and Methods .....	3
Laboratory Procedures .....	4
Identification .....	5
Computer Entry and Editing .....	10
Species Summary .....	10
Explanation of Tables .....	11
Acknowledgments .....	12
Literature Cited .....	13
Figures .....	16
Tables .....	22
Index .....	216

## LIST OF FIGURES

	Page
Figure 1. Composite arrangement of diagrammatic charts showing areas sampled on each CalCOFI cruise during 1965 .....	16
Figure 2. Station pattern for CalCOFI Cruise 6501 showing tracks for each vessel .....	17
Figure 3. Station pattern for CalCOFI Cruise 6504 .....	18
Figure 4. Station pattern for CalCOFI Cruise 6507 .....	19
Figure 5. Station pattern for CalCOFI Cruise 6509 .....	20
Figure 6. The basic station plan for CalCOFI cruises from 1950 to the present .....	21

## LIST OF TABLES

	Page
Table 1. Station and plankton tow data for CalCOFI cruises in 1965 .....	22
Table 2. Pooled occurrences of fish larvae taken during CalCOFI cruises in 1965 .....	48
Table 3. Pooled numbers of fish larvae taken during CalCOFI cruises in 1965 .....	52
Table 4. Numbers of fish larvae taken on stations occupied during CalCOFI cruises in 1965 .....	56
Table 5. Summary of pooled occurrences of fish larvae taken on CalCOFI cruises from 1961-1969 .....	211
Table 6. List of stations with multiple occupancies in one month during 1965 .....	215



## ABSTRACT

This report provides ichthyoplankton and associated station and tow data from California Cooperative Oceanic Fisheries Investigations (CalCOFI) cruises conducted off California and Baja California in 1965. It is the fifteenth report in a series that presents these data for all biological-oceanographic CalCOFI surveys from 1951 to the present. A total of 1086 stations was occupied during 4 quarterly multivessel cruises over a survey area which extended from Pt. Reyes, California to Cape San Lazaro, Mexico and seaward to several hundred miles. The data are listed in a series of 6 tables; the background, methodology, and information necessary for interpretation and quantitative analysis of the data are presented in an accompanying text. All pertinent station and tow data, including volumes of water strained and standard haul factors, are listed in the first table. Another key table lists, by station and month, standardized counts of each of the 161 larval fish categories identified from survey samples. This and previous and subsequent reports make the CalCOFI ichthyoplankton and station data available to all investigators and serve as guides to the newly developed computer data base.

## INTRODUCTION

This report, the fifteenth of a series, provides ichthyoplankton and associated station and tow data from California Cooperative Oceanic Fisheries Investigations (CalCOFI) joint biological-oceanographic survey cruises conducted in 1965. This program was initiated in 1949, under the sponsorship of the Marine Research Committee of the State of California, to study the population fluctuations of the Pacific sardine (*Sardinops sagax*) and the environmental factors that may play a role in such fluctuations. CalCOFI, known as the California Cooperative Sardine Research Program from 1949 to 1953, was made up of representatives of the South Pacific Fisheries Investigations (SPFI) of the U.S. Fish and Wildlife Service [now the La Jolla Laboratory, National Marine Fisheries Service (NMFS)], the Scripps Institution of Oceanography (SIO), the California Department of Fish and Game (CDFG), the California Academy of Sciences (CAS) and the Hopkins Marine Station of Stanford University. The first three of these agencies supplied ships and personnel to conduct the sea surveys. NMFS processed the plankton samples and analyzed the ichthyoplankton from them. SIO processed and analyzed the hydrographic samples and measurements and also analyzed invertebrate groups from the plankton samples.

The boundaries, station placement, and sampling frequency for the CalCOFI survey area were based on the results of joint biological and oceanographic cruises conducted by NMFS and SIO during 1939-41. Those cruises were designed to collect sardine eggs and larvae and associated hydrographic data over the entire areal and seasonal spawning range of the species. On these survey cruises, plankton tows were made to 70 m, a depth which

encompassed the vertical distribution of sardine eggs and larvae. Wide-ranging joint biological and oceanographic survey cruises were resumed in 1949 with sardine as the focus; however, an increasing interest in other biological components resulted in the deepening of standard tows to 140 m in 1951. This marked the beginning of truly quantitative ichthyoplankton sampling on CalCOFI surveys.

Data resulting from CalCOFI surveys in 1965 have been published in a number of forms. Hydrographic data (Univ. of Calif., SIO, 1965, 1967) and zooplankton volumes (Smith, 1971) were presented in standard formats. Distributional maps of larvae of 5 taxa taken on CalCOFI surveys during 1965 are presented in the CalCOFI Atlas series: northern anchovy (*Engraulis mordax*), Kramer and Ahlstrom, 1968; jack mackerel (*Trachurus symmetricus*) and Pacific hake (*Merluccius productus*), Ahlstrom, 1969; Pacific sardine (*Sardinops sagax*), Kramer, 1970; rockfish (*Sebastes spp.*), Ahlstrom et al., 1978.

A computer data base for eggs and larvae of sardine and anchovy, for larvae of hake, jack mackerel and Pacific mackerel (*Scomber japonicus*), and for eggs of Pacific saury (*Cololabis saira*) was established in 1969. The development of a data base for other fish larvae is a complex undertaking because competency of identification has evolved steadily over the past 38 years. We began the task of producing a CalCOFI ichthyoplankton data base and associated data report series in 1983. All available original records for 1965 were subjected to an extensive verification and editing process to produce this report. This and previous (Ambrose et al., 1987a,b,c; 1988; Sandknop et al., 1987a,b; 1988a,b; Stevens et al., 1987a,b,c; Sumida et al., 1987a,b; 1988) and subsequent reports make the CalCOFI ichthyoplankton and station data available to all investigators and serve as guides to the computer data base. The data base will be modified when additional errors are discovered and when composite taxa from the earlier years are reidentified. These reports are the fundamental reference documents against which subsequent changes in the data base can be compared.

#### SAMPLING AREA AND PATTERN

In 1965, CalCOFI survey cruises were conducted at quarterly intervals. A total of 1086 stations included in this data base was occupied on 4 cruises, with an average of 272 stations per cruise (range 193-311). Coverage of the survey station pattern varied among cruises and the entire survey area was not covered on any single cruise (Figures 1-5, Table 1). The area off northern California (lines 40-57) was not covered in 1965. The area off central California (lines 60-77) was covered in January, April, and July. The area between Pt. Conception, California and Pt. San Juanico, Baja California (lines 80-137) was surveyed on all cruises and coverage extended to line 140, off Cape San Lazaro, in February and August. Coverage extended seaward to station 200 (approximately 600-700 miles offshore) on lines 60,

70, 80, and 90 during Cruises 6501 and 6507 but typically did not extend beyond station 90 or 100 (approximately 200-300 miles offshore)<sup>1</sup>. Some inshore stations were occupied during 1965 which were not covered on early CalCOFI surveys. These stations were included in the data base but omitted from the station plots (Figures 2-5).

Two vessels were employed on these cruises: the *Black Douglas* of NMFS and the *Alexander Agassiz* of SIO. Both vessels participated on each cruise except Cruise 6509, in which only the *Black Douglas* was used (Univ. of Calif., SIO 1965, 1967).

#### SAMPLING GEAR AND METHODS

The standard CalCOFI net used from 1949 to 1969 had a 1-m diameter mouth opening ( $0.785 \text{ m}^2$  area) and an overall length of about 5 m. The net was constructed of 30xxx gauze, a heavy duty grade of silk bolting cloth, with a mesh size of 0.55 mm after shrinkage. The last 40 cm of the cone and the cod end were constructed of 56xxx grit gauze which had a mesh size of 0.25 mm after shrinkage. The net ring was fastened to a short 3-lead bridle connected to several meters of line which attached to the towing cable by a clamp. A current meter was suspended in the center of the net mouth to measure volume of water filtered (see Kramer et al., 1972, for further details).

The standard tow from 1951 through 1968 was an oblique haul to 140 m depth (to 15 m of the bottom in shallow areas) designed to filter a constant amount of water per depth interval (ca.  $3\text{m}^3/\text{m}$  of depth) over the vertical range of most ichthyoplankters. Hauls were made at a ship speed of 1.5-2.0 knots and initiated by clamping the net line to the towing cable with the 45 kg terminal weight about 10-15 m below the surface. The net was lowered to 140 m depth by paying out 200 m of wire over a 4 minute period (35 m of depth/min.). After fishing at depth for 30 seconds, the net was retrieved at 20 m/min. (14 m depth/min.). The angle of stray of the towing cable was recorded every 30 seconds and

---

<sup>1</sup>CalCOFI lines (Figure 6) are arranged perpendicular to the coastline and extend from the Canadian border (line 10) to below Cape San Lucas, Baja California (line 157). Stations were established on the basis of a perpendicular to line 80 (off Pt. Conception) at a point designated as station 60. Stations were plotted seaward and shoreward from station 60 on each line. Cardinal CalCOFI lines (those ending in "0") are 120 miles apart and usually bracket two ordinal lines (ending in "3" or "7"), so that lines are 40 miles apart over most of the pattern. Cardinal stations are 40 miles apart and typically these are separated by a station number ending in "5" so that stations are 20 miles apart out to station 90 on most lines. Stations are placed at closer intervals near the coast and islands to accommodate these features (see Kramer et al., 1972 for further details).

maintained at 45° (+3°) by adjusting the ship speed and course. After reaching the surface, the net was washed down and the samples preserved in 5% formalin buffered with sodium borate. Flowmeter readings were made at the beginning and end of each tow. Detailed descriptions of gear and methods are given by Ahlstrom (1953), Kramer et al. (1972), and Smith and Richardson (1977).

#### LABORATORY PROCEDURES

Laboratory processing began with the determination of a displacement volume for each sample (methods described in Staff, SPFI, 1953 and Kramer et al., 1972). Zooplankton volumes (including ichthyoplankton) of samples collected in 1965 are presented graphically in Smith (1971).

Sorting involved the removal of ichthyoplankton from the sample and identification and separation of: eggs and larvae of Pacific sardine and northern anchovy; larvae of Pacific hake; and eggs of Pacific saury. In 1965, each sample was sorted completely.

A "standard haul factor" (SHF) was calculated for each tow to make them comparable and allow estimations of areal abundance. This factor adjusts the number of eggs or larvae in a haul to the number in 10 m<sup>3</sup> of water strained per meter of depth fished. If the vertical distribution of the species has been encompassed, then the adjusted value is equivalent to the number under 10 m<sup>2</sup> of sea surface. The SHF is calculated for each haul by the formula:

$$SHF = \frac{10 D}{V}$$

where D = depth of haul = cosine of the average angle of stray of the towing cable multiplied by cable length (m)

V = total volume of water (m<sup>3</sup>) strained during the haul

$$V = R \cdot a \cdot p$$

where R = total number of revolutions of the current meter during the haul

a = area (m<sup>2</sup>) of the mouth of the net

p = length of column of water (m) needed to produce one revolution of the current meter.

Tow depth, volume of water strained, and standard haul factor are listed in Table 1 for each tow taken during 1965. Detailed

descriptions of factors involved in calculating these values are presented in Ahlstrom (1948), Kramer et al. (1972), and Smith and Richardson (1977).

## IDENTIFICATION

Identification of ichthyoplankton species beyond those separated during the sorting process was carried out by a separate group of specialists. Ontogenetic stages of fishes are inherently difficult to identify and this is further complicated by the large number and diversity of species which contribute to the ichthyoplankton of the California Current region. Most identifications were accomplished by establishing ontogenetic series on the basis of morphology, meristics, and pigmentation and then identifying these series by relating them to known metamorphic, juvenile, or adult stages with overlapping features (Powles and Markle, 1984). A total of 159 taxa was identified for 1965, with 89 taken to species, 32 to genus, 32 to family, and 6 to order or suborder. Beginning in 1961, larvae in the families Paralepididae and Labridae were identified to genus or species.

The task of producing a reliable and equitable ichthyoplankton data base required extensive procedures to verify, correct, and edit the original identifications. The primary data source was the original identification sheets (see Kramer et al., 1972, for examples); however, a critical resource used in all phases of this process was the CalCOFI ichthyoplankton collection in which the samples are archived. Throughout the course of CalCOFI ichthyoplankton studies, samples have been identified to the lowest taxon possible. In reviewing these identifications for the data base, our approach has been conservative and we have preserved those identifications and counts which we could confirm, while correcting as many of the errors as possible. After computer entry, taxonomic errors and inconsistencies in the data base were corrected and the most obvious identification errors were corrected. Our current knowledge of ichthyoplankton techniques coupled with a precise understanding of the development of identification competency in the program over the years allowed us to critically judge the historical records. Identifications were changed to different taxa, lumped to a higher taxonomic category, or given a more precise taxonomic name. In some cases, identifications of a taxon were inconsistent among cruises in a year. These records were made equitable by lumping to the higher taxonomic category to avoid biases that could result in quantitative misinterpretations.

Next, statistical, seasonal, and geographic outliers were identified, employing a series of graphic summaries and listings. Examination of geographic outliers proved to be especially effective because of our accumulated knowledge of species distributions. In the course of examining samples for these outliers, other identification errors were discovered and

eventually all taxa were scrutinized to some extent. Lastly, certain taxa were reexamined in all samples for the entire CalCOFI time series. These taxa were selected because of their commercial, ecological, phylogenetic, or zoogeographic importance or because taxonomic confusion was at the ordinal level. The following is a list of the taxa for 1965 which received special attention, with explanations and caveats intended to aid in quantitative interpretations:

*Anguilliformes* - tentative and sporadic identifications to family or lower taxon lumped to order.

*Sardinops sagax* - all specimens south of line 120 checked for misidentification of *Opisthonema* spp. Four large samples on line 133 in August (stations 137.20, 137.21, 137.22 and 137.23) appear to be a mixture of primarily small *S. sagax* and *Opisthonema* spp. which cannot be separated at this time.

*Engraulis mordax* - some nearshore samples of small *E. mordax* may contain other anchovy genera which could not be differentiated.

*Nansenia* spp. - all specimens checked and identified as *N. candida* or *N. crassa*; all specimens of these species near their range boundaries checked.

*Bathylagus* spp. - includes small and/or disintegrated specimens of *Bathylagus* or *Leuroglossus stilbius*.

*Bathylagus milleri* - all specimens checked.

*Stomiiformes* - all specimens checked and identified to genus or species; residuals are small, poorly preserved specimens.

*Vinciguerria lucetia* - specimens taken seaward of station 100 checked for misidentification of *V. poweriae*; some *V. poweriae* may remain in *V. lucetia* samples from these stations because small larvae of the two species could not be differentiated; sporadic identification of *V. poweriae* began in 1961.

*Sternopychidae* - tentative and sporadic identifications of hatchetfishes to genus were lumped to family.

*Bathophilus* spp. - all specimens checked.

*Eustomias* spp. - all specimens checked.

*Photonectes* spp. - all specimens checked.

*Tactostoma macropus* - all specimens checked.

*Paralepididae* - all specimens examined and identified to species; residuals are small, poorly preserved or unavailable specimens.

*Scopelarchidae* - tentative and sporadic identifications to genus lumped to family.

*Lampanyctus* spp. - tentative and sporadic identifications to species lumped to genus.

*Lampanyctus regalis* - underrepresented because of inability to differentiate small larvae (<5 mm) from those of other species of the genus; counts may include other species of the genus because of difficulty in identifying larvae of this large and complex genus.

*Lampanyctus ritteri* - comment for *L. regalis* applies to this species.

*Stenobrachius leucopsarus* - all specimens taken seaward of station 100 checked.

*Triphoturus mexicanus* - specimens taken seaward of station 100 checked for misidentification of *T. nigrescens*.

*Centrobranchus* spp. - all specimens checked.

*Diogenichthys atlanticus* - all specimens at margins of range checked.

*Diogenichthys laternatus* - all specimens at margins of range checked.

*Electrona rissoii* - recognition of this species was inconsistent and others may be included in *Protomyctophum crockeri* or Myctophidae.

*Hygophum* spp. - all specimens reidentified to species; residuals are small, poorly preserved or unavailable specimens.

*Hygophum atratum* - all specimens checked.

*Hygophum reinhardtii* - all specimens checked.

*Protomyctophum crockeri* - some samples on northern lines may contain *P. thompsoni*, which was not identified originally.

*Physiculus* spp. - all specimens examined.

*Ophidiiformes* - this category did not exist originally and ophidiiform larvae were included in *Brosmophycis marginata*, "Otophidium", "Zoarcidae", and "blenny"; identifications of *B. marginata* proved to be mostly correct and "Zoarcidae" to be a yet unidentified ophidiiform species; all "Otophidium" and "blenny" were reexamined and the former included *Ophidion scrippsae*, *Chilara taylori* and other ophidiiform taxa (moved to order); "blenny" contained *O. scrippsae*, *C. taylori*, and other ophidiiform taxa in addition to true blennioids.

Trachipteridae - tentative and sporadic identifications to genus were lumped to family.

*Melamphaes* spp. - all identifications ascribed to Melamphaidae were reexamined and assigned to genus (*Melamphaes*, *Poromitra*) or species (*Scopeloberyx robustus*, *Scopelogadus bispinosus*). Larvae originally identified as *Melamphaes* spp. were not reexamined and this category may contain other melamphaid genera.

Cottidae - all specimens checked.

*Oxylebius pictus* - all specimens checked.

*Zaniolepis* spp. - all specimens checked.

*Sebastes* spp. - category may contain other scorpaenid genera, particularly in samples south of line 120.

Blennioidei - this is the residual of the completely reexamined "blenny" category, which also contained various misidentified ophidiiforms, and is now restricted to members of northern stichaeioid families and true blennioids (other than *Hypsoblennius* spp.) in the southern part of the pattern).

Labridae - all specimens originally identified to family were reexamined and assigned to genus (*Halichoeres*) or species (*Oxyjulis californica*, *Semicossyphus pulcher*); residuals are small, poorly preserved or unavailable specimens.

Pomacentridae - all specimens checked; now includes species other than *Chromis punctipinnis*, primarily in the south.

*Chromis punctipinnis* - all specimens taken south of line 120 checked.

*Mugil* spp. - specimen checked.

*Howella brodiei* - all specimens checked; originally included in Apogonidae; in this report we list *H. brodiei* in the family Apogonidae for convenience, recognizing that its systematic affinities are not resolved.

Carangidae - all specimens checked; tentative and sporadic identifications to genus or species (except *Trachurus symmetricus* and *Seriola lalandi*) were lumped to family.

*Seriola lalandi* - all specimens checked.

Gerreidae - tentative and sporadic identifications to genus were lumped to family.

Haemulidae - tentative and sporadic identifications to genus lumped to family.

*Girella nigricans* - all specimens checked.

*Medialuna californiensis* - all specimens checked.

*Caulolatilus princeps* - all specimens checked.

*Sciaenidae* - tentative and sporadic identifications to genus lumped to family.

*Scombridae* - all larvae identified to this family or constituent taxa (except *Scomber japonicus*) were reexamined and reassigned; residuals are small, poorly preserved or unavailable specimens.

*Nameidae* - tentative identifications to genus lumped to family.

*Pleuronectiformes* - all specimens of this category (originally called "flatfish") were examined and reidentified; residuals are small, poorly preserved or unavailable specimens.

*Bothidae* - all specimens examined and reassigned; most were assigned to various paralichthyid genera.

*Citharichthys* spp. - all larvae identified to species were lumped to genus except *C. stigmaeus*; category includes larvae of *Etropus* spp.

*Citharichthys stigmaeus* - includes larvae larger than ca. 4.5 mm; smaller larvae are in *Citharichthys* spp.

*Paralichthys* spp. - all specimens of this genus were examined and most were assigned to *P. californicus* or *Xystreurus liolepis*.

*Syacium ovale* - all specimens examined.

*Xystreurus liolepis* - originally misidentified as *Paralichthys californicus*; all specimens reidentified.

*Glyptocephalus zachirus* - all specimens examined.

*Hypsopsetta guttulata* - specimens were originally identified as *Pleuronichthys* spp.

*Microstomus pacificus* - all specimens examined.

*Pleuronichthys* spp. - all larvae of this genus and constituent species were examined and assigned to species; residuals are small, poorly preserved or unavailable specimens.

*Psettichthys melanostictus* - all specimens examined.

## COMPUTER ENTRY AND EDITING

Each taxon on the original identification sheets was given a 3-digit code based on the list of codes in Haight et al. (1979). Taxon codes and counts from these sheets were keypunched by cruise and station, along with pertinent station and tow data, and entered into the VAX 11/780 computer at the University of California, San Diego, Computing Center. After entries were completed for an entire year, print-out listings of taxa and counts on each station were compared with the original data sheets to eliminate keypunch errors. Next, data in the file were cross-checked with data on an existing file which contained: station and tow data; numbers of eggs of sardine, anchovy, and saury; numbers of larvae of sardine, anchovy, hake, jack mackerel, and Pacific mackerel; total number of fish eggs; and total number of fish larvae.

Discrepancies in ichthyoplankton data in these two files were corrected by inspecting original records from the sorting laboratory, the original ichthyoplankton identification sheets, and the samples themselves. Station and tow data discrepancies between the two files were corrected by reviewing ships' logs and deck tow sheets, original records from the sorting laboratory, cruise announcements, publications, header information on the ichthyoplankton identification sheets, and station plots generated for each cruise. Eventually all station and tow data were checked by comparing these sources.

The corrected ichthyoplankton data base was then examined statistically and outliers were found and checked as above. Distributional plots were then prepared for each taxon and these were checked by reviewing the data sources mentioned above and by examining archived specimens. A listing of each taxon by station (Table 4) was produced, which became the primary document for subsequent checks. Misidentifications found in geographic outlier checks and other misidentifications and data problems discovered in the course of examining archived samples resulted in several iterations of Table 4. Finally, totals in Table 4 were checked against annual summaries of incidence and abundance (Tables 2 and 3). Ecological analyses of the data were conducted concurrently with editing procedures and provided cross-checks that allowed correction of errors.

## SPECIES SUMMARY

Larvae of northern anchovy (*Engraulis mordax*) represented 57% of all fish larvae taken on CalCOFI cruises during 1965 and numbered eight times as many as the gonostomatid *Vinciguerria lucetia*, the next most abundant species with 7% of the total larvae (Table 2, 3). Northern anchovy also ranked first in incidence; *V. lucetia* ranked 3rd. Rockfish larvae, a composite of about 70 species of the genus *Sebastes*, ranked 3rd in number (4.4% of the total larvae) and 4th in incidence. Pacific hake, *Merluccius productus*, with 3.9% of the total larvae, ranked 4th

in number and 8th in incidence. The lanternfish *Triphoturus mexicanus* ranked 5th in number and 2nd in incidence and the deepsea smelt *Leuroglossus stilbius* ranked 6th in number and 7th in incidence; each contributed 3.2% to the total number of larvae. The Pacific sardine, *Sardinops sagax*, ranked 7th with 3.0% of the total number, but ranked 27th in incidence, an indication of several large samples. The sanddab genus *Citharichthys* ranked 8th in number (2.6%) and 5th in incidence. The remaining top 10 taxa were the lanternfish *Stenobrachius leucopsarus* (2.2%) and the jack mackerel, *Trachurus symmetricus* (1.5%). These two species ranked 10th and 17th, respectively, in incidence. The 10 taxa contributed 88% to the total larvae collected in 1965. The remaining 12% was distributed among 149 taxa, plus the unidentified and disintegrated categories. The top 10 taxa included 4 midwater species, 3 coastal pelagic species, and 3 coastal demersal taxa.

#### EXPLANATION OF TABLES

Table 1 - This table lists by cruise the pertinent station and tow data for 1965, the volume of water filtered and standard haul factor for each tow, the percent of sample sorted, and the total numbers of fish eggs and larvae. CalCOFI cruises are designated by four digits; the first two indicate the year and the second two the month. Within each cruise the data are listed in order of increasing line and station number (southerly and seaward directions); the order of station occupancy is shown on the station charts (Figures 2-5). Stations are designated by two groups of digits; the first set indicates the line and decimal fraction and the second set indicates the station on the line. Time is listed as Pacific Standard Time at the start of each tow in 24-hour designation. Methods for determining tow depth, volume of water strained, standard haul factor, and percent sorted were described in the methods section. The values for total fish eggs and larvae represent raw counts (unadjusted for percent sorted or standard haul factor). Ship codes are as follows: BD, *Black Douglas*; AX, *Alexander Agassiz*.

Table 2 - This table lists pooled occurrences of all larval fish taxa taken during 1965 in ranked order.

Table 3 - This table lists pooled counts of all larval fish taxa taken during 1965 in ranked order. Numbers are adjusted for percent sorted and standard haul factors.

Table 4 - This table gives numbers of fish larvae for each taxon, listed by station and calendar month in which the tow was taken. Counts are adjusted for percent of sample sorted and standard haul factor. Average values are given for stations occupied more than once during a month. See Table 1 for station and tow data and Table

6 for listing of stations with multiple occupancies during a month. Multiple occupancies occurred when a station was occupied more than once during a calendar month. The orders are listed in "phylogenetic" sequence modified from Nelson (1984). Subtaxa within each order are listed alphabetically. Page numbers for each taxon are given in the index at the end of the report.

Table 5 - This table is a summary of pooled occurrences of all larval fish taxa taken on CalCOFI surveys from 1961 to 1969. Taxa are listed in the same order as in Table 4.

Table 6 - List of stations with multiple occupancies in one month during 1965.

#### ACKNOWLEDGMENTS

Elaine Sandknop and the senior author originally identified larvae from CalCOFI cruises of 1965. Ronald Whyte coded each larval fish taxon or type and Rita Ford entered them into the computer. Debby Snow efficiently assisted in all aspects of data editing and retrieval. Cindy Meyer and James Ryan provided programming assistance. Dorothy Roll designed the CalCOFI data acquisition system and provided data processing support. Ken Raymond, Roy Allen, and Henry Orr helped with graphics and production of the report. Lorraine Prescott and Diane Forsythe prepared the manuscript for printing. Paul Smith determined statistical outliers, provided assistance during geographical outlier checks and offered helpful suggestions throughout the project. Izadore Barrett, Director of the Southwest Fisheries Center and Reuben Lasker, Chief, Coastal Fisheries Resources Division, SWFC, provided the support critical to the completion of the project. James Thrailkill planned CalCOFI surveys and supervised cruises, data handling, and plankton sorting from 1949 to 1986 and is largely responsible for the high quality of these operations. Without the vision and direction of Elbert Ahlstrom and Elton Sette and the dedicated efforts of the many people who collected, processed, and analyzed the samples, this data base would not exist.

## LITERATURE CITED

- Ahlstrom, E. H. 1948. A record of pilchard eggs and larvae collected during surveys made in 1939 to 1941. U.S. Fish Wildl. Serv. SSRF 54, 82 p.
- Ahlstrom, E. H. 1953. Pilchard eggs and larvae and other fish larvae, Pacific Coast - 1951. U.S. Fish Wildl. Serv. SSRF 102, 55 p.
- Ahlstrom, E. H. 1969. Distributional atlas of fish larvae in the California Current region: jack mackerel, *Trachurus symmetricus*, and Pacific hake, *Merluccius productus*, 1951 through 1966. CalCOFI Atlas No. 11:xi + 187 p.
- Ahlstrom, E. H., H. G. Moser, and E. M. Sandknop. 1978. Distributional atlas of fish larvae in the California Current region: rockfishes, *Sebastodes* spp., 1950 through 1975. CalCOFI Atlas No. 26:xxi + 178 p.
- Ambrose, D. A., R. L. Charter, H. G. Moser, and C. R. Santos Methot. 1987a. Ichthyoplankton and station data for California Cooperative Oceanic Fisheries Investigations survey cruises in 1951. U.S. Dep. Commer., NOAA Tech. Memo., NMFS, SWFC, No. 79, 196 p.
- Ambrose, D. A., R. L. Charter, H. G. Moser, and C. R. Santos Methot. 1987b. Ichthyoplankton and station data for California Cooperative Oceanic Fisheries Investigations survey cruises in 1955. U.S. Dep. Commer., NOAA Tech. Memo., NMFS, SWFC, No. 83, 185 p.
- Ambrose, D. A., R. L. Charter, H. G. Moser, and C. R. Santos Methot. 1987c. Ichthyoplankton and station data for California Cooperative Oceanic Fisheries Investigations survey cruises in 1960. U.S. Dep. Commer., NOAA Tech. Memo., NMFS, SWFC, No. 88, 253 p.
- Ambrose, D.A., R. L. Charter, H. G. Moser, and B. S. Earhart. 1988. Ichthyoplankton and station data for California Cooperative Oceanic Fisheries Investigations survey cruises in 1963. U.S. Dep. Commer., NOAA Tech. Memo., NMFS, SWFC, No. 94, 209 p.
- Haight, C. A., H. G. Moser, and P. E. Smith. 1979. Data entry programs: CalCOFI. II. Fish eggs and larvae identification sheet. National Marine Fisheries Service, Southwest Fisheries Center, La Jolla, Admin. Rept. No. LJ-79-25.
- Kramer, D. 1970. Distributional atlas of fish eggs and larvae in the California current region: Pacific sardine, *Sardinops caerulea* (Girard), 1951 through 1966. CalCOFI Atlas No. 12:vi + 277 p.

Kramer, D. and E. H. Ahlstrom. 1968. Distributional atlas of fish larvae in the California Current region: northern anchovy, *Engraulis mordax* Girard, 1951 through 1965. CalCOFI Atlas No. 9:xi + 269 p.

Kramer, D., M. Kalin, E. G. Stevens, J. R. Threlkild, and J. R. Zweifel. 1972. Collecting and processing data on fish eggs and larvae in the California Current Region. NOAA Tech. Rep. NMFS Circ. 370, 38 p.

Nelson, J. S. 1984. Fishes of the world. John Wiley and Sons, N.Y., 523 p.

Powles, H. and D. F. Markle. 1984. Identification of larvae, p. 31-33. In: Ontogeny and systematics of fishes. H. G. Moser, W. J. Richards, D. M. Cohen, M. P. Fahay, A. W. Kendall, Jr., and S. L. Richardson (eds.). Spec. Publ. No. 1. Amer. Soc. Ichthyol. Herpetol., 760 p.

Sandknop, E. M., R. L. Charter, H. G. Moser, and J. D. Ryan. 1987a. Ichthyoplankton and station data for California Cooperative Oceanic Fisheries Investigations survey cruises in 1952. U.S. Dep. Commer., NOAA Tech. Memo., NMFS, SWFC, No. 80, 207 p.

Sandknop, E. M., R. L. Charter, H. G. Moser, and J. D. Ryan. 1987b. Ichthyoplankton and station data for California Cooperative Oceanic Fisheries Investigations survey cruises in 1958. U.S. Dep. Commer. NOAA Tech. Memo., NMFS, SWFC, No. 86, 248 p.

Sandknop, E. M., R. L. Charter, H. G. Moser, C. A. Meyer, and A. E. Hays. 1988a. Ichthyoplankton and station data for California Cooperative Oceanic Fisheries Investigations survey cruises in 1961. U.S. Dep. Commer., NOAA Tech. Memo., NMFS, SWFC, No. 92, 167 p.

Sandknop, E. M., R. L. Charter, H. G. Moser, C. A. Meyer, and A. E. Hays. 1988b. Ichthyoplankton and station data for California Cooperative Oceanic Fisheries Investigations survey cruises in 1964. U.S. Dep. Commer. NOAA Tech. Memo., NMFS, SWFC, No. 95, 222 p.

Smith, P. E. 1971. Distributional atlas of zooplankton volume in the California Current region, 1951 through 1966. CalCOFI Atlas No. 13:xvi + 144 p.

Smith, P. E. and S. L. Richardson. 1977. Standard techniques for pelagic fish egg and larva surveys. FAO Fish. Tech. Pap. No. 175, 100 p.

Staff, South Pacific Fishery Investigations. 1953. Zooplankton volumes off the Pacific Coast, 1952. U.S. Fish Wildl. Serv. SSRF 100, 41 p.

Stevens, E. G., R. L. Charter, H. G. Moser, and M. S. Busby.  
1987a. Ichthyoplankton and station data for California  
Cooperative Oceanic Fisheries Investigations survey cruises  
in 1953. U.S. Dep. Commer., NOAA Tech. Memo., NMFS, SWFC,  
No. 81, 186 p.

Stevens, E. G., R. L. Charter, H. G. Moser, and M. S. Busby.  
1987b. Ichthyoplankton and station data for California  
Cooperative Oceanic Fisheries Investigations survey cruises  
in 1956. U.S. Dep. Commer., NOAA Tech. Memo., NMFS, SWFC,  
No. 84, 189 p.

Stevens, E. G., R. L. Charter, H. G. Moser, and M. S. Busby.  
1987c. Ichthyoplankton and station data for California  
Cooperative Oceanic Fisheries Investigations survey cruises  
in 1959. U.S. Dep. Commer., NOAA Tech. Memo., NMFS, SWFC,  
No. 87, 273 p.

Sumida, B. Y., R. L. Charter, H. G. Moser, and D. L. Snow.  
1987a. Ichthyoplankton and station data for California  
Cooperative Oceanic Fisheries Investigations survey cruises  
in 1954. U.S. Dep. Commer., NOAA Tech. Memo., NMFS, SWFC,  
No. 82, 207 p.

Sumida, B. Y., R. L. Charter, H. G. Moser, and D. L. Snow.  
1987b. Ichthyoplankton and station data for California  
Cooperative Oceanic Fisheries Investigations survey cruises  
in 1957. U.S. Dep. Commer., NOAA Tech. Memo., NMFS, SWFC,  
No. 85, 225 p.

Sumida, B. Y., R. L. Charter, H. G. Moser, and D. L. Snow.  
1988. Ichthyoplankton and station data for California  
Cooperative Oceanic Fisheries Investigations survey cruises  
in 1962. U.S. Dep. Commer., NOAA Tech. Memo., NMFS, SWFC,  
No. 93, 179 p.

University of California, Scripps Institution of Oceanography.  
1965. Data report: physical and chemical data, CalCOFI  
Cruise 6501. SIO Ref. 66-4.

University of California, Scripps Institution of Oceanography.  
1967. Data report: physical and chemical data, CalCOFI  
Cruise 6504; 6507, 6509. SIO Ref. 67-16; 67-17.






Figure 1. Composite arrangement of diagrammatic charts showing areas sampled on each CalCOFI cruise during 1965.




Figure 2. Station pattern for CalCOFI Cruise 6501 showing tracks for each vessel. Stations with plankton tows are indicated by a dot; circles designate hydrographic stations and diamonds signify STD recordings. Figures 2-5 modified from charts in Univ of Calif., SIO (1965, 1967) to include only those stations listed in Table 1 of this report; see Table 1 for inshore stations not shown on charts.




Figure 3. Station pattern for CalCOFI Cruise 6504. Symbols as in Figure 2.




Figure 4. Station pattern for CalCOFI Cruise 6507. Symbols as in Figure 2; additional symbol, an enlarged diamond and circle, indicates deep hydrographic station with STD recording.




Figure 5. Station pattern for CalCOFI Cruise 6509. Symbols as in Figure 2.




Figure 6. The basic station plan for CalCOFI cruises from 1950 to the present.

TABLE 1. Station and plankton tow data for CalCOFI cruises in 1965. Counts for fish eggs and larvae are not adjusted for standard haul factor or percent of sample sorted.

CalCOFI Cruise 6501

Line	Station	Lat. (N) deg. min.	Long. (W) deg. min.	Ship Code	Tow Date yr. mo. day	Time (PST)	Tow Depth (m)	Water Strained (cu. m)	Stand- ard Haul Factor	Percent Sorted	Total Larvae	Total Eggs
60.0	49.5	37 59.0	122 49.5	AX	65 01 10	1314	16	192	0.86	100.0	49	35
60.0	50.0	37 57.4	122 53.2	AX	65 01 10	1344	28	216	1.30	100.0	35	248
60.0	51.0	37 55.4	122 57.3	AX	65 01 10	1454	40	249	1.62	100.0	177	932
60.0	52.0	37 53.9	123 01.8	AX	65 01 10	1604	52	233	2.25	100.0	83	654
60.0	55.0	37 47.5	123 15.0	AX	65 01 10	1814	113	366	3.09	100.0	335	174
60.0	60.0	37 37.5	123 35.4	AX	65 01 10	2136	135	510	2.65	100.0	553	333
60.0	70.0	36 17.0	124 20.5	AX	65 01 11	0341	148	457	3.24	100.0	32	24
60.0	80.0	36 56.3	125 02.3	AX	65 01 11	0901	130	512	2.54	100.0	37	16
60.0	90.0	36 37.0	125 46.5	AX	65 01 11	1456	144	430	3.35	100.0	14	44
60.0	100.0	36 17.0	126 30.0	AX	65 01 11	2016	141	451	3.12	100.0	11	10
60.0	120.0	35 39.0	127 54.0	AX	65 01 12	0621	149	432	2.46	100.0	7	21
60.0	140.0	34 56.2	129 18.9	AX	65 01 12	1436	148	418	3.54	100.0	7	11
60.0	160.0	34 17.8	130 40.5	AX	65 01 12	2341	148	459	3.23	100.0	23	6
60.0	180.0	33 37.8	132 05.0	AX	65 01 13	0836	141	479	2.95	100.0	15	4
60.0	200.0	32 56.0	133 27.0	AX	65 01 13	1756	143	459	3.11	100.0	38	20
63.0	49.7	37 24.0	122 26.4	AX	65 01 09	0830	13	223	0.58	100.0	149	553
63.0	50.0	37 23.6	122 27.8	AX	65 01 09	0800	19	230	0.82	100.0	366	1434
63.0	51.0	37 22.0	122 32.5	AX	65 01 09	0704	34	128	2.63	100.0	110	2333
63.0	52.0	37 19.3	122 36.6	AX	65 01 09	0613	69	238	2.90	100.0	366	2374
63.0	55.0	37 12.7	122 49.6	AX	65 01 09	0331	144	455	3.11	100.0	589	34
63.0	60.0	37 01.2	123 17.2	AX	65 01 09	0006	144	441	3.26	100.0	44	20
67.0	47.3	36 54.4	121 53.5	AX	65 01 08	0804	14	211	0.68	100.0	30	744
67.0	48.0	36 52.8	121 55.7	AX	65 01 08	0844	28	230	1.23	100.0	5	127
67.0	49.0	36 50.8	122 00.4	AX	65 01 08	0953	74	230	3.23	100.0	95	883
67.0	50.0	36 49.0	122 04.6	AX	65 01 08	1118	64	271	2.35	100.0	302	227
67.0	55.0	36 38.9	122 25.9	AX	65 01 08	1506	143	432	3.32	100.0	182	94
67.0	60.0	36 29.0	122 47.5	AX	65 01 08	1806	141	469	3.01	100.0	99	101
70.0	50.0	36 11.4	121 43.8	AX	65 01 17	1603	85	289	2.93	100.0	587	17
70.0	51.0	36 10.4	121 45.7	AX	65 01 17	1426	139	454	3.07	100.0	248	12
70.0	52.0	36 08.7	121 50.0	AX	65 01 17	1326	141	446	3.15	100.0	211	83
70.0	53.0	36 06.5	121 54.0	AX	65 01 17	1201	136	463	2.95	100.0	145	715
70.0	55.0	36 02.7	122 02.0	AX	65 01 17	1016	136	470	2.90	100.0	148	214
70.0	60.0	35 53.0	122 24.1	AX	65 01 17	0701	140	490	2.86	100.0	31	36
70.0	70.0	35 32.0	123 05.3	AX	65 01 17	0141	141	521	2.71	100.0	166	22
70.0	80.0	35 12.8	123 47.0	AX	65 01 16	2011	139	498	2.78	100.0	14	3
70.0	90.0	34 53.8	124 28.1	AX	65 01 16	1526	136	473	2.87	100.0	7	2
70.0	100.0	34 33.2	125 09.3	AX	65 01 16	1001	141	494	2.85	100.0	3	4
70.0	120.0	33 54.0	126 36.0	AX	65 01 16	0031	142	478	2.98	100.0	45	9
70.0	160.0	32 34.8	129 21.6	AX	65 01 15	0526	140	471	2.97	100.0	45	53
70.0	180.0	31 54.0	130 43.0	AX	65 01 14	1651	140	471	2.98	100.0	12	12
70.0	200.0	31 10.5	132 12.0	AX	65 01 14	0736	141	462	3.06	100.0	34	27
73.0	49.4	35 38.3	121 15.0	AX	65 01 17	2010	7	230	0.31	100.0	142	1926
73.0	50.0	35 37.2	121 17.1	AX	65 01 17	2053	63	273	2.32	100.0	567	144

TABLE 1. (cont.)

Line	Station	Lat. (N) deg. min.	Long. (W) deg. min.	Ship Code	Tow Date yr. mo. day	Time (PST)	Tow Depth (m)	Water Strained (cu. m)	Vol. Stand- ard Haul Factor	Percent Sorted	Total Larvae	Total Eggs				
											Tow Time (PST)	Depth (m)	Water Strained (cu. m)	Vol.	Stand- ard Haul Factor	Percent Sorted
73.0	51.0	35 35.2	121 21.0	AX	65 01 17	2126	142	458	3.09	100.0	322	260				
73.0	53.0	35 31.7	121 28.6	AX	65 01 17	2336	133	448	2.73	100.0	503	439				
73.0	60.0	35 17.8	121 57.4	AX	65 01 18	0346	139	497	2.79	100.0	163	114				
77.0	47.6	35 08.4	120 43.9	AX	65 01 18	2235	7	249	0.28	100.0	269	4912				
77.0	48.0	35 08.8	120 42.2	AX	65 01 18	2320	7	249	0.29	100.0	526	3700				
77.0	51.0	35 01.8	120 56.3	AX	65 01 19	0231	141	475	2.97	100.0	375	76				
77.0	55.0	34 54.2	121 13.2	AX	65 01 18	1236	141	488	2.88	100.0	65	350				
77.0	57.0	34 50.2	121 21.2	AX	65 01 18	0926	140	469	2.99	100.0	80	150				
80.0	50.2	34 27.7	120 29.0	AX	65 01 19	0705	7	192	0.36	100.0	292	280				
80.0	51.0	34 26.1	120 32.5	AX	65 01 19	0836	141	447	3.16	100.0	382	521				
80.0	52.0	34 24.4	120 36.6	AX	65 01 19	1016	140	427	3.29	100.0	297	417				
80.0	55.0	34 18.8	120 48.5	AX	65 01 19	1321	144	461	3.11	100.0	256	1151				
80.0	60.0	34 08.7	121 09.0	AX	65 01 19	1656	133	516	2.57	100.0	80	73				
80.0	65.0	33 59.0	121 29.9	AX	65 01 19	2006	141	452	3.12	100.0	45	55				
80.0	70.0	33 48.1	121 51.0	AX	65 01 19	2301	139	488	2.84	100.0	13	8				
80.0	80.0	33 30.0	122 32.9	AX	65 01 20	0526	140	467	2.99	100.0	19	32				
80.0	90.0	33 08.1	123 13.0	AX	65 01 20	1001	140	486	2.88	100.0	10	5				
80.0	100.0	32 49.9	123 53.0	AX	65 01 20	1541	138	454	3.05	100.0	5	8				
80.0	120.0	32 09.0	125 15.0	AX	65 01 21	0111	141	478	2.95	100.0	81	3				
80.0	140.0	31 29.2	126 36.2	AX	65 01 21	1156	136	498	2.73	100.0	12	15				
80.0	160.0	30 47.9	127 56.0	AX	65 01 21	1941	140	481	2.92	100.0	131	16				
80.0	180.0	30 08.1	129 16.9	AX	65 01 22	0501	138	505	2.74	100.0	31	21				
80.0	200.0	29 29.5	130 36.0	AX	65 01 22	1426	139	492	2.83	100.0	21	6				
82.0	47.0	34 15.0	119 59.0	BD	65 01 20	1831	142	497	2.86	100.0	179	1813				
83.0	39.0	34 15.5	119 17.7	BD	65 01 20	0949	14	88	1.60	100.0	24	354				
83.0	40.0	34 13.7	119 22.0	BD	65 01 20	1114	22	113	1.91	100.0	169	251				
83.0	43.0	34 08.0	119 34.0	BD	65 01 20	1356	143	282	2.96	100.0	146	554				
83.0	51.0	33 52.0	120 07.6	BD	65 01 20	2222	102	432	2.36	100.0	53	275				
83.0	55.0	33 44.0	120 24.5	BD	65 01 21	1051	141	485	2.90	100.0	309	4499				
83.0	60.0	33 34.0	120 45.0	BD	65 01 21	1436	142	467	3.04	100.0	101	1921				
83.0	65.0	33 25.0	121 05.8	BD	65 01 21	1711	134	468	2.87	100.0	139	349				
83.0	70.0	33 15.2	121 26.3	BD	65 01 21	2046	141	441	3.19	100.0	48	161				
83.0	80.0	32 53.0	122 07.3	BD	65 01 22	0111	142	402	3.53	100.0	58	18				
83.0	90.0	32 31.5	122 46.0	BD	65 01 22	0736	142	467	3.03	100.0	23	13				
87.0	32.5	33 54.0	118 26.5	BD	65 01 24	0120	12	77	1.58	100.0	605	439				
87.0	33.0	33 54.2	118 29.4	BD	65 01 24	0030	44	221	1.99	100.0	2821	1500				
87.0	34.0	33 52.2	118 33.4	BD	65 01 23	2338	62	231	2.67	100.0	1004	875				
87.0	35.0	33 50.0	118 37.5	BD	65 01 23	2216	140	475	2.94	100.0	1669	1591				
87.0	40.0	33 40.0	118 58.0	BD	65 01 23	1851	138	484	2.85	100.0	1352	1128				
87.0	45.0	33 30.0	119 19.0	BD	65 01 23	1501	142	466	3.06	100.0	699	3111				
87.0	50.0	33 20.0	119 39.5	BD	65 01 23	1148	70	178	3.94	100.0	391	487				
87.0	55.0	33 11.2	120 01.3	BD	65 01 23	0841	140	469	2.98	100.0	954	2892				
87.0	60.0	33 00.0	120 21.5	BD	65 01 23	0531	139	463	3.00	100.0	47	1756				
87.0	65.0	32 49.5	120 41.5	BD	65 01 22	0211	145	469	3.08	100.0	12	1030				
87.0	70.0	32 39.5	121 02.0	BD	65 01 22	2311	142	460	3.10	100.0	39	5				

TABLE 1. (cont.)

## CalCOFI Cruise 6501

Line	Station	Lat. (N) deg. min.	Long. (W) deg. min.	Ship Code	Tow yr. mo. day	Date yr. mo. day	Tow Time (PST)	Depth (m)	Water vol. (cu. m)	Strained Haul (cu. m)	Percent Sorted	Total Larvae	Total Eggs
87.0	80.0	32 20.0	121 43.0	BD	65 01	22	1756	142	469	3.03	100.0	48	17
87.0	90.0	32 29.2	122 23.0	BD	65 01	22	1211	142	476	2.99	100.0	21	14
90.0	27.5	33 29.2	117 44.6	AX	65 01	27	1214	17	249	0.57	100.0	319	1537
90.0	28.0	33 28.5	117 46.6	AX	65 01	27	1111	137	490	2.79	100.0	586	957
90.0	32.0	33 21.1	118 01.6	AX	65 01	27	0816	136	505	2.70	100.0	741	2995
90.0	37.0	33 10.9	118 22.3	AX	65 01	27	0421	140	466	2.99	100.0	1373	3840
90.0	45.0	32 54.5	118 55.4	AX	65 01	26	2336	132	520	2.54	100.0	922	2191
90.0	53.0	32 39.1	119 28.0	AX	65 01	26	1831	143	444	3.23	100.0	277	3960
90.0	60.0	32 24.1	119 57.2	AX	65 01	26	1341	138	463	2.99	100.0	199	697
90.0	65.0	32 15.4	120 18.3	AX	65 01	26	0911	141	436	3.23	100.0	19	12
90.0	70.0	32 04.2	120 39.0	AX	65 01	26	0621	142	458	3.10	100.0	9	118
90.0	80.0	31 45.0	121 19.5	AX	65 01	25	2321	131	492	2.66	100.0	27	38
90.0	90.0	31 24.0	121 59.0	AX	65 01	25	1716	142	472	3.00	100.0	8	20
90.0	100.0	31 04.5	122 40.3	AX	65 01	25	1036	138	500	2.75	100.0	23	29
90.0	120.0	30 24.2	123 59.1	AX	65 01	25	0021	137	507	2.71	100.0	25	14
90.0	140.0	29 45.0	125 19.9	AX	65 01	24	0816	142	543	2.61	100.0	1	7
90.0	160.0	29 07.0	126 39.3	AX	65 01	23	2321	135	522	2.60	100.0	29	3
90.0	180.0	28 24.5	127 58.5	AX	65 01	23	1346	140	513	2.73	100.0	19	11
90.0	200.0	27 42.8	129 18.5	AX	65 01	23	0516	139	497	2.81	100.0	31	27
93.0	26.5	32 56.8	117 16.7	BD	65 01	24	1015	15	62	2.40	100.0	104	714
93.0	27.0	32 56.0	117 19.0	BD	65 01	24	1048	77	272	2.83	100.0	43	213
93.0	28.0	32 54.7	117 21.8	BD	65 01	25	1411	143	463	3.09	100.0	187	313
93.0	30.0	32 50.2	117 31.0	BD	65 01	25	1700	143	470	3.04	100.0	1220	3383
93.0	35.0	32 40.5	117 51.5	BD	65 01	25	2001	143	475	3.00	100.0	1355	2331
93.0	40.0	32 30.0	118 11.5	BD	65 01	26	0016	144	478	3.01	100.0	232	5052
93.0	45.0	32 21.5	118 34.0	BD	65 01	26	0326	143	472	3.04	100.0	742	3490
93.0	50.0	32 11.5	118 55.3	BD	65 01	26	0706	143	459	3.11	100.0	230	2793
93.0	55.0	32 00.0	119 13.5	BD	65 01	26	1001	139	453	3.07	100.0	147	1615
93.0	60.0	31 50.5	119 33.5	BD	65 01	26	1316	142	467	3.05	100.0	182	5950
93.0	65.0	31 42.5	119 55.2	BD	65 01	26	1631	143	454	3.15	100.0	40	262
93.0	70.0	31 31.0	120 14.0	BD	65 01	26	1946	142	484	2.94	100.0	31	70
93.0	80.0	31 15.0	120 53.5	BD	65 01	27	0151	141	469	3.01	100.0	29	202
93.0	90.0	31 00.0	121 31.5	BD	65 01	27	0656	142	446	3.19	100.0	20	53
93.0	96.5	32 19.4	117 03.8	BD	65 01	29	0230	11	91	1.18	100.0	868	267
96.5	28.7	32 17.5	117 04.7	BD	65 01	29	0149	35	108	1.87	100.0	1792	534
97.0	29.0	32 17.5	117 04.7	BD	65 01	29	0053	49	226	2.16	100.0	1404	275
97.0	30.0	32 16.0	117 07.0	BD	65 01	28	2306	140	449	3.12	100.0	1870	3142
97.0	32.0	32 12.0	117 15.2	BD	65 01	28	2051	140	457	3.07	100.0	1024	1569
97.0	35.0	32 06.5	117 28.5	BD	65 01	28	1811	141	449	3.15	100.0	1029	1697
97.0	40.0	31 58.0	117 51.0	BD	65 01	28	1416	142	468	3.04	100.0	2287	3041
97.0	45.0	31 46.0	118 08.5	BD	65 01	28	1126	141	449	3.15	100.0	1528	1674
97.0	50.0	31 32.1	118 27.2	BD	65 01	28	0741	141	472	3.00	100.0	216	5019
97.0	55.0	31 25.5	118 49.5	BL	65 01	28	0506	141	478	2.96	100.0	326	1294
97.0	60.0	31 16.2	119 10.0	BD	65 01	28	0151	142	462	3.07	100.0	695	450
97.0	65.0	31 06.8	119 30.8	BD	65 01	27	2311	140	462	3.07	100.0	100.0	450
97.0	70.0	30 56.3	119 50.8	BD	65 01	27						100.0	88

TABLE 1. (cont.)

## CalCOFI Cruise 6501

Line	Station	Lat. (N) deg. min.	Long. (W) deg. min.	Ship Code	Tow Date yr. mo. day	Time (PST)	Tow Depth (m)	Vol. Water (cu. m)	Stand- ard Strained Factor	Percent Sorted	Total Larvae	Total Eggs
97.0	80.0	30 36.0	120 30.5	BD	65 01 27	1741	141	422	3.35	100.0	12	23
97.0	90.0	30 18.0	121 08.3	BD	65 01 27	1236	140	461	3.05	100.0	14	34
100.0	29.0	31 42.2	116 43.5	AX	65 01 29	2232	104	277	3.75	100.0	636	658
100.0	30.0	31 40.5	116 47.0	AX	65 01 30	0031	139	495	2.81	100.0	361	1053
100.0	35.0	31 31.0	117 06.9	AX	65 01 30	0401	141	484	2.92	100.0	993	1164
100.0	40.0	31 20.8	117 27.0	AX	65 01 30	0806	140	477	2.94	100.0	500	329
100.0	45.0	31 12.4	117 48.6	AX	65 01 30	1026	137	442	3.11	100.0	524	1888
100.0	50.0	31 01.0	118 07.5	AX	65 01 30	1351	134	481	2.79	100.0	417	1331
100.0	55.0	30 50.2	118 29.5	AX	65 01 30	1511	139	460	3.03	100.0	649	1195
100.0	60.0	30 41.0	118 48.2	AX	65 01 30	2016	141	475	2.98	100.0	234	34
100.0	65.0	30 29.6	119 09.3	AX	65 01 30	2226	140	471	2.98	100.0	231	48
100.0	70.0	30 17.8	119 30.2	AX	65 01 31	0211	140	455	3.08	100.0	440	178
100.0	80.0	29 52.1	120 09.2	AX	65 01 31	0731	140	482	2.91	100.0	27	14
100.0	90.0	29 40.8	120 46.9	AX	65 01 31	2326	141	467	3.03	100.0	21	14
100.0	100.0	29 21.0	121 29.5	AX	65 02 02	0456	141	478	2.95	100.0	67	46
100.0	120.0	28 40.8	122 46.7	AX	65 02 02	1421	142	477	2.97	100.0	9	10
103.0	28.5	31 07.8	116 19.2	BD	65 01 30	2250	12	96	1.25	100.0	203	97
103.0	29.0	31 07.0	116 21.0	BD	65 01 30	2344	22	98	2.29	100.0	110	214
103.0	30.0	31 05.2	116 25.2	BD	65 01 31	0043	54	220	2.46	100.0	195	33
103.0	35.0	30 55.2	116 45.2	BD	65 01 31	0341	138	470	2.94	100.0	1297	525
103.0	40.0	30 46.3	117 04.5	BD	65 01 31	0616	141	481	2.92	100.0	259	280
103.0	45.0	30 36.0	117 24.0	BD	65 01 31	0906	141	479	2.95	100.0	23	12
103.0	50.0	30 26.5	117 45.0	BD	65 01 31	1141	142	486	2.92	100.0	12	8
103.0	55.0	30 17.0	118 05.0	BD	65 01 31	1456	141	491	2.87	100.0	109	24
103.0	60.0	30 07.0	118 25.0	BD	65 01 31	1736	141	464	3.04	100.0	162	109
103.0	65.0	29 57.0	118 44.3	BD	65 01 31	2026	137	482	2.83	100.0	354	76
103.0	70.0	29 47.2	119 04.0	BD	65 01 31	2256	142	471	3.01	100.0	285	155
103.0	80.0	29 26.5	119 40.0	BD	65 02 01	0346	141	470	3.00	100.0	39	8
103.0	90.0	29 04.3	120 16.0	BD	65 02 01	0831	143	468	3.04	100.0	30	25
105.0	140.0	27 09.3	123 26.1	BD	65 02 02	0026	141	508	2.78	100.0	48	16
107.0	30.0	30 30.0	116 03.3	BD	65 02 02	2250	13	84	1.50	100.0	104	222
107.0	31.0	30 27.8	116 07.0	BD	65 02 02	2159	33	155	2.15	100.0	331	66
107.0	32.0	30 25.8	116 11.0	BD	65 02 02	2036	143	464	3.09	100.0	956	2503
107.0	35.0	30 21.5	116 22.5	BD	65 02 02	1831	143	446	3.20	100.0	299	1212
107.0	40.0	30 11.5	116 42.5	BD	65 02 02	1531	144	489	2.94	100.0	253	210
107.0	45.0	30 01.5	117 02.0	BD	65 02 02	1246	142	477	2.99	100.0	2	13
107.0	50.0	29 50.1	117 22.9	BD	65 02 02	1006	142	486	2.93	100.0	17	26
107.0	55.0	29 41.5	117 42.5	BD	65 02 02	0736	142	482	2.95	100.0	7	17
107.0	60.0	29 32.0	118 01.5	BD	65 02 02	0436	142	467	3.03	100.0	185	107
107.0	65.0	29 22.0	118 21.0	BD	65 02 02	0136	142	483	2.94	100.0	14	85
107.0	70.0	29 12.0	118 41.0	BD	65 02 01	2231	144	452	3.18	100.0	105	236
107.0	80.0	28 53.2	119 20.5	BD	65 02 01	1736	142	490	2.90	100.0	32	63
107.0	90.0	28 32.0	119 59.0	BD	65 02 01	1251	141	476	2.96	100.0	41	11
110.0	32.0	29 52.1	115 47.8	AX	65 02 05	1434	14	212	0.67	100.0	330	532
110.0	35.0	29 46.3	116 00.0	AX	65 02 05	1201	141	496	2.83	100.0	374	796

TABLE 1. (cont.)

## CalCOFI Cruise 6501

Line	Station	Lat. (N) deg. min.	Long. (W) deg. min.	Ship Code	Tow Date yr. mo. day	Tow Time (PST)	Depth (m)	Vol. water (cu. m)	Stand- ard Haul Factor	Total Larvae	Total Eggs
110.0	40.0	29 37.4	116 26.1	AX	65 02	05	0746	139	502	2.77	1063
110.0	45.0	29 27.0	116 43.7	AX	65 02	05	0421	139	498	2.80	1000
110.0	50.0	29 16.0	116 59.3	AX	65 02	05	0206	141	504	2.80	1000
110.0	55.0	29 06.0	117 19.0	AX	65 02	04	2211	141	515	2.73	1000
110.0	60.0	28 57.0	117 38.9	AX	65 02	04	1951	141	516	2.72	1000
110.0	65.0	28 46.1	117 58.0	AX	65 02	04	1551	140	528	2.66	1000
110.0	70.0	28 36.0	118 18.0	AX	65 02	04	1321	140	500	2.80	1000
110.0	80.0	28 16.0	118 58.1	AX	65 02	04	0746	140	545	2.57	1000
110.0	90.0	27 55.8	119 36.7	AX	65 02	04	0221	141	535	2.64	1000
110.0	100.0	27 38.0	120 16.1	AX	65 02	03	1926	141	542	2.60	1000
110.0	120.0	26 55.6	121 32.5	AX	65 02	03	1016	139	540	2.58	1000
110.0	140.0	26 17.2	122 50.4	AX	65 02	03	0026	141	524	2.70	1000
110.0	160.0	25 37.2	124 02.8	AX	65 02	02	1526	140	498	2.80	1000
113.0	28.5	29 25.0	115 11.3	BD	65 02	03	0750	9	106	0.88	1000
113.0	29.0	29 24.2	115 13.2	BD	65 02	03	0829	20	92	2.15	1000
113.0	30.0	29 21.3	115 18.1	AX	65 02	05	1924	46	207	2.24	1000
113.0	30.0	29 21.5	115 17.8	BD	65 02	03	0924	48	196	2.47	1000
113.0	35.0	29 11.2	115 38.5	BD	65 02	03	1156	143	472	3.02	1000
113.0	40.0	29 01.0	115 59.8	BD	65 02	03	1441	141	484	2.93	1000
113.0	45.0	28 51.2	116 22.0	BD	65 02	03	1721	142	460	3.09	1000
113.0	50.0	28 41.8	116 39.5	BD	65 02	03	1951	142	469	3.04	1000
113.0	55.0	28 32.0	116 57.0	BD	65 02	03	2231	143	480	2.98	1000
113.0	60.0	28 22.5	117 18.5	BD	65 02	04	0116	141	480	2.94	1000
113.0	65.0	28 13.0	117 38.0	BD	65 02	04	0401	143	457	3.12	1000
113.0	70.0	28 03.0	117 59.0	BD	65 02	04	0646	136	498	2.73	1000
113.0	80.0	27 42.0	118 33.5	BD	65 02	04	1106	139	490	2.84	1000
113.0	90.0	27 22.5	119 12.5	BD	65 02	04	1536	143	505	2.83	1000
115.0	35.0	28 54.5	115 27.2	AX	65 02	05	2341	141	494	2.85	1000
115.0	24.8	28 58.5	114 36.3	BD	65 02	06	1204	9	92	0.99	1000
117.0	25.0	28 58.0	114 37.0	BD	65 02	06	1104	49	186	2.61	1000
117.0	26.0	28 56.0	114 41.5	BD	65 02	06	0958	70	251	2.80	1000
117.0	30.0	28 48.0	114 56.5	BD	65 02	06	0752	97	335	2.89	1000
117.0	35.0	28 38.0	115 16.0	BD	65 02	06	0506	141	449	3.14	1000
117.0	40.0	28 28.0	115 35.5	BD	65 02	05	2346	142	464	3.05	1000
117.0	45.0	28 19.5	116 00.2	BD	65 02	05	2056	140	474	2.96	1000
117.0	50.0	28 09.0	116 07.0	BD	65 02	05	1751	145	451	3.20	1000
117.0	55.0	27 57.5	116 35.5	BD	65 02	05	1451	144	468	3.07	1000
117.0	60.0	27 49.0	116 53.3	BD	65 02	05	1206	143	454	3.16	1000
117.0	65.0	27 40.0	117 12.2	BD	65 02	05	0921	145	467	3.11	1000
117.0	70.0	27 30.0	117 31.0	BD	65 02	05	146	146	474	3.08	1000
117.0	80.0	27 08.5	118 05.5	BD	65 02	04	0101	141	475	2.97	1000
117.0	90.0	26 48.5	118 50.0	BD	65 02	04	2006	144	489	2.94	1000
117.0	95.0	28 18.5	115 23.7	BD	65 02	06	0226	141	453	3.10	1000
117.0	100.0	28 19.0	114 52.9	AX	65 02	07	0343	71	268	2.66	1000
120.0	22.5	28 28.0	114 04.0	BD	65 02	06	1705	12	1.63	1.63	1000

TABLE 1. (cont.)

CalCOFI Cruise 6501

Line	Station	Lat. (N) deg. min.	Long. (W) deg. min.	Ship Code	Tow Date yr. mo. day	Time (PST)	Tow Depth (m)	Vol. Water Strained (cu. m)	Stand- ard Haul Factor	Percent Sorted	Total Larvae	Total Eggs
120.0	23.0	28 26.9	114 06.7	BD	65 02 06	1744	15	122	1.25	100.0	50	231
120.0	24.0	28 25.0	114 10.7	BD	65 02 06	1834	28	105	2.65	100.0	867	379
120.0	25.0	28 22.5	114 15.0	BD	65 02 06	1919	50	137	3.66	100.0	267	320
120.0	30.0	28 13.0	114 34.0	BD	65 02 07	0352	77	266	2.90	100.0	670	1501
120.0	35.0	28 03.0	114 54.0	BD	65 02 07	0633	79	258	3.05	100.0	250	67
120.0	40.0	27 56.5	115 14.0	BD	65 02 07	0909	37	139	2.65	100.0	117	2
120.0	45.0	27 43.1	115 33.2	AX	65 02 07	1026	140	485	2.88	100.0	21	114
120.0	50.0	27 33.8	115 52.4	AX	65 02 07	1411	144	470	3.07	100.0	37	72
120.0	55.0	27 23.0	116 11.4	AX	65 02 07	1636	138	492	2.80	100.0	3	5
120.0	60.0	27 13.8	116 30.9	AX	65 02 07	2031	140	486	2.88	100.0	24	48
120.0	65.0	27 02.8	116 49.8	AX	65 02 07	2311	141	504	2.79	100.0	117	15
120.0	70.0	26 52.7	117 09.5	AX	65 02 08	0256	141	482	2.92	100.0	20	23
120.0	80.0	26 32.4	117 49.0	AX	65 02 08	0841	140	498	2.81	100.0	8	20
120.0	90.0	26 12.8	118 25.8	AX	65 02 08	1441	142	487	2.91	100.0	10	21
120.0	100.0	25 52.1	119 06.5	AX	65 02 08	1946	142	474	2.99	100.0	62	38
120.0	110.0	25 11.6	120 24.3	AX	65 02 09	0526	141	487	2.90	100.0	93	347
120.0	120.0	27 26.2	120 36.0	BD	65 02 07	1504	35	131	2.67	100.0	25	98
123.0	36.0	27 24.0	114 40.0	BD	65 02 07	1543	66	230	2.86	100.0	66	12
123.0	37.0	27 14.0	114 59.2	AX	65 02 16	1856	144	437	3.29	100.0	928	427
123.0	42.0	27 14.0	114 59.2	BD	65 02 07	1841	141	478	2.96	100.0	197	729
123.0	42.0	27 08.0	115 11.5	BD	65 02 07	2021	143	490	2.91	100.0	274	83
123.0	45.0	27 08.0	115 11.5	BD	65 02 07	2321	142	485	2.93	100.0	36	20
123.0	50.0	26 57.8	115 30.5	BD	65 02 07	1543	66	230	2.86	100.0	9	32
123.0	55.0	26 48.0	115 51.8	BD	65 02 08	0141	143	481	2.98	100.0	8	12
123.0	60.0	26 39.0	116 12.8	BD	65 02 08	0441	142	486	2.92	100.0	3	35
123.0	65.0	26 31.0	116 28.8	BD	65 02 08	0646	142	486	2.93	100.0	11	5
123.0	70.0	26 21.2	116 49.5	BD	65 02 08	0946	142	503	2.82	100.0	46	173
123.0	80.0	25 58.5	117 24.0	BD	65 02 08	1421	141	482	2.92	100.0	538	61
123.0	85.0	27 24.2	114 31.8	BD	65 02 07	1420	14	63	2.22	100.0	128	269
123.0	35.6	26 58.3	114 00.6	BD	65 02 09	2140	12	82	1.49	100.0	458	1264
127.0	32.6	26 34.0	114 48.5	BD	65 02 09	2059	50	176	2.85	100.0	1796	2309
127.0	33.0	26 57.5	114 02.2	BD	65 02 09	2003	72	274	2.63	100.0	832	46
127.0	34.0	26 55.0	114 06.5	BD	65 02 09	1656	140	496	2.83	100.0	538	61
127.0	40.0	26 43.5	114 29.0	BD	65 02 09	1241	139	477	3.00	100.0	18	35
127.0	40.0	26 43.9	114 27.6	AX	65 02 16	1426	142	475	3.00	100.0	21	26
127.0	45.0	26 34.0	114 48.5	BD	65 02 09	1131	143	485	2.94	100.0	32	16
127.0	50.0	26 24.8	115 06.3	BD	65 02 09	0916	142	486	2.92	100.0	40	13
127.0	55.0	26 09.3	115 25.5	BD	65 02 09	0611	138	504	2.74	100.0	61	538
127.0	60.0	26 06.8	115 41.5	BD	65 02 09	0326	142	491	2.89	100.0	30	22
127.0	65.0	25 55.8	116 02.0	BD	65 02 09	0011	142	484	2.94	100.0	15	7
127.0	70.0	25 45.0	116 22.3	BD	65 02 09	1921	141	489	2.88	100.0	40	13
127.0	80.0	25 24.0	117 02.5	BD	65 02 08	0230	14	225	0.63	100.0	61	2478
130.0	0	25.5	113 11.5	AX	65 02 12	0200	21	221	0.96	100.0	76	3193
130.0	0	26.0	113 13.2	AX	65 02 12	0034	32	150	2.14	100.0	232	852
130.0	0	28.0	113 21.2	AX	65 02 11	2304	33	138	2.43	100.0	87	309
130.0	0	30.0	113 29.0	AX	65 02 11	1921	147	442	3.33	100.0	974	974

TABLE 1. (cont.)

CalCOFI Cruise 6501									
Line	Station	Lat. (N) deg. min.	Long. (W) deg. min.	Ship Code	Tow Date yr. mo. day	Tow Depth (PST) (m)	Vol. Water Strained (cu. m)	Stand- ard Haul Factor	Total Larvae
130.0	40.0	26 08.8	114 26.7	AX	65 02 11	1656	143	468	3.06
130.0	45.0	25 58.8	114 47.4	AX	65 02 11	1241	140	502	2.79
130.0	50.0	25 49.9	115 01.0	AX	65 02 11	1006	139	484	2.88
130.0	55.0	25 40.9	115 21.0	AX	65 02 11	0611	142	497	2.87
130.0	60.0	25 30.0	116 01.2	AX	65 02 10	0336	141	504	2.79
130.0	70.0	25 09.6	116 09.0	AX	65 02 10	2151	141	493	2.87
130.0	80.0	24 49.2	116 39.2	AX	65 02 10	1636	139	500	2.78
130.0	90.0	24 28.5	117 17.7	AX	65 02 10	1016	138	518	2.67
130.0	100.0	24 06.4	117 54.5	AX	65 02 10	0506	140	506	2.77
130.0	120.0	23 28.4	119 09.0	AX	65 02 09	1911	139	504	2.76
133.0	19.5	26 12.6	112 25.8	BD	65 02 10	0659	14	81	1.70
133.0	21.0	26 12.6	112 32.3	BD	65 02 10	0754	38	189	2.00
133.0	23.0	26 08.5	112 40.2	BD	65 02 10	0853	77	290	2.65
133.0	25.0	26 04.5	112 48.0	BD	65 02 10	1023	78	288	2.70
133.0	30.0	25 54.5	113 07.5	BD	65 02 10	1846	141	509	2.77
133.0	35.0	25 44.5	113 26.5	BD	65 02 10	2136	142	493	2.88
133.0	40.0	25 34.8	113 44.7	BD	65 02 10	2351	142	476	2.98
133.0	45.0	25 24.8	114 03.2	BD	65 02 11	0251	142	485	2.93
133.0	50.0	25 14.3	114 21.2	BD	65 02 11	0506	141	500	2.83
133.0	55.0	25 04.5	114 40.0	BD	65 02 11	0801	140	512	2.74
133.0	60.0	24 54.2	115 01.0	BD	65 02 11	1021	141	491	2.88
133.0	65.0	24 44.2	115 20.8	BD	65 02 11	1341	143	497	2.87
133.0	70.0	24 34.8	115 39.2	BD	65 02 11	1601	141	496	2.84
133.0	80.0	24 14.5	116 17.0	BD	65 02 11	2041	142	492	2.89
137.0	20.0	25 40.1	112 07.2	AX	65 02 12	1059	7	268	0.26
137.0	21.0	25 38.1	112 11.1	AX	65 02 12	1140	14	219	0.63
137.0	22.0	25 36.1	112 14.8	AX	65 02 12	1239	27	208	1.30
137.0	23.0	25 34.0	112 18.8	AX	65 02 12	1344	40	216	1.85
137.0	30.0	25 19.8	112 45.7	AX	65 02 12	1741	139	453	3.08
137.0	35.0	25 10.0	113 04.5	AX	65 02 12	2026	140	449	3.11
137.0	40.0	25 00.0	113 21.9	AX	65 02 12	2336	139	451	3.08
137.0	45.0	24 49.6	113 40.9	AX	65 02 13	0226	137	474	2.90
137.0	50.0	24 38.8	114 00.5	AX	65 02 13	0636	143	432	3.31
137.0	55.0	24 29.7	114 19.8	AX	65 02 13	0856	144	467	3.08
137.0	70.0	24 02.2	115 24.5	AX	65 02 13	1746	141	471	3.00
137.0	80.0	23 40.0	115 55.0	AX	65 02 13	2321	141	478	2.95
140.0	28.8	24 48.0	112 19.2	AX	65 02 15	0424	14	208	0.68
140.0	29.0	24 47.4	112 20.3	AX	65 02 15	0354	32	136	2.34
140.0	30.0	24 45.3	112 24.0	AX	65 02 15	0243	68	252	2.71
140.0	35.0	24 34.0	112 41.9	AX	65 02 14	2341	141	465	3.03
140.0	40.0	24 27.0	112 56.5	AX	65 02 14	2121	139	473	2.95
140.0	45.0	24 17.4	113 22.9	AX	65 02 14	1701	141	460	3.06
140.0	50.0	24 05.2	113 40.7	AX	65 02	1401	141	2.96	2.96

TABLE 1. (cont.)

CalCOFI Cruise 6504									
Line Station	Lat. (N) deg. min.	Long. (W) deg. min.	Ship Code	Tow Date yr. mo. day	Time (PST)	Tow Depth (m)	Water Strained (cu. m)	Stand- ard Haul Factor	Total Eggs
63.0 49.7	37 24.0	122 26.4	AX	65 04 15	0550	14	161	0.88	16 214
63.0 50.0	37 23.5	122 27.8	AX	65 04 15	0524	19	238	0.80	20 142
63.0 51.0	37 21.4	122 32.0	AX	65 04 15	0409	32	146	2.18	7 30
63.0 52.0	37 18.9	122 36.0	AX	65 04 15	0253	55	267	2.05	42 149
63.0 55.0	37 13.2	122 49.0	AX	65 04 15	0036	114	609	1.86	270 887
63.0 60.0	37 02.7	123 11.2	AX	65 04 14	2018	134	530	2.52	44 58
63.0 65.0	36 53.0	123 33.3	AX	65 04 14	1716	144	497	2.91	15 19
63.0 70.0	36 42.7	123 55.0	AX	65 04 14	1446	127	559	2.27	19 18
63.0 80.0	36 23.0	124 39.3	AX	65 04 14	0956	146	500	2.93	43 14
67.0 47.3	36 54.4	121 52.8	AX	65 04 13	0954	11	277	0.38	100.0 23
67.0 48.0	36 52.8	121 56.0	AX	65 04 13	1054	32	241	1.32	100.0 88
67.0 49.0	36 50.8	122 00.4	AX	65 04 13	1148	72	254	2.83	100.0 42
67.0 50.0	36 49.3	122 04.8	AX	65 04 13	1333	84	313	2.70	100.0 62
67.0 55.0	36 39.0	122 26.0	AX	65 04 13	1601	141	521	2.70	100.0 23
67.0 60.0	36 27.8	122 47.3	AX	65 04 13	1906	136	492	2.75	100.0 179
67.0 65.0	36 19.0	123 09.0	AX	65 04 13	2131	141	486	2.91	100.0 120
67.0 70.0	36 06.0	123 30.0	AX	65 04 14	0026	139	489	2.84	100.0 121
67.0 80.0	35 48.5	124 12.0	AX	65 04 14	0451	138	521	2.65	100.0 64
67.0 85.0	36 11.3	121 43.9	AX	65 04 13	0156	116	520	2.23	100.0 54
70.0 51.0	36 10.4	121 45.9	AX	65 04 13	0331	141	513	2.76	100.0 134
70.0 52.0	36 08.5	121 50.0	AX	65 04 12	2331	134	520	2.58	100.0 201
70.0 53.0	36 06.5	121 54.0	AX	65 04 12	2216	136	506	2.70	100.0 109
70.0 60.0	35 54.6	122 24.0	AX	65 04 12	1901	136	477	2.86	100.0 89
70.0 65.0	35 42.9	122 45.4	AX	65 04 12	1546	138	513	2.69	100.0 68
70.0 70.0	35 35.8	123 08.7	AX	65 04 12	1306	142	488	2.90	100.0 29
70.0 80.0	35 13.5	123 47.5	AX	65 04 12	0646	123	449	2.74	100.0 10
73.0 49.4	35 37.8	121 15.2	AX	65 04 11	0440	14	205	0.69	100.0 147
73.0 50.0	35 37.8	121 15.2	AX	65 04 11	0538	70	255	2.77	100.0 57
73.0 51.0	35 35.2	121 21.1	AX	65 04 11	0626	140	501	2.79	100.0 119
73.0 53.0	35 31.7	121 28.6	AX	65 04 11	0821	148	493	3.01	100.0 102
73.0 60.0	35 18.0	121 57.5	AX	65 04 11	1126	139	491	2.82	100.0 143
73.0 65.0	35 08.0	122 19.3	AX	65 04 11	1441	143	498	2.88	100.0 429
73.0 70.0	34 58.2	122 40.5	AX	65 04 11	1721	141	508	2.77	100.0 8
73.0 73.0	34 38.8	123 22.5	AX	65 04 11	2226	136	524	2.59	100.0 24
77.0 47.6	35 09.0	120 42.1	AX	65 04 10	1930	14	224	0.63	100.0 661
77.0 48.0	35 08.2	120 43.4	AX	65 04 10	1820	14	226	0.62	100.0 260
77.0 49.0	35 06.3	120 47.8	AX	65 04 10	1728	14	206	0.67	100.0 252
77.0 51.0	35 01.9	120 55.7	AX	65 04 10	1543	72	238	3.01	100.0 449
77.0 55.0	34 53.9	121 12.4	AX	65 04 10	1346	142	392	3.62	100.0 54
77.0 57.0	34 50.3	121 21.0	AX	65 04 10	1136	116	473	2.45	100.0 130
77.0 60.0	34 44.2	121 33.6	AX	65 04 10	0941	138	442	3.13	100.0 237
77.0 65.0	34 34.2	121 54.6	AX	65 04 10	0530	140	551	2.55	100.0 374
77.0 70.0	34 24.2	122 16.3	AX	65 04 10	0131	120	520	2.30	100.0 26
77.0 80.0	34 03.5	122 57.2	AX	65 04 09	2026	132	446	2.83	100.0 19
80.0 50.2	34 27.8	120 29.3	AX	65 04 08	1734	14	188	0.75	100.0 1743

TABLE 1. (cont.)

## CalCOFI Cruise 6504

Line	Station	Lat. (N) deg. min.	Long. (W) deg. min.	Ship Code	Tow Date yr. no. day	Tow Time (PST)	Vol. Water Strained (cu. m)	Tow Depth (m)	Stand- ard Haul Factor	Percent Sorted	Total Larvae	Total Eggs
80.0	51.0	34 26.3	120 32.5	AX	65 04	08	1823	71	202	3.51	100.0	898
80.0	52.0	34 24.3	120 36.5	AX	65 04	08	1951	138	413	3.33	100.0	2895
80.0	55.0	34 18.7	120 48.4	AX	65 04	08	2141	139	435	3.20	100.0	717
80.0	60.0	34 09.2	121 05.9	AX	65 04	09	0051	143	421	3.39	100.0	1041
80.0	65.0	33 59.4	121 30.2	AX	65 04	09	0356	142	420	3.37	100.0	4865
80.0	70.0	33 48.0	121 50.2	AX	65 04	09	0711	144	425	3.40	100.0	136
80.0	80.0	33 27.4	122 32.0	AX	65 04	09	1221	145	450	3.22	100.0	188
80.0	82.0	47.0	119 58.6	AX	65 04	08	1321	139	447	3.10	100.0	161
83.0	40.0	34 15.1	119 21.8	AX	65 04	07	2110	14	184	0.77	100.0	90
83.0	43.0	34 07.7	119 34.3	AX	65 04	07	1911	132	470	2.80	100.0	940
83.0	51.0	33 52.0	120 07.6	AX	65 04	07	1508	66	255	2.59	100.0	826
83.0	55.0	33 43.8	120 24.3	AX	65 04	07	1216	147	465	3.16	100.0	3980
83.0	60.0	33 33.8	120 45.0	AX	65 04	07	0941	138	460	3.00	100.0	695
83.0	65.0	33 24.2	121 05.5	AX	65 04	07	0611	137	470	2.91	100.0	722
83.0	70.0	33 15.1	121 25.3	AX	65 04	07	0346	142	452	3.15	100.0	292
83.0	80.0	32 55.7	122 07.0	AX	65 04	06	2256	136	474	2.87	100.0	143
87.0	32.5	33 55.5	118 27.4	AX	65 04	05	0104	14	193	0.73	100.0	101
87.0	33.0	33 54.1	118 29.4	AX	65 04	05	0159	32	123	2.63	100.0	478
87.0	34.0	33 52.3	118 33.4	AX	65 04	05	0244	32	130	2.45	100.0	1878
87.0	35.0	33 50.1	118 37.6	AX	65 04	05	0356	147	430	3.41	100.0	646
87.0	40.0	33 40.0	118 58.0	AX	65 04	05	0621	147	354	4.14	100.0	329
87.0	45.0	33 29.8	119 18.8	AX	65 04	05	0930	131	383	3.42	100.0	177
87.0	50.0	33 20.0	119 39.4	AX	65 04	05	1208	64	272	2.31	100.0	723
87.0	55.0	33 09.9	120 00.0	AX	65 04	05	1510	135	481	2.80	100.0	528
87.0	60.0	32 59.0	120 20.0	AX	65 04	05	1846	128	535	2.39	100.0	187
87.0	65.0	32 48.0	120 43.0	AX	65 04	05	2156	130	486	2.68	100.0	46
87.0	70.0	32 38.1	121 03.8	AX	65 04	05	026	141	441	3.21	100.0	947
87.0	80.0	32 19.5	121 43.5	AX	65 04	06	0506	138	458	3.01	100.0	251
87.0	90.0	32 00.0	122 24.0	AX	65 04	06	0900	136	480	2.82	100.0	46
87.0	100.0	31 40.0	123 04.0	AX	65 04	06	1351	139	445	3.12	100.0	107
90.0	27.5	33 29.5	117 44.7	AX	65 04	04	2014	14	218	0.65	100.0	3373
90.0	28.0	33 28.5	117 47.0	AX	65 04	04	1901	143	466	3.06	100.0	771
90.0	32.0	33 21.5	118 01.4	AX	65 04	04	1636	144	471	3.05	100.0	898
90.0	37.0	33 11.0	118 22.5	AX	65 04	04	1256	143	464	3.09	100.0	786
90.0	45.0	32 54.5	118 55.5	AX	65 04	04	0856	143	456	3.13	100.0	610
90.0	53.0	32 39.0	119 28.7	AX	65 04	04	0426	146	439	3.32	100.0	1221
90.0	60.0	32 25.7	119 55.5	AX	65 04	04	0056	146	438	3.34	100.0	642
90.0	65.0	32 16.0	120 17.0	AX	65 04	03	2101	141	479	2.94	100.0	459
90.0	70.0	32 05.0	120 38.8	AX	65 04	03	1836	147	472	3.11	100.0	321
90.0	80.0	31 46.9	121 15.5	AX	65 04	03	1301	146	452	3.22	100.0	416
90.0	90.0	31 26.0	121 59.0	AX	65 04	03	0711	144	450	3.20	100.0	56
90.0	100.0	31 05.8	122 37.0	AX	65 04	03	0151	143	458	3.12	100.0	442
93.0	27.0	32 56.0	117 18.8	AX	65 03	31	1258	74	187	3.96	100.0	867
93.0	28.0	32 54.8	117 22.2	AX	65 03	31	1515	145	437	3.32	100.0	343
93.0	30.0	32 50.3	117 31.0	AX	65 03	31	1750	137	478	2.87	100.0	347

TABLE 1. (cont.)

## CalCOFI Cruise 6504

Line	Station	Lat. (N) deg. min.	Long. (W) deg. min.	Ship Code	Tow yr. mo.	Date day	Time (PST)	Tow Depth (m)	Water Strained (cu. m)	Stand- ard Factor	Percent Sorted	Total Larvae	Total Eggs
93.0	35.0	32 41.0	117 52.3	AX	65 04	31	2156	145	500	2.91	100.0	250	407
93.0	40.0	32 30.4	118 11.3	AX	65 04	01	0156	148	449	3.29	100.0	266	346
93.0	45.0	32 21.5	118 33.0	AX	65 04	01	0550	148	463	3.19	100.0	62	297
93.0	50.0	32 09.5	118 53.0	AX	65 04	01	1120	135	392	3.45	100.0	135	121
93.0	55.0	32 00.0	119 13.0	AX	65 04	01	1606	146	352	4.16	100.0	253	544
93.0	60.0	31 48.3	119 33.0	AX	65 04	01	1946	153	454	3.38	100.0	714	274
93.0	65.0	31 40.0	119 54.0	AX	65 04	01	2301	139	467	2.98	100.0	739	477
93.0	70.0	31 30.0	120 14.2	AX	65 04	02	0206	146	439	3.33	100.0	2094	2602
93.0	80.0	31 10.2	120 55.0	AX	65 04	02	0811	146	461	3.17	100.0	258	470
93.0	90.0	30 48.2	121 36.2	AX	65 04	02	1441	148	441	3.36	100.0	110	116
93.0	100.0	30 31.1	122 13.0	AX	65 04	02	1956	138	487	2.83	100.0	113	115
96.8	28.7	32 19.4	117 03.8	BD	65 03	30	1820	5	140	0.37	100.0	45	590
97.0	29.0	32 17.5	117 04.7	BD	65 03	30	1928	49	202	2.42	100.0	157	3060
97.0	30.0	32 16.0	117 07.0	BD	65 03	30	1953	60	236	2.53	100.0	250	2641
97.0	32.0	32 12.0	117 15.2	BD	65 03	30	2110	140	450	3.10	100.0	997	8885
97.0	35.0	32 04.8	117 28.0	BD	65 03	30	2330	142	452	3.15	100.0	452	336
97.0	40.0	31 55.0	117 50.0	BD	65 03	31	0215	143	448	3.19	100.0	99	109
97.0	45.0	31 46.0	118 10.0	BD	65 03	31	0525	143	415	3.45	100.0	262	179
97.0	50.0	31 36.0	118 30.5	BD	65 03	31	0906	145	450	3.22	100.0	180	198
97.0	65.0	31 05.0	119 31.0	BD	65 03	31	1906	145	420	3.45	100.0	439	600
97.0	75.0	30 42.5	120 08.5	BD	65 04	01	0135	128	410	3.11	100.0	891	2711
97.0	85.0	30 20.0	120 46.5	BD	65 04	01	0846	139	389	3.58	100.0	208	246
97.0	95.0	30 06.5	121 32.5	BD	65 04	01	1526	139	382	3.63	100.0	31	282
97.0	100.0	29 58.0	121 56.5	BD	65 04	01	1826	140	400	3.50	100.0	17	124
97.0	129.0	31 42.2	116 43.4	BD	65 04	03	1242	117	403	2.91	100.0	149	216
100.0	30.0	31 40.8	116 47.2	BD	65 04	03	1146	138	432	3.21	100.0	237	148
100.0	35.0	31 30.5	117 07.0	BD	65 04	03	0841	139	442	3.15	100.0	50	154
100.0	40.0	31 19.0	117 24.3	BD	65 04	03	0616	141	437	3.22	100.0	33	33
100.0	45.0	31 10.0	117 46.5	BD	65 04	03	0306	116	512	2.27	100.0	78	130
100.0	50.0	31 00.5	118 08.5	BD	65 04	03	0036	137	455	3.01	100.0	246	43
100.0	55.0	30 52.5	118 28.0	BD	65 04	02	2121	140	462	3.02	100.0	179	98
100.0	60.0	30 44.0	118 47.0	BD	65 04	02	1906	142	424	3.35	100.0	116	116
100.0	65.0	30 32.5	119 07.0	BD	65 04	02	1556	139	436	3.19	100.0	448	433
100.0	70.0	30 20.0	119 26.5	BD	65 04	02	1341	142	426	3.34	100.0	721	372
100.0	75.0	30 07.8	119 46.0	BD	65 04	02	1016	142	422	3.37	100.0	240	336
100.0	80.0	29 59.5	120 04.5	BD	65 04	02	0741	142	426	3.34	100.0	53	181
100.0	85.0	29 50.0	120 26.7	BD	65 04	02	0431	144	416	3.45	100.0	259	633
100.0	90.0	29 41.0	120 47.0	BD	65 04	02	0143	141	427	3.31	100.0	196	141
103.0	28.5	31 07.8	116 19.2	BD	65 04	03	1806	14	76	1.81	100.0	55	513
103.0	29.0	31 07.0	116 21.0	BD	65 04	03	1834	16	102	1.59	100.0	113	157
103.0	30.0	31 06.0	116 24.5	BD	65 04	03	1933	62	204	3.06	100.0	327	129
103.0	35.0	30 56.0	116 45.0	BD	65 04	03	2156	140	450	3.11	100.0	161	242
103.0	40.0	30 46.0	117 04.5	BD	65 04	04	0110	140	457	3.06	100.0	80	73
103.0	45.0	30 36.0	117 24.0	BD	65 04	04	0336	139	459	3.03	100.0	63	556
103.0	50.0	30 25.0	117 43.5	BD	65 04	04	0645	140	426	3.29	100.0	51	51

TABLE 1. (cont.)

## CalCOFI Cruise 6504

Line	Station	Lat. (N) deg. min.	Long. (W) deg. min.	Ship Code	Tow Date yr. mo. day	Tow Time (PST)	Tow Depth (m)	Vol. Water (cu. m)	Strained Depth (m)	Haul Factor	Percent Sorted	Total Larvae	Total Eggs
103.0	55.0	30 15.2	118 03.5	BD	65 04	04	0911	143	428	3.33	100.0	23	23
103.0	60.0	30 07.0	118 21.0	BD	65 04	04	1221	142	424	3.35	100.0	10	245
103.0	65.0	29 58.0	118 40.0	BD	65 04	04	1506	142	449	3.15	100.0	38	113
103.0	70.0	29 49.2	119 05.5	BD	65 04	04	1844	143	459	3.11	100.0	114	128
103.0	75.0	29 39.0	119 25.4	BD	65 04	04	2126	141	442	3.19	100.0	121	438
103.0	80.0	29 27.5	119 44.0	BD	65 04	05	0036	139	469	2.97	100.0	107	328
103.0	85.0	29 17.2	120 05.0	BD	65 04	05	0306	141	484	2.90	100.0	161	210
103.0	90.0	30 07.5	120 25.8	BD	65 04	05	0606	143	455	3.15	100.0	86	184
107.0	30.0	30 30.0	116 03.3	BD	65 04	06	2325	14	92	1.53	100.0	99	442
107.0	31.0	30 27.8	116 07.0	BD	65 04	06	2234	43	153	2.83	100.0	139	93
107.0	32.0	30 25.8	116 11.0	BD	65 04	06	2136	142	465	3.06	100.0	489	426
107.0	35.0	30 21.0	116 22.0	BD	65 04	06	1921	143	454	3.15	100.0	143	186
107.0	40.0	30 11.0	116 39.5	BD	65 04	06	1706	141	469	3.00	100.0	29	140
107.0	45.0	30 00.5	117 00.5	BD	65 04	06	1356	142	477	2.97	100.0	35	663
107.0	50.0	29 51.0	117 22.5	BD	65 04	06	1119	141	468	3.01	100.0	20	109
107.0	55.0	29 41.0	117 43.5	BD	65 04	06	0756	144	453	3.18	100.0	18	152
107.0	60.0	29 31.5	118 01.5	BD	65 04	06	0526	146	451	3.24	100.0	24	152
107.0	65.0	29 21.0	118 21.0	BD	65 04	06	0206	145	464	3.12	100.0	11	118
107.0	70.0	29 11.3	118 41.0	BD	65 04	05	2316	144	466	3.06	100.0	33	160
107.0	75.0	29 01.5	119 00.5	BD	65 04	05	2026	144	468	3.08	100.0	27	175
107.0	80.0	28 52.0	119 19.5	BD	65 04	05	1706	140	472	2.97	100.0	21	126
107.0	85.0	28 42.0	119 39.5	BD	65 04	05	1416	136	466	2.91	100.0	36	937
107.0	90.0	28 31.5	119 59.7	BD	65 04	05	1056	141	462	3.06	100.0	145	401
110.0	32.0	29 52.0	115 47.8	BD	65 04	10	0544	21	87	2.41	100.0	65	20
110.0	35.0	29 46.0	116 00.0	BD	65 04	10	0746	140	454	3.08	100.0	2	10
110.0	40.0	29 38.2	116 19.5	BD	65 04	10	1021	138	486	2.85	100.0	18	37
110.0	45.0	29 27.0	116 38.0	BD	65 04	10	1341	138	434	3.19	100.0	4	18
110.0	50.0	29 16.7	116 58.2	BD	65 04	10	1601	137	494	2.78	100.0	7	56
110.0	55.0	29 06.5	117 19.5	BD	65 04	10	1924	140	474	2.95	100.0	17	8
110.0	60.0	28 57.5	117 41.3	BD	65 04	10	2151	142	467	3.05	100.0	49	10
110.0	65.0	28 46.0	117 58.0	BD	65 04	11	0041	140	478	2.93	100.0	24	46
110.0	70.0	28 36.5	118 18.0	BD	65 04	11	0306	141	435	3.24	100.0	17	229
110.0	75.0	28 26.0	118 36.5	BD	65 04	11	0621	142	446	3.20	100.0	30	282
110.0	80.0	28 17.0	118 56.5	BD	65 04	11	0846	141	462	3.05	100.0	7	32
110.0	85.0	28 06.5	119 16.0	BD	65 04	11	1150	141	460	3.07	100.0	22	118
110.0	90.0	27 56.5	119 35.0	BD	65 04	11	1411	141	457	3.09	100.0	26	212
113.0	28.5	29 25.0	115 11.3	BD	65 04	13	0700	14	63	2.24	100.0	10	46
113.0	32.0	29 24.2	115 13.2	BD	65 04	13	0614	14	116	1.22	100.0	42	47
113.0	36.0	29 22.0	115 18.0	BD	65 04	13	0518	14	205	2.78	100.0	22	127
113.0	40.0	29 11.5	115 38.0	BD	65 04	13	0211	148	426	3.49	100.0	18	3
113.0	45.0	28 52.0	116 18.0	BD	65 04	12	2333	145	448	3.23	100.0	28	7
113.0	50.0	28 42.0	116 40.0	BD	65 04	12	2021	143	458	3.11	100.0	77	13
113.0	55.0	28 31.7	116 57.2	BD	65 04	12	1721	142	462	3.08	100.0	53	42
113.0	60.0	28 22.0	117 16.5	BD	65 04	12	1426	144	404	3.57	100.0	72	154

TABLE 1. (cont.)

## CalCOFI Cruise 6504

Line Station	Lat. (N) deg. min.	Long. (W) deg. min.	Ship Code yr. mo. day	Tow Date yr. mo. day	Time (PST)	Tow Depth (m)	Water Strained (cu. m)	Vol. Haul Factor	Stand- ard Factor	Percent Sorted	Total Larvae	Total Eggs
113.0	65.0	28 12.0	117 33.0	BD	65 04 12	0906	143	434	3.29	100.0	38	707
113.0	70.0	28 03.7	117 53.0	BD	65 04 12	0650	141	457	3.08	100.0	105	211
113.0	75.0	27 55.7	118 09.5	BD	65 04 12	0351	144	436	3.30	100.0	37	122
113.0	80.0	27 46.8	118 30.5	BD	65 04 12	0106	141	457	3.09	100.0	79	202
113.0	85.0	27 35.0	118 50.5	BD	65 04 11	2146	139	467	2.99	100.0	49	309
113.0	90.0	27 23.0	119 11.0	BD	65 04 11	1856	147	434	3.39	100.0	18	102
117.0	24.8	28 58.5	114 36.3	BD	65 04 13	1130	10	93	1.11	100.0	29	254
117.0	25.0	28 58.0	114 37.0	BD	65 04 13	1209	50	179	2.79	100.0	149	2828
117.0	26.0	28 56.0	114 41.5	BD	65 04 13	1258	72	240	3.01	100.0	174	896
117.0	30.0	28 48.0	114 56.5	BD	65 04 13	1542	97	297	3.26	100.0	45	3195
117.0	35.0	28 38.0	115 16.0	BD	65 04 13	1751	142	438	3.25	100.0	252	136
117.0	40.0	28 28.0	115 35.5	BD	65 04 13	2301	141	460	3.06	100.0	91	103
117.0	45.0	28 18.0	115 56.0	BD	65 04 14	0155	141	483	2.91	100.0	45	5
117.0	50.0	28 08.0	116 15.0	BD	65 04 14	0411	139	478	2.91	100.0	129	44
117.0	55.0	27 58.0	116 39.2	BD	65 04 14	0716	140	482	2.91	100.0	223	1
117.0	60.0	27 48.5	116 56.3	BD	65 04 14	0916	139	475	2.93	100.0	31	10
117.0	65.0	27 37.2	117 11.0	BD	65 04 14	1201	138	490	2.83	100.0	37	55
117.0	70.0	27 28.0	117 32.5	BD	65 04 14	1441	138	493	2.80	100.0	11	309
117.0	75.0	27 19.0	117 53.8	BD	65 04 14	1750	141	464	3.04	100.0	23	27
117.0	80.0	27 10.5	118 12.0	BD	65 04 14	2015	140	459	3.06	100.0	139	47
117.0	85.0	26 59.0	118 33.0	BD	65 04 14	2231	141	485	2.91	100.0	97	65
117.0	90.0	26 48.0	118 53.0	BD	65 04 15	0101	139	493	2.82	100.0	122	71
118.0	39.0	28 18.5	115 23.7	BD	65 04 13	2106	141	456	3.09	100.0	850	818
119.0	33.0	28 19.0	114 53.0	BD	65 04 16	1317	113	390	2.90	100.0	35	1896
120.0	22.5	28 28.0	114 04.0	BD	65 04 16	1835	14	68	2.01	100.0	24	573
120.0	23.0	28 26.9	114 06.7	BD	65 04 16	1904	22	79	2.82	100.0	27	120
120.0	24.0	28 25.0	114 10.7	BD	65 04 16	1949	26	124	2.08	100.0	241	321
120.0	25.0	28 22.5	114 15.0	BD	65 04 16	2038	53	202	2.65	100.0	162	253
120.0	30.0	28 13.0	114 34.0	BD	65 04 16	2306	84	294	2.85	100.0	565	1057
120.0	35.0	28 03.0	114 54.0	BD	65 04 16	1046	77	275	2.79	100.0	347	1177
120.0	40.0	27 56.5	115 14.0	BD	65 04 16	0759	30	160	1.90	100.0	229	136
120.0	45.0	27 43.0	115 33.0	BD	65 04 16	0452	139	450	3.09	100.0	30	40
120.0	50.0	27 32.0	115 50.0	BD	65 04 16	0146	141	476	2.97	100.0	64	96
120.0	55.0	27 22.2	116 09.5	BD	65 04 15	2301	141	472	2.98	100.0	38	58
120.0	60.0	27 15.0	116 30.0	BD	65 04 15	1936	139	477	2.91	100.0	80	41
120.0	65.0	27 04.0	116 52.0	BD	65 04 15	1636	141	460	3.06	100.0	56	205
120.0	70.0	26 53.0	117 10.0	BD	65 04 15	1311	144	436	3.30	100.0	68	55
120.0	75.0	26 46.2	117 28.5	BD	65 04 15	1052	140	452	3.09	100.0	14	26
120.0	80.0	26 32.2	117 49.5	BD	65 04 15	0726	148	433	3.43	100.0	27	26
123.0	36.0	27 26.2	114 36.0	BD	65 04 17	0814	39	163	2.41	100.0	64	64
123.0	37.0	27 24.0	114 40.0	BD	65 04 17	0853	62	254	2.46	100.0	21	57
123.0	42.0	27 13.4	114 59.2	BD	65 04 17	1140	139	484	2.86	100.0	791	490
123.0	45.0	27 09.0	115 08.2	BD	65 04 17	1556	140	474	2.91	100.0	1696	862
123.0	50.0	26 58.0	115 30.5	BD	65 04 17	1851	143	431	3.33	100.0	28	230
123.0	55.0	26 48.5	115 49.5	BD	65 04 17					100.0	19	19

TABLE 1. (cont.)

CalCOFI Cruise 6504									
Line	Station	Lat. (N) deg. min.	Long. (W) deg. min.	Ship Code yr. no.	Date mo. day	Tow Depth (PST) (m)	Time Depth (cu. m)	Vol. Water Strained (cu. m)	Stand- ard Haul Factor
123.0	65.0	26 29.0	116 27.5	BD	65 04 18	0001	141	452	3.12
123.0	70.0	26 18.2	116 46.5	BD	65 04 18	0216	142	460	3.08
123.0	75.0	26 08.5	117 04.3	BD	65 04 18	0506	141	452	3.12
123.0	80.0	25 58.7	117 22.8	BD	65 04 18	0731	142	461	3.08
123.0	85.0	25 24.2	114 31.8	BD	65 04 17	0715	10	86	1.22
123.5	35.6	26 58.3	114 00.6	BD	65 04 19	1130	12	75	1.66
127.0	32.6	26 57.5	114 02.2	BD	65 04 19	1053	56	210	2.69
127.0	33.0	26 57.5	114 06.5	BD	65 04 19	0943	67	276	2.44
127.0	34.0	26 55.0	114 26.5	BD	65 04 19	0656	140	472	2.97
127.0	40.0	26 44.5	114 47.0	BD	65 04 19	0356	141	454	3.11
127.0	45.0	26 34.0	115 08.0	BD	65 04 19	0118	141	464	3.03
127.0	50.0	26 23.5	115 21.5	BD	65 04 19	0118	141	476	2.97
127.0	55.0	26 13.5	115 47.5	BD	65 04 18	2206	141	462	3.08
127.0	60.0	26 06.0	116 06.0	BD	65 04 18	1931	142	462	3.08
127.0	65.0	25 55.0	116 06.0	BD	65 04 18	1616	142	481	2.94
127.0	70.0	25 45.3	116 21.5	BD	65 04 18	1401	141	488	2.88
130.0	25.5	26 38.0	113 11.1	BD	65 04 19	2227	15	80	1.82
130.0	26.0	26 37.0	113 13.1	BD	65 04 19	2250	30	123	2.41
130.0	28.0	26 33.0	113 21.0	BD	65 04 20	0014	47	175	2.71
130.0	30.0	26 29.0	113 29.0	BD	65 04 20	0118	71	247	2.86
130.0	35.0	26 19.0	113 48.0	BD	65 04 20	0406	142	454	3.12
130.0	40.0	26 09.8	114 08.4	BD	65 04 20	0626	142	451	3.15
130.0	45.0	25 58.5	114 24.8	BD	65 04 20	0936	141	464	3.04
130.0	50.0	25 48.5	114 46.5	BD	65 04 20	1156	140	475	2.95
130.0	55.0	25 39.0	115 04.5	BD	65 04 20	1501	141	482	2.93
130.0	60.0	25 29.0	115 23.5	BD	65 04 20	1721	142	461	3.09
130.0	65.0	25 19.7	115 43.0	BD	65 04 20	2006	142	460	3.08
130.0	70.0	25 11.0	115 59.2	BD	65 04 20	2216	142	460	3.08
133.0	19.5	26 13.6	112 25.8	BD	65 04 22	0905	14	77	1.76
133.0	21.0	26 12.6	112 32.3	BD	65 04 22	0754	38	172	2.18
133.0	23.0	26 08.5	112 40.2	BD	65 04 22	0648	62	236	2.63
133.0	25.0	26 02.5	112 46.8	BD	65 04 22	0501	130	429	3.04
133.0	30.0	25 51.5	113 06.8	BD	65 04 22	0212	121	389	3.10
133.0	35.0	25 42.6	113 26.7	BD	65 04 21	2306	139	452	3.09
133.0	40.0	25 34.5	113 45.5	BD	65 04 21	2041	141	459	3.08
133.0	45.0	25 25.0	114 06.5	BD	65 04 21	1726	142	472	3.02
133.0	50.0	25 14.5	114 23.0	BD	65 04 21	1451	141	476	2.96
133.0	55.0	25 03.8	114 42.8	BD	65 04 21	1141	141	476	2.96
133.0	60.0	24 55.0	115 01.0	BD	65 04 21	0906	141	467	3.02
133.0	65.0	24 46.0	115 20.5	BD	65 04 21	0606	141	458	3.09
133.0	70.0	24 35.0	115 38.5	BD	65 04 21	0323	141	473	3.02
137.0	20.0	25 07.2	112 07.2	BD	65 04 22	1310	13	76	1.66
137.0	21.0	25 38.1	112 11.0	BD	65 04 22	1349	20	91	2.18
137.0	22.0	25 36.1	112 14.8	BD	65 04 22	1429	45	198	2.27
137.0	23.0	25 34.0	112 19.0	BD	65 04 22	1533	63	238	2.64
137.0	30.0	25 20.0	112 46.0	BD	65 04 22	1841	142	456	3.11

TABLE 1. (cont.)

## CalCOFI Cruise 6504

Line	Station	Lat. (N) deg. min.	Long. (W) deg. min.	Ship Code	Tow yr. mo. day	Time (PST)	Tow Depth (m)	Water Strained (cu. m)	Vol. Haul	Stand- ard Factor	Total Larvae	Total Eggs
137.0	35.0	25 09.8	113 04.5	BD	65	04 22	2138	140	464	3.01	100.0	60
137.0	40.0	25 00.0	113 23.5	BD	65	04 22	2356	140	465	3.00	100.0	52
137.0	45.0	24 49.8	113 42.5	BD	65	04 23	0256	141	464	3.03	100.0	28
137.0	50.0	24 38.0	114 02.5	BD	65	04 23	0506	135	497	2.72	100.0	34
137.0	55.0	24 29.5	114 21.8	BD	65	04 23	0951	142	446	3.19	100.0	23
137.0	60.0	24 21.0	114 40.0	BD	65	04 23	1201	142	462	3.07	100.0	30
137.0	65.0	24 10.5	114 58.2	BD	65	04 23	1501	145	447	3.23	100.0	65
137.0	70.0	23 59.5	115 17.5	BD	65	04 23	1721	144	457	3.16	100.0	48

TABLE 1. (cont.)

CALCOFI Cruise 6507		Line	Station	Lat. (N) deg. min.	Long. (W) deg. min.	Ship Code	Tow Date yr. mo. day	Time (PST)	Tow Depth (m)	Vol. Water Strained (cu. m)	Stand- ard Haul Factor	Percent Sorted	Total Larvae	Total Eggs
60.0	49.2	37	59.0	122 49.5	AX	65 07 16	2034	11	109	0.97	100.0	41	29	
60.0	50.0	37	57.6	122 53.3	AX	65 07 16	2114	28	133	2.09	100.0	0	0	
60.0	51.0	37	55.5	122 57.4	AX	65 07 16	2159	31	152	2.02	100.0	5	260	
60.0	52.0	37	53.5	123 01.8	AX	65 07 16	2323	68	268	2.52	100.0	1	228	
60.0	55.0	37	47.5	123 15.0	AX	65 07 17	0053	68	261	2.62	100.0	35	131	
60.0	60.0	37	36.9	123 37.0	AX	65 07 17	0441	140	495	2.84	100.0	15	389	
60.0	60.0	37	36.9	123 37.0	AX	65 07 17	0931	137	484	2.82	100.0	37	127	
60.0	70.0	37	17.0	124 21.2	AX	65 07 17	1451	148	483	3.06	100.0	13	25	
60.0	80.0	36	57.0	125 04.2	AX	65 07 17	1921	140	503	2.78	100.0	12	19	
60.0	90.0	36	37.0	125 46.7	AX	65 07 17	0056	134	510	2.63	100.0	16	10	
60.0	100.0	36	16.9	126 30.0	AX	65 07 18	0916	137	512	2.68	100.0	8	27	
60.0	120.0	35	36.8	127 55.0	AX	65 07 18	0236	135	516	2.61	100.0	88	30	
60.0	140.0	34	56.5	129 19.0	AX	65 07 18	0656	142	497	2.86	100.0	11	31	
60.0	160.0	34	16.0	130 43.0	AX	65 07 19	1516	141	522	2.70	100.0	15	11	
60.0	180.0	33	36.5	132 05.7	AX	65 07 19	0106	138	506	2.74	100.0	180	24	
60.0	200.0	32	56.2	133 27.5	AX	65 07 20	0916	137	512	2.68	100.0	0	27	
63.0	50.0	37	26.8	122 28.2	AX	65 07 14	2014	9	108	0.86	100.0	0	210	
63.0	51.0	37	23.6	122 33.8	AX	65 07 14	1934	33	139	2.40	100.0	11	5	
63.0	52.0	37	21.0	122 37.7	AX	65 07 14	1829	24	137	1.77	100.0	68	62	
63.0	55.0	37	12.5	122 49.6	AX	65 07 14	1636	140	500	2.80	100.0	77	24	
63.0	60.0	37	02.5	123 11.6	AX	65 07 14	1301	144	506	2.85	100.0	72	14	
63.0	64.0	36	54.4	121 52.9	AX	65 07 13	2330	14	196	0.72	100.0	10	687	
67.0	47.2	36	53.1	121 55.9	AX	65 07 14	0019	27	164	1.64	100.0	3	148	
67.0	48.0	36	51.0	122 00.3	AX	65 07 14	0048	72	236	3.06	100.0	31	144	
67.0	49.0	36	49.0	122 04.5	AX	65 07 14	0208	70	254	2.76	100.0	77	24	
67.0	50.0	36	38.1	122 27.3	AX	65 07 14	0451	142	494	2.88	100.0	42	144	
67.0	55.0	36	38.1	122 27.3	AX	65 07 14	0811	142	465	3.05	100.0	29	174	
67.0	60.0	36	29.0	122 47.6	AX	65 07 14	1836	145	464	3.13	100.0	8	29	
70.0	50.0	36	11.1	121 43.8	AX	65 07 13	1743	73	250	2.90	100.0	25	3	
70.0	51.0	36	10.4	121 45.8	AX	65 07 13	1456	143	497	2.88	100.0	21	6	
70.0	53.0	36	06.5	121 53.8	AX	65 07 13	1136	137	517	2.65	100.0	24	150	
70.0	60.0	35	52.8	122 23.0	AX	65 07 13	0551	146	474	3.07	100.0	22	16	
70.0	70.0	35	33.0	123 06.0	AX	65 07 13	0101	145	459	3.16	100.0	14	77	
70.0	80.0	35	12.8	123 48.0	AX	65 07 12	1746	141	512	2.75	100.0	44	40	
70.0	90.0	34	53.0	124 29.6	AX	65 07 12	1658	73	252	2.90	100.0	23	53	
70.0	100.0	34	32.8	125 12.0	AX	65 07 08	1816	150	472	3.17	100.0	47	13	
70.0	120.0	33	53.5	126 35.0	AX	65 07 12	0206	142	518	2.74	100.0	30	32	
70.0	200.0	31	13.0	132 01.9	AX	65 07 20	1356	138	520	2.65	100.0	53	254	
73.0	49.4	35	38.2	121 14.4	AX	65 07 08	1530	14	252	0.56	100.0	1	150	
73.0	50.0	35	37.3	121 17.1	AX	65 07 08	1658	73	252	2.90	100.0	11	17	
73.0	51.0	35	35.3	121 21.4	AX	65 07 08	1816	150	472	3.17	100.0	18	14	
73.0	53.0	35	31.8	121 28.5	AX	65 07 08	2011	140	502	2.78	100.0	30	32	
73.0	60.0	35	17.9	121 57.3	AX	65 07 08	2326	140	498	2.82	100.0	31	10	
77.0	47.6	35	09.0	120 42.1	AX	65 07 09	1400	7	243	0.29	100.0	3	837	
77.0	48.0	35	08.2	120 43.7	AX	65 07 09	1330	14	232	0.61	100.0	4	877	
77.0	49.0	35	06.3	120 47.8	AX	65 07 09	1208	42	241	1.76	100.0	7	18	
77.0	51.0	35	02.1	120 56.4	AX	65 07 09	1056	492	492	2.87	100.0	78	9	

TABLE 1. (cont.)

## CalCOFI Cruise 6507

Line Station	Lat. (N) deg. min.	Long. (W) deg. min.	Ship Code	Tow Date yr. mo. day	Time (PST)	Tow Depth (m)	Vol. Water (cu. m)	Strained Factor	Stand- ard Haul Factor	Percent Sorted	Total Larvae	Total Eggs
77.0	55.0	34 54.2	121	14.0	AX	65	07	09	0726	145	466	3.10
77.0	57.0	34 49.0	121	22.8	AX	65	07	09	0451	138	498	2.76
80.0	50.2	34 28.2	120	29.4	AX	65	07	09	1850	11	236	0.45
80.0	51.0	34 26.4	120	32.9	AX	65	07	09	1932	71	447	1.59
80.0	52.0	34 24.4	120	36.6	AX	65	07	09	2041	118	549	2.15
80.0	55.0	34 19.7	120	48.3	AX	65	07	09	2246	125	522	2.40
80.0	60.0	34 08.0	121	09.3	AX	65	07	10	0416	136	488	2.80
80.0	65.0	33 58.7	121	32.8	AX	65	07	10	0701	146	492	2.97
80.0	70.0	33 48.7	121	50.5	AX	65	07	10	1056	145	471	3.07
80.0	80.0	33 29.0	122	32.5	AX	65	07	10	1556	146	492	2.96
80.0	90.0	33 09.0	123	12.8	AX	65	07	10	2136	136	487	2.79
80.0	100.0	32 49.0	123	53.1	AX	65	07	11	0236	143	476	3.01
80.0	120.0	32 09.0	125	14.7	AX	65	07	11	1141	143	486	2.95
80.0	200.0	29 28.3	130	38.6	AX	65	07	21	0246	140	498	2.81
82.0	47.0	34 15.0	119	59.0	BD	65	06	16	1356	134	493	2.72
83.0	39.0	34 15.6	119	17.7	BD	65	06	16	0624	112	94	1.27
83.0	40.0	34 13.7	119	21.6	BD	65	06	16	0719	14	91	1.55
83.0	43.0	34 07.8	119	34.2	BD	65	06	16	0916	142	452	3.15
83.0	51.0	33 52.0	120	07.5	BD	65	06	16	1747	100	318	3.15
83.0	55.0	33 45.5	120	21.2	BD	65	06	16	2056	141	425	3.32
83.0	60.0	33 34.0	120	45.0	BD	65	06	17	0006	145	419	3.45
83.0	65.0	33 24.0	121	06.0	BD	65	06	17	0306	143	435	3.29
83.0	70.0	33 14.2	121	26.0	BD	65	06	17	0621	143	424	3.37
83.0	80.0	32 57.0	122	09.0	BD	65	06	17	1231	142	351	4.05
83.0	90.0	32 33.8	122	50.0	BD	65	06	17	1751	140	435	3.21
83.0	92.0	33 54.1	118	27.0	BD	65	06	19	1202	21	121	1.74
87.0	32.5	33 54.2	118	29.4	BD	65	06	19	1119	49	182	2.69
87.0	34.0	33 52.2	118	33.4	BD	65	06	19	1033	68	258	2.64
87.0	35.0	33 50.0	118	37.5	BD	65	06	19	0911	141	458	3.07
87.0	40.0	33 40.0	118	58.0	BD	65	06	19	0606	142	433	3.29
87.0	45.0	33 30.0	119	19.0	BD	65	06	19	0211	138	413	3.35
87.0	50.0	33 20.0	119	39.5	BD	65	06	18	2258	60	222	2.71
87.0	55.0	33 10.0	120	01.0	BD	65	06	18	1931	142	435	3.27
87.0	60.0	33 00.0	120	21.5	BD	65	06	18	1706	143	440	3.25
87.0	65.0	32 49.0	120	42.5	BD	65	06	18	1256	145	452	3.20
87.0	70.0	32 36.0	120	59.0	BD	65	06	18	1031	141	463	3.04
87.0	80.0	32 18.3	121	40.0	BD	65	06	18	0501	143	437	3.26
87.0	90.0	31 59.0	122	25.0	BD	65	06	17	2331	143	459	3.12
90.0	27.5	33 27.6	117	44.0	AX	65	07	25	1004	14	186	0.76
90.0	28.0	33 28.3	117	47.0	AX	65	07	25	0901	130	554	2.35
90.0	32.0	33 21.8	118	02.0	AX	65	07	25	0656	133	520	2.55
90.0	37.0	33 11.0	118	22.6	AX	65	07	25	0311	116	553	2.09
90.0	45.0	32 54.5	118	55.4	AX	65	07	24	2251	130	508	2.56
90.0	53.0	32 40.4	119	27.8	AX	65	07	24	481	138	229	2.87
90.0	60.0	32 25.0	119	57.6	AX	65	07	24	1326	140	516	2.71

TABLE 1. (cont.)

CalCOFI Cruise 6507									
Line	Station	Lat. (N) deg. min.	Long. (W) deg. min.	Ship Code	Tow Date yr. mo. day	Time (PST)	Tow Depth (m)	Strained (cu. m)	Stand- ard Factor
90.0	65.0	32 14.3	120 17.8	AX	65 07 24	0936	138	520	2.65
90.0	70.0	32 04.7	120 38.4	AX	65 07 24	0716	141	508	2.77
90.0	80.0	31 44.2	121 19.1	AX	65 07	0136	136	520	2.62
90.0	90.0	31 24.0	122 01.1	AX	65 07	23	2031	130	562
90.0	100.0	31 04.5	122 38.8	AX	65 07	23	1506	142	519
90.0	120.0	30 25.6	123 59.1	AX	65 07	23	0356	139	518
90.0	140.0	29 46.0	125 20.1	AX	65 07	22	1826	138	515
90.0	160.0	29 05.2	126 40.0	AX	65 07	22	0936	140	540
90.0	180.0	28 24.5	127 58.5	AX	65 07	22	0056	141	505
90.0	200.0	27 45.0	129 18.3	AX	65 07	21	1626	142	516
90.0	226.5	32 56.8	117 16.7	BD	65 06	19	2050	14	110
93.0	27.0	32 56.0	117 19.0	BD	65 06	19	2133	68	238
93.0	28.0	32 54.5	117 22.8	BD	65 06	19	2241	140	436
93.0	30.0	32 50.5	117 31.0	BD	65 06	20	0026	139	451
93.0	35.0	32 40.5	117 51.5	BD	65 06	20	0316	141	464
93.0	40.0	32 30.0	118 11.5	BD	65 06	20	0656	141	448
93.0	45.0	32 19.8	118 30.8	BD	65 06	20	0931	141	430
93.0	50.0	32 09.0	118 49.0	BD	65 06	20	1226	141	427
93.0	55.0	32 00.0	119 09.8	BD	65 06	20	1526	142	430
93.0	60.0	31 50.0	119 34.0	BD	65 06	20	1851	141	423
93.0	65.0	31 39.0	119 52.8	BD	65 06	20	2141	138	426
93.0	70.0	31 28.0	120 13.0	BD	65 06	21	0051	142	428
93.0	80.0	31 05.5	120 50.5	BD	65 06	21	0626	141	448
93.0	90.0	30 44.0	121 28.5	BD	65 06	21	1126	141	450
93.0	96.8	32 19.4	117 04.0	BD	65 06	24	1400	14	143
97.0	29.0	32 17.5	117 04.7	BD	65 06	24	1419	41	161
97.0	30.0	32 16.0	117 07.0	BD	65 06	24	1523	50	178
97.0	32.0	32 12.0	117 15.2	BD	65 06	24	1621	141	418
97.0	35.0	32 05.5	117 28.5	BD	65 06	23	0206	142	437
97.0	40.0	31 57.0	117 51.0	BD	65 06	22	2246	141	431
97.0	45.0	31 46.5	118 09.5	BD	65 06	22	1901	139	436
97.0	50.0	31 35.5	118 31.5	BD	65 06	22	1631	141	425
97.0	55.0	31 25.5	118 50.2	BD	65 06	22	1301	141	465
97.0	60.0	31 20.0	119 11.2	BD	65 06	22	1036	142	452
97.0	65.0	31 10.0	119 31.3	BD	65 06	22	0656	139	473
97.0	70.0	30 58.0	119 51.3	BD	65 06	22	0426	142	462
97.0	80.0	30 36.0	120 32.5	BD	65 06	21	2226	142	440
97.0	90.0	30 15.5	121 10.5	BD	65 06	21	1616	143	461
97.0	96.8	31 42.3	116 43.5	AX	65 07	26	2103	65	156
100.0	29.0	31 40.5	116 46.8	AX	65 07	26	2236	129	512
100.0	30.0	31 30.5	117 07.1	AX	65 07	27	0126	137	494
100.0	40.0	31 21.0	117 27.0	AX	65 07	27	0431	138	501
100.0	45.0	31 11.0	117 47.0	AX	65 07	27	0631	138	470
100.0	50.0	31 02.2	118 08.8	AX	65 07	27	1006	144	490
100.0	55.0	30 50.0	118 27.6	AX	65 07	27	1226	138	477

TABLE 1. (cont.)

## CALCOFI Cruise 6507

Line	Station	Lat. (N) deg. min.	Long. (W) deg. min.	Ship Code	Tow Date yr. mo. day	Tow Time (PST)	Vol. Water (cu. m)	Stand- ard Haul Factor	Total Larvae	Total Eggs
100.0	60.0	30 39.8	118 48.6	AX	65 07	27	1616	1.40	454	3.08
100.0	65.0	30 31.0	119 08.1	AX	65 07	27	1826	1.44	470	3.06
100.0	70.0	30 21.3	119 28.0	AX	65 07	28	0334	1.30	556	2.34
100.0	80.0	30 01.4	120 06.9	AX	65 07	28	0806	1.31	512	2.56
100.0	90.0	29 40.0	120 46.8	AX	65 07	28	1341	1.43	487	2.93
100.0	100.0	29 21.4	121 27.0	AX	65 07	28	1826	1.43	447	2.99
100.0	120.0	28 37.7	122 44.5	AX	65 07	29	0626	1.43	477	3.00
103.0	28.5	31 07.8	116 19.2	BD	65 06	25	0109	1.3	126	1.04
103.0	29.0	31 07.0	116 21.0	BD	65 06	25	0154	28	168	1.68
103.0	30.0	31 06.0	116 24.5	BD	65 06	25	0253	57	217	2.63
103.0	35.0	30 55.8	116 46.5	BD	65 06	25	0516	142	446	3.18
103.0	40.0	30 47.0	117 07.7	BD	65 06	25	0821	142	426	3.33
103.0	45.0	30 39.7	117 24.0	BD	65 06	25	1031	141	419	3.38
103.0	50.0	30 26.0	117 44.5	BD	65 06	25	1406	147	402	3.67
103.0	55.0	30 15.8	118 02.2	BD	65 06	25	1641	145	432	3.36
103.0	60.0	30 05.8	118 20.7	BD	65 06	25	2006	142	431	3.30
103.0	65.0	29 55.0	118 39.2	BD	65 06	25	2241	142	431	3.30
103.0	70.0	29 44.5	118 58.0	BD	65 06	26	0211	144	428	3.36
103.0	80.0	29 24.3	119 44.0	BD	65 06	26	0756	143	476	3.01
103.0	90.0	29 03.3	120 21.2	BD	65 06	26	1416	145	487	3.33
103.0	140.0	27 09.3	123 28.1	AX	65 07	29	1536	138	494	2.79
107.0	30.0	30 30.0	116 03.3	BD	65 06	28	0815	13	122	1.10
107.0	31.0	30 27.8	116 07.0	BD	65 06	28	0729	22	117	1.88
107.0	32.0	30 25.8	116 11.0	BD	65 06	28	0551	141	410	3.44
107.0	35.0	30 21.5	116 22.5	BD	65 06	28	0356	143	414	3.46
107.0	40.0	30 12.4	116 43.0	BD	65 06	28	0051	142	427	3.33
107.0	45.0	30 01.5	117 02.0	BD	65 06	27	2206	144	417	3.45
107.0	50.0	29 50.5	117 22.0	BD	65 06	27	1846	142	426	3.33
107.0	55.0	29 41.0	117 42.0	BD	65 06	27	1606	143	432	3.31
107.0	60.0	29 32.0	118 01.5	BD	65 06	27	1256	144	428	3.37
107.0	65.0	29 21.0	118 21.0	BD	65 06	27	1006	143	449	3.17
107.0	70.0	29 11.0	118 41.0	BD	65 06	27	0621	142	464	3.07
107.0	80.0	28 50.0	119 23.5	BD	65 06	27	0036	144	446	3.22
107.0	90.0	28 31.5	119 59.0	BD	65 06	26	1831	144	431	3.34
107.0	32.0	29 52.0	115 47.9	AX	65 08	02	0120	14	185	0.76
110.0	35.0	29 46.0	116 00.3	AX	65 08	01	2316	139	489	2.85
110.0	40.0	29 36.0	116 19.5	AX	65 08	01	1956	135	479	2.82
110.0	45.0	29 26.5	116 39.0	AX	65 08	01	1706	132	514	2.57
110.0	50.0	29 16.2	116 59.2	AX	65 08	01	1346	141	497	2.84
110.0	55.0	29 06.6	117 20.0	AX	65 08	01	1046	138	486	2.84
110.0	60.0	28 56.1	117 38.1	AX	65 08	01	0736	138	496	2.78
110.0	65.0	28 46.5	117 58.2	AX	65 08	01	0411	142	472	3.03
110.0	70.0	28 36.2	118 18.1	AX	65 08	01	0056	145	481	3.02
110.0	80.0	28 16.4	118 56.7	AX	65 07	31	1941	138	517	2.67
110.0	90.0	27 57.0	119 35.8	AX	65 07	31	1331	140	503	2.79

TABLE 1. (cont.)

## CalCOFI Cruise 6507

Line	Station	Lat. (N) deg. min.	Long. (W) deg. min.	Ship Code	Tow yr. mo. day	Date Time (PST)	Tow Depth (m)	Vol. Water (cu. m)	Tow Strained Factor	Stand- ard Haul	Percent Sorted	Total Larvae	Total Eggs
110.0	100.0	27 36.6	120 16.0	AX	65 07	31	0906	120	565	2.12	100.0	87	1149
110.0	120.0	26 56.0	121 33.0	AX	65 07	30	2326	139	515	2.71	100.0	251	245
110.0	140.0	26 16.8	122 50.0	AX	65 07	30	1501	140	509	2.75	100.0	268	196
110.0	160.0	25 36.0	124 06.3	AX	65 07	30	0206	143	490	2.92	100.0	1043	18
113.0	28.5	25 25.0	115 11.3	BD	65 06	28	1735	15	115	1.29	100.0	14	275
113.0	29.0	29 24.2	115 13.2	BD	65 06	28	1754	21	119	1.80	100.0	14	253
113.0	30.0	29 22.5	115 20.0	BD	65 06	28	1914	51	171	2.98	100.0	25	408
113.0	30.0	30.0	29 22.0	AX	65 08	02	0554	26	165	1.56	100.0	395	2018
113.0	35.0	29 11.5	115 18.2	BD	65 06	28	2131	142	436	3.25	100.0	89	26
113.0	40.0	29 02.0	115 57.0	BD	65 06	29	0056	142	438	3.24	100.0	54	35
113.0	45.0	28 52.0	116 18.0	BD	65 06	29	0336	143	463	3.08	100.0	88	154
113.0	50.0	28 45.0	116 41.0	BD	65 06	29	0641	141	439	3.21	100.0	15	29
113.0	55.0	28 37.0	117 02.0	BD	65 06	29	0911	141	477	2.95	100.0	52	64
113.0	60.0	28 29.0	117 18.0	BD	65 06	29	1206	142	420	3.39	100.0	12	115
113.0	65.0	28 12.0	117 36.0	BD	65 06	29	1501	132	435	3.03	100.0	21	138
113.0	70.0	28 02.0	117 54.5	BD	65 06	29	1746	143	419	3.40	100.0	16	275
113.0	80.0	27 42.0	118 32.0	BD	65 06	29	2151	142	414	3.44	100.0	27	422
113.0	90.0	27 22.0	119 09.0	BD	65 06	30	0246	142	436	3.27	100.0	47	217
115.0	35.0	28 54.0	115 27.1	AX	65 08	02	0926	132	507	2.61	100.0	135	23
117.0	24.8	28 58.2	114 36.0	BD	65 07	02	0115	14	72	1.91	100.0	268	99
117.0	25.0	28 58.0	114 36.5	BD	65 07	02	0229	41	174	2.36	100.0	408	170
117.0	26.0	28 56.0	114 41.5	BD	65 07	01	2338	58	217	2.67	100.0	154	162
117.0	30.0	28 48.0	114 56.5	BD	65 07	01	2113	72	245	2.94	100.0	38	865
117.0	35.0	28 38.0	115 16.0	BD	65 07	01	1826	144	503	2.86	100.0	23	13
117.0	40.0	28 28.0	115 35.5	BD	65 07	01	1216	138	533	2.59	100.0	57	16
117.0	45.0	28 18.0	115 56.0	BD	65 07	01	0831	142	422	3.36	100.0	20	30
117.0	50.0	28 09.0	116 15.0	BD	65 07	01	0456	144	433	3.32	100.0	63	206
117.0	55.0	27 58.0	116 34.5	BD	65 07	01	0216	143	432	3.30	100.0	122	40
117.0	60.0	27 48.0	116 53.0	BD	65 06	30	2241	142	370	3.85	100.0	160	55
117.0	65.0	27 37.3	117 13.0	BD	65 06	30	2011	143	429	3.33	100.0	38	118
117.0	70.0	27 28.0	117 32.5	BD	65 06	30	1646	143	459	3.10	100.0	48	296
117.0	80.0	27 08.0	118 10.5	BD	65 06	30	1151	143	432	3.30	100.0	30	743
117.0	90.0	26 49.0	118 42.0	BD	65 06	30	0701	143	439	3.26	100.0	89	2987
118.0	39.0	28 18.5	115 23.7	BD	65 07	01	1436	142	508	2.80	100.0	151	71
119.0	33.0	28 18.7	114 53.0	AX	65 08	02	1443	62	228	2.17	100.0	626	240
120.0	30.0	28 13.0	114 34.0	BD	65 07	02	1043	69	255	2.72	100.0	416	610
120.0	22.5	28 28.0	114 04.0	BD	65 07	02	0610	15	144	1.03	100.0	150	811
120.0	23.0	28 26.0	114 06.7	BD	65 07	02	0659	21	145	1.47	100.0	125	684
120.0	24.0	28 24.0	114 10.7	BD	65 07	02	0740	28	237	1.16	100.0	687	431
120.0	25.0	28 22.5	114 15.0	BD	65 07	02	0843	56	281	2.00	100.0	1470	441
120.0	26.0	28 21.5	114 19.0	BD	65 07	02	1043	69	255	2.72	100.0	416	610
120.0	27.0	28 03.0	114 54.0	BD	65 07	02	1343	70	269	2.60	100.0	654	68
120.0	40.0	27 56.5	115 14.0	BD	65 07	02	1544	29	113	2.58	100.0	802	77
120.0	45.0	27 43.0	115 33.0	AX	65 08	03	0556	139	487	2.86	100.0	179	679
120.0	50.0	27 33.0	115 52.0	AX	65 08	03	1816	141	467	3.01	100.0	53	41
120.0	55.0	27 23.2	116 11.2	AX	65 08	03	523	134	2026	2.56	100.0	199	56

TABLE 1. (cont.)

## CalCOFI Cruise 6507

Line	Station	Lat. (N) deg. min.	Long. (W) deg. min.	Ship Code	Tow Date yr. mo. day	Tow Time (PST)	Water Depth (m)	Strained Vol. (cu. m)	Total Larvae	Stand- ard Haul Factor	Percent Sorted	Total Eggs
120.0	60.0	27 13.0	1116 30.7	AX	65 08	03	2326	139	492	2.82	100.0	144
120.0	65.0	27 03.0	1116 49.7	AX	65 08	04	0216	134	506	2.65	100.0	94
120.0	70.0	26 52.0	1117 09.8	AX	65 08	04	0606	139	491	2.83	100.0	94
120.0	80.0	26 32.5	1117 48.4	AX	65 08	04	1036	135	502	2.69	100.0	544
120.0	100.0	25 52.2	1119 05.7	AX	65 08	04	2016	136	524	2.60	100.0	97
120.0	120.0	25 13.0	120 22.0	AX	65 08	05	0501	140	511	2.75	100.0	79
120.0	136.0	27 26.2	1114 36.0	BD	65 07	02	2209	21	137	1.53	100.0	77
1223.0	37.0	27 24.0	1114 40.0	BD	65 07	02	2318	57	252	2.25	100.0	42
1223.0	42.0	27 14.0	1114 59.0	AX	65 08	11	1121	135	496	3.15	100.0	94
1223.0	42.0	27 14.0	1114 59.0	BD	65 07	03	0146	141	449	3.15	100.0	33
1223.0	45.0	27 08.0	1115 11.5	BD	65 07	03	0411	144	435	3.32	100.0	124
1223.0	50.0	26 58.0	1115 31.0	BD	65 07	03	0621	143	504	2.84	100.0	151
1223.0	55.0	26 47.0	1115 50.0	BD	65 07	03	0946	143	501	2.84	100.0	49
1223.0	60.0	26 35.0	1116 06.0	BD	65 07	03	1206	141	513	2.74	100.0	64
1223.0	65.0	26 25.0	1116 28.0	BD	65 07	03	1506	142	505	2.81	100.0	107
1223.0	70.0	26 19.0	1116 47.0	BD	65 07	03	1716	142	513	2.77	100.0	61
1223.0	80.0	26 00.5	1117 22.0	BD	65 07	03	2206	143	508	2.82	100.0	153
1223.5	35.6	27 24.2	1114 31.8	BD	65 07	03	2140	14	110	1.27	100.0	130
1227.0	32.6	26 58.3	1114 00.6	BD	65 07	05	0525	11	82	1.37	100.0	26
1227.0	33.0	26 57.5	1114 02.2	BD	65 07	05	0454	48	161	2.97	100.0	5
1227.0	34.0	26 55.0	1114 06.0	BD	65 07	05	0333	63	261	2.42	100.0	71
1227.0	40.0	26 43.5	1114 29.0	AX	65 08	11	0616	136	499	2.73	100.0	47
1227.0	40.0	26 45.5	1114 28.0	BD	65 07	05	0046	142	501	2.84	100.0	47
1227.0	45.0	26 33.5	1114 48.5	BD	65 07	04	2121	142	441	3.23	100.0	144
1227.0	50.0	26 23.0	1115 08.0	BD	65 07	04	1836	144	494	2.91	100.0	144
1227.0	55.0	26 13.5	1115 27.0	BD	65 07	04	1516	143	523	2.74	100.0	62
1227.0	60.0	26 04.4	1115 45.0	BD	65 07	04	1246	144	491	2.93	100.0	32
1227.0	65.0	25 52.5	1116 05.5	BD	65 07	04	0941	143	497	2.88	100.0	49
1227.0	70.0	25 46.5	1116 22.0	BD	65 07	04	0716	144	506	2.84	100.0	265
1227.0	80.0	25 26.0	1116 57.0	BD	65 07	04	0211	142	521	2.73	100.0	630
130.0	25.5	26 38.0	1113 11.0	AX	65 08	07	0850	14	173	0.81	100.0	7800
130.0	26.0	26 37.0	1113 13.0	AX	65 08	07	1031	137	524	2.62	100.0	458
130.0	28.0	26 33.3	1113 20.5	AX	65 08	07	0726	141	499	2.83	100.0	325
130.0	30.0	26 29.2	1113 29.0	AX	65 08	07	0451	140	511	2.48	100.0	140
130.0	35.0	26 19.0	1113 47.6	AX	65 08	07	1554	44	217	2.02	100.0	140
130.0	40.0	26 08.8	1114 07.2	AX	65 08	07	0211	142	514	2.55	100.0	132
130.0	45.0	25 58.9	1114 26.5	AX	65 08	07	1031	137	524	2.62	100.0	89
130.0	50.0	25 49.0	1114 46.0	AX	65 08	07	0726	141	499	2.83	100.0	102
130.0	55.0	25 39.2	1115 04.7	AX	65 08	07	0146	140	512	2.73	100.0	77
130.0	60.0	25 29.2	1115 24.0	AX	65 08	06	2301	140	498	2.81	100.0	603
130.0	70.0	25 09.0	1116 01.5	AX	65 08	06	1711	134	540	2.49	100.0	110
130.0	80.0	24 49.2	1116 39.2	AX	65 08	06	1226	140	512	2.73	100.0	73
130.0	90.0	24 28.8	1117 17.9	AX	65 08	06	0726	141	519	2.71	100.0	257
130.0	100.0	24 09.0	1117 55.0	AX	65 08	06	0246	142	499	2.84	100.0	261
130.0	120.0	23 30.0	1119 10.0	AX	65 08	05	1736	141	519	2.71	100.0	26

TABLE 1. (cont.)

## CalCOFI Cruise 6507

Line	Station	Lat. (N) deg. min.	Long. (W) deg. min.	Ship Code	Tow yr. mo. day	Date (PST)	Time (PST)	Tow Depth (m)	Water Strained (cu. m)	Stand- ard Haul Factor	Percent Sorted	Total Larvae	Total Eggs
133.0	19.5	26 13.6	112 25.8	BD	65 07	05	1625	10	127	0.83	100.0	28	5084
133.0	21.0	26 12.6	112 32.3	BD	65 07	05	1709	42	210	2.02	100.0	32	865
133.0	23.0	26 08.5	112 40.2	BD	65 07	05	1843	71	282	2.51	100.0	2	20
133.0	25.0	26 04.5	112 48.0	BD	65 07	05	1948	78	289	2.69	100.0	10	12
133.0	30.0	25 54.5	113 07.5	BD	65 07	05	2256	141	476	2.96	100.0	153	235
133.0	35.0	25 45.0	113 27.0	BD	65 07	06	0116	140	475	2.96	100.0	29	187
133.0	40.0	25 36.0	113 47.0	BD	65 07	06	0401	140	483	2.91	100.0	316	164
133.0	45.0	25 24.5	114 05.0	BD	65 07	06	0631	138	519	2.66	100.0	92	96
133.0	50.0	25 13.0	114 22.5	BD	65 07	06	0936	138	526	2.62	100.0	44	6
133.0	55.0	25 02.0	114 42.0	BD	65 07	06	1201	141	532	2.66	100.0	43	28
133.0	60.0	24 54.5	115 02.0	BD	65 07	06	1531	142	508	2.80	100.0	33	60
133.0	65.0	24 44.5	115 20.5	BD	65 07	06	1746	141	533	2.64	100.0	38	133
133.0	70.0	24 34.5	115 39.0	BD	65 07	06	2051	140	524	2.68	100.0	284	874
133.0	80.0	24 14.5	116 17.0	BD	65 07	07	0116	142	556	2.55	100.0	117	243
137.0	20.0	25 40.0	112 07.3	AX	65 08	08	0254	7	168	0.42	100.0	4536	>999999
137.0	21.0	25 38.0	112 10.9	AX	65 08	08	0324	21	154	1.37	100.0	6374	51318
137.0	22.0	25 36.0	112 14.8	AX	65 08	08	0419	32	151	2.14	100.0	2696	7202
137.0	23.0	25 34.6	112 18.8	AX	65 08	08	0518	45	237	1.90	100.0	171	2113
137.0	30.0	25 20.1	112 46.0	AX	65 08	08	0906	138	501	2.76	100.0	85	27
137.0	35.0	25 10.5	113 04.2	AX	65 08	08	1131	136	514	2.65	100.0	3	11
137.0	40.0	25 00.2	113 23.6	AX	65 08	08	1511	142	499	2.84	100.0	5	13
137.0	45.0	24 50.4	113 42.2	AX	65 08	08	1726	141	433	3.25	100.0	118	175
137.0	50.0	24 40.0	114 02.0	AX	65 08	08	2046	135	518	2.60	100.0	167	969
137.0	55.0	24 30.0	114 20.0	AX	65 08	08	2306	137	496	2.76	100.0	101	640
137.0	60.0	24 20.0	114 39.4	AX	65 08	09	0306	121	560	2.16	100.0	240	216
137.0	70.0	24 15.0	115 18.0	AX	65 08	09	0746	130	535	2.44	100.0	20	205
137.0	80.0	23 39.8	115 55.5	AX	65 08	09	1236	131	527	2.49	100.0	108	185
140.0	28.8	24 48.0	112 19.3	AX	65 08	10	1400	9	178	0.48	100.0	100	985
140.0	29.0	24 47.5	112 20.0	AX	65 08	10	1334	33	168	1.96	100.0	241	1078
140.0	30.0	24 45.5	112 24.0	AX	65 08	10	1218	72	254	2.85	100.0	30	91
140.0	35.0	24 35.4	112 42.9	AX	65 08	10	0946	133	500	2.66	100.0	28	67
140.0	40.0	24 25.5	113 01.8	AX	65 08	10	0651	139	494	2.81	100.0	21	24
140.0	45.0	24 15.5	113 20.6	AX	65 08	10	0341	139	503	2.76	100.0	146	59
140.0	50.0	24 05.0	113 40.0	AX	65 08	10	0056	128	553	2.31	100.0	365	100

TABLE 1. (cont.)

## CalCOFI Cruise 6509

Line	Station	Lat. (N) deg. min.	Long. (W) deg. min.	Ship Code yr.	Tow Date mo. day	Time (PST)	Tow Depth (m)	Water Strained (cu. m)	Vol. Water (cu. m)	Stand- ard Haul Factor	Percent Sorted	Total Larvae	Total Eggs
80.0	50.2	34 28.0	120 29.1	BD	65 08	31	1215	14	96	1.50	100.0	0	47
80.0	51.0	34 26.0	120 32.5	BD	65 08	31	1251	143	568	2.51	100.0	2	34
80.0	52.0	34 24.3	120 36.5	BD	65 08	31	1406	141	532	2.65	100.0	37	35
80.0	55.0	34 18.8	120 48.2	BD	65 08	31	1536	142	539	2.64	100.0	4	4
80.0	60.0	34 09.0	121 09.0	BD	65 08	31	1816	141	523	2.69	100.0	24	0
80.0	65.0	33 59.0	121 30.0	BD	65 08	31	2046	142	480	2.97	100.0	48	6
80.0	70.0	33 48.7	121 51.0	BD	65 08	31	2326	142	523	2.71	100.0	66	0
80.0	75.0	33 38.7	122 11.5	BD	65 09	01	0136	142	521	2.73	100.0	44	1
80.0	80.0	33 28.7	122 32.0	BD	65 09	01	0411	144	478	3.02	100.0	17	5
82.0	47.0	34 15.0	119 59.0	BD	65 08	31	0741	139	550	2.53	100.0	45	2
83.0	39.0	34 15.5	119 17.7	BD	65 09	02	0825	14	73	1.93	100.0	40	541
83.0	40.0	34 13.6	119 21.8	BD	65 09	02	0735	16	68	2.36	100.0	41	978
83.0	43.0	34 08.0	119 34.0	BD	65 09	02	0516	142	492	2.88	100.0	229	75
83.0	51.0	33 52.0	120 08.5	BD	65 09	02	0018	72	292	2.48	100.0	31	155
83.0	55.0	33 45.3	120 21.3	BD	65 09	01	2216	141	460	3.06	100.0	159	6
83.0	60.0	33 34.0	120 45.0	BD	65 09	01	1916	140	528	2.66	100.0	77	5
83.0	65.0	33 24.0	121 06.0	BD	65 09	01	1636	142	503	2.82	100.0	2	1
83.0	70.0	33 12.0	121 24.0	BD	65 09	01	1411	143	516	2.77	100.0	3	3
83.0	75.0	33 04.5	121 46.5	BD	65 09	01	1121	144	504	2.86	100.0	4	6
83.0	80.0	32 54.0	122 07.0	BD	65 09	01	0831	142	503	2.83	100.0	4	5
87.0	32.5	33 55.2	118 26.5	BD	65 09	02	1409	14	81	1.66	100.0	122	1042
87.0	33.0	33 54.2	118 29.4	BD	65 09	02	1449	44	170	2.58	100.0	71	142
87.0	34.0	33 52.2	118 33.4	BD	65 09	02	1528	63	240	2.64	100.0	133	164
87.0	35.0	33 50.0	118 37.5	BD	65 09	02	1616	141	466	3.02	100.0	134	329
87.0	40.0	33 40.0	118 58.0	BD	65 09	02	1936	144	472	3.05	100.0	207	48
87.0	45.0	33 30.0	119 19.0	BD	65 09	02	2206	143	446	3.20	100.0	260	22
87.0	50.0	33 20.0	119 39.5	BD	65 09	03	0048	69	260	2.65	100.0	421	14
87.0	55.0	33 10.0	120 00.0	BD	65 09	03	0326	139	494	2.82	100.0	19	1
87.0	60.0	33 00.0	120 21.5	BD	65 09	03	0556	142	468	3.03	100.0	5	2
87.0	65.0	32 50.0	120 41.0	BD	65 09	03	0811	140	484	2.89	100.0	8	6
87.0	70.0	32 40.0	121 00.2	BD	65 09	03	1031	142	468	3.03	100.0	5	9
87.0	75.0	32 31.0	121 19.0	BD	65 09	03	1256	148	458	3.22	100.0	4	8
87.0	80.0	32 19.5	121 43.0	BD	65 09	03	1556	141	472	2.99	100.0	4	2
90.0	27.5	33 29.5	117 44.7	BD	65 09	05	0800	13	99	1.30	100.0	35	81
90.0	28.0	33 28.5	117 46.7	BD	65 09	05	0706	142	496	2.85	100.0	27	9
90.0	32.0	33 21.2	118 01.6	BD	65 09	05	0406	140	498	2.82	100.0	50	123
90.0	37.0	33 11.0	118 22.5	BD	65 09	05	0126	142	485	2.93	100.0	26	30
90.0	45.0	32 54.6	118 55.5	BD	65 09	04	2131	142	468	3.03	100.0	13	4
90.0	53.0	32 39.0	119 28.5	BD	65 09	04	1746	142	499	2.84	100.0	6	3
90.0	60.0	32 25.0	119 57.5	BD	65 09	04	1411	143	482	2.97	100.0	2	6
90.0	65.0	32 15.0	120 18.8	BD	65 09	04	1146	143	483	2.96	100.0	7	16
90.0	70.0	32 10.0	120 41.5	BD	65 09	04	0856	142	477	2.97	100.0	84	187
90.0	75.0	31 59.0	121 00.0	BD	65 09	04	0646	142	491	2.88	100.0	4	109
90.0	80.0	31 47.2	121 22.2	BD	65 09	04	0351	143	465	3.08	100.0	5	131
90.0	85.0	31 35.8	121 41.0	BD	65 09	04	0106	146	450	3.24	100.0	19	67

TABLE 1. (cont.)

## CalCOFI Cruise 6509

Line	Station	Lat. (N) deg. min.	Long. (W) deg. min.	Ship Code	Tow yr. no.	Date day	Tow Time (PST)	Depth (m)	Strained (cu. m)	Tow Water Factor	Stand- ard Factor	Haul Percent Sorted	Total Larvae	Total Eggs
90.0	90.0	31 24.0	122 01.0	BD	65	09 03	2216	144	468	3.08	100.0	66	227	
93.0	26.5	32 56.8	117 16.7	BD	65	09 05	1229	13	82	1.58	100.0	441	200	
93.0	27.0	32 56.0	117 19.0	BD	65	09 05	1301	99	455	2.18	100.0	32	67	
93.0	28.0	32 54.7	117 21.8	BD	65	09 05	1341	135	506	2.67	100.0	43	39	
93.0	30.0	32 50.5	117 31.0	BD	65	09 05	1536	141	479	2.95	100.0	485	36	
93.0	35.0	32 40.5	117 51.5	BD	65	09 05	1826	158	422	3.73	100.0	5	19	
93.0	40.0	32 30.0	118 11.5	BD	65	09 05	2121	144	456	3.16	100.0	20	5	
93.0	45.0	32 20.0	118 32.0	BD	65	09 06	0001	142	466	3.05	100.0	15	1	
93.0	50.0	32 10.0	118 52.5	BD	65	09 06	0231	141	486	2.90	100.0	10	1	
93.0	55.0	32 00.0	119 13.5	BD	65	09 06	0506	142	491	2.88	100.0	5	1	
93.0	60.0	31 50.0	119 34.0	BD	65	09 06	0716	141	506	2.78	100.0	4	31	
93.0	65.0	31 41.3	119 54.0	BD	65	09 06	0931	142	479	2.96	100.0	8	15	
93.0	70.0	31 28.5	120 12.8	BD	65	09 06	1206	141	510	2.76	100.0	12	39	
93.0	75.0	31 19.5	120 35.0	BD	65	09 06	1426	142	496	2.86	100.0	21	54	
93.0	80.0	31 08.2	120 52.8	BD	65	09 06	1641	140	478	2.94	100.0	121	79	
93.0	85.0	31 00.0	121 13.5	BD	65	09 06	1931	141	472	2.99	100.0	198	42	
93.0	90.0	30 50.7	121 34.5	BD	65	09 06	2156	139	487	2.85	100.0	132	133	
96.8	28.7	32 19.4	117 03.8	BD	65	09 10	1504	13	84	1.54	100.0	24	130	
97.0	29.0	32 17.5	117 04.7	BD	65	09 10	1533	48	195	2.44	100.0	25	170	
97.0	30.0	32 16.0	117 07.0	BD	65	09 10	1609	53	233	2.27	100.0	24	76	
97.0	32.0	32 12.0	117 15.2	BD	65	09 08	0526	142	487	3.08	100.0	26	28	
97.0	35.0	32 07.2	117 29.0	BD	65	09 08	0331	143	489	2.91	100.0	33	3	
97.0	40.0	31 56.0	117 48.0	BD	65	09 08	0046	145	477	3.03	100.0	38	5	
97.0	45.0	31 46.0	118 08.5	BD	65	09 07	2201	143	471	3.04	100.0	46	1	
97.0	50.0	31 36.0	118 29.0	BD	65	09 07	1926	143	482	2.96	100.0	26	3	
97.0	55.0	31 25.5	118 49.5	BD	65	09 07	1651	142	498	2.84	100.0	2	5	
97.0	60.0	31 15.5	119 10.0	BD	65	09 07	1421	142	505	2.81	100.0	7	127	
97.0	65.0	31 01.5	119 31.0	BD	65	09 07	1146	140	505	2.78	100.0	2	20	
97.0	70.0	30 53.0	119 52.0	BD	65	09 07	0911	143	494	2.89	100.0	9	30	
97.0	75.0	30 45.0	120 11.0	BD	65	09 07	0651	142	440	3.23	100.0	17	44	
97.0	80.0	30 35.0	120 31.0	BD	65	09 11	0406	140	482	2.90	100.0	105	105	
100.0	29.0	31 42.2	116 43.4	BD	65	09 10	2053	72	173	4.19	100.0	1	5	
100.0	30.0	31 40.5	116 46.5	BD	65	09 10	2131	139	506	2.75	100.0	39	5	
100.0	35.0	31 30.5	117 07.0	BD	65	09 11	0041	143	472	3.04	100.0	76	49	
100.0	40.0	31 22.5	117 27.5	BD	65	09 11	0311	144	458	3.14	100.0	56	2	
100.0	45.0	31 12.0	117 48.5	BD	65	09 11	0531	145	468	3.10	100.0	67	1	
100.0	50.0	31 02.5	118 11.0	BD	65	09 11	0836	139	455	3.05	100.0	7	47	
100.0	55.0	30 50.5	118 27.5	BD	65	09 11	1106	144	476	3.02	100.0	25	47	
100.0	60.0	30 41.0	118 47.0	BD	65	09 11	1336	142	497	2.86	100.0	29	92	
100.0	65.0	30 29.0	119 09.5	BD	65	09 11	1611	143	474	3.02	100.0	126	339	
100.0	70.0	30 19.0	119 29.0	BD	65	09 11	1831	147	474	3.09	100.0	134	65	
103.0	28.5	31 07.8	116 19.2	BD	65	09 12	2309	12	125	1.00	100.0	27	1258	
103.0	29.0	31 07.0	116 21.0	BD	65	09 12	2234	27	156	1.71	100.0	76	167	
103.0	30.0	31 06.0	116 24.5	BD	65	09 12	2143	50	192	2.59	100.0	50	6	
103.0	35.0	30 51.6	116 49.0	BD	65	09 12	1726	141	494	2.85	100.0	9	2	

TABLE 1. (cont.)

## CALCOFI CRUISE 6509

Line	Station	Lat. (N) deg. min.	Long. (W) deg. min.	Ship Code	Tow Date yr. mo. day	Time (PST)	Tow Depth (m)	Water Strained (cu. m)	Vol. (cu. m)	Stand- ard Factor	Percent Sorted	Total Larvae	Total Eggs
103.0	40.0	30 45.0	117 04.7	BD	65 09 12	1446	138	440	3.13	100.0	136	7	7
103.0	45.0	30 36.5	117 21.5	BD	65 09 12	1226	145	447	3.24	100.0	7	0	0
103.0	50.0	30 26.0	117 44.5	BD	65 09 12	0951	143	456	3.14	100.0	12	8	8
103.0	55.0	30 16.0	118 05.0	BD	65 09 12	0701	142	459	3.09	100.0	14	17	17
103.0	60.0	30 06.0	118 25.0	BD	65 09 12	0436	145	470	3.09	100.0	209	71	71
103.0	65.0	29 56.5	118 44.0	BD	65 09 12	0201	143	449	3.19	100.0	171	108	108
103.0	70.0	29 46.5	119 04.0	BD	65 09 11	2321	140	482	2.90	100.0	127	121	121
107.0	30.0	30 30.0	116 03.3	BD	65 09 13	0354	9	145	0.61	100.0	7	341	341
107.0	31.0	30 27.8	116 07.0	BD	65 09 13	0444	36	150	2.38	100.0	39	128	128
107.0	32.0	30 25.8	116 11.0	BD	65 09 13	0526	140	467	3.00	100.0	21	12	12
107.0	35.0	30 21.5	116 22.5	BD	65 09 13	0741	142	480	2.96	100.0	9	0	0
107.0	40.0	30 12.2	116 41.7	BD	65 09 13	1011	140	480	2.93	100.0	7	0	0
107.0	45.0	30 03.2	116 59.5	BD	65 09 13	1236	147	448	3.28	100.0	14	1	1
107.0	50.0	29 54.0	117 20.0	BD	65 09 13	1521	145	453	3.19	100.0	15	24	24
107.0	55.0	29 44.0	117 39.0	BD	65 09 13	1751	140	458	3.06	100.0	21	59	59
107.0	60.0	29 32.5	118 01.7	BD	65 09 13	2056	141	477	2.95	100.0	49	69	69
107.0	65.0	29 21.0	118 21.0	BD	65 09 13	2341	143	435	3.20	100.0	210	84	84
107.0	70.0	29 11.0	118 41.0	BD	65 09 14	0216	141	465	3.03	100.0	140	64	64
110.0	32.0	29 52.0	115 47.8	BD	65 09 15	0430	14	158	0.86	100.0	77	622	622
110.0	35.0	29 46.0	116 00.0	BD	65 09 15	0201	140	469	2.98	100.0	20	12	12
110.0	40.0	29 36.0	116 18.2	BD	65 09 14	2316	142	451	3.14	100.0	33	1	1
110.0	45.0	29 25.7	116 37.8	BD	65 09 14	2036	143	427	3.35	100.0	41	5	5
110.0	50.0	29 17.0	116 55.5	BD	65 09 14	1751	142	462	3.08	100.0	10	10	10
110.0	55.0	29 06.4	117 06.4	BD	65 09 14	1511	143	466	3.06	100.0	43	55	55
110.0	60.0	28 56.5	117 38.0	BD	65 09 14	1226	142	477	2.98	100.0	75	21	21
110.0	65.0	28 46.0	117 59.0	BD	65 09 14	0936	142	458	3.11	100.0	21	35	35
113.0	28.5	29 25.0	115 11.3	BD	65 09 15	1005	10	92	1.12	100.0	66	297	297
113.0	29.0	29 24.2	115 13.2	BD	65 09 15	1034	20	110	1.78	100.0	119	71	71
113.0	30.0	29 22.0	115 18.0	BD	65 09 15	1118	50	185	2.68	100.0	19	123	123
113.0	35.0	29 11.5	115 22.0	BD	65 09 15	1821	141	467	3.01	100.0	35	27	27
113.0	40.0	29 02.0	115 57.0	BD	65 09 15	2131	141	455	3.09	100.0	88	4	4
113.0	45.0	28 52.0	116 18.0	BD	65 09 16	0006	131	530	2.47	100.0	203	8	8
113.0	50.0	28 41.5	116 36.5	BD	65 09 16	0226	141	470	3.00	100.0	554	11	11
113.0	55.0	28 32.0	116 57.0	BD	65 09 16	0501	142	478	2.97	100.0	97	23	23
113.0	60.0	28 22.0	117 16.5	BD	65 09 16	0711	140	485	2.88	100.0	32	1	1
115.0	35.0	28 54.5	115 25.0	BD	65 09 15	1526	143	457	3.13	100.0	50	53	53
117.0	24.8	28 58.5	114 36.3	BD	65 09 17	1015	10	87	1.15	100.0	38	372	372
117.0	25.0	28 58.0	114 37.0	BD	65 09 17	0949	42	164	2.57	100.0	73	1147	1147
117.0	26.0	28 56.0	114 41.5	BD	65 09 17	0828	65	248	2.63	100.0	247	171	171
117.0	30.0	28 48.0	114 56.5	BD	65 09 17	0648	68	260	2.60	100.0	58	236	236
117.0	35.0	28 38.0	115 16.0	BD	65 09 17	0336	140	479	2.92	100.0	98	35	35
117.0	40.0	28 28.0	115 35.5	BD	65 09 16	2236	142	458	3.11	100.0	42	8	8
117.0	45.0	28 18.0	115 56.0	BD	65 09 16	1946	140	483	2.89	100.0	107	31	31
117.0	50.0	28 08.0	116 15.0	BD	65 09 16	1656	139	495	2.80	100.0	107	5	5
117.0	55.0	27 58.0	116 34.5	BD	65 09 16	1421	141	483	2.91	100.0	187	187	187

TABLE 1. (cont.)

Line	Station	Lat. (N) deg. min.	Long. (W) deg. min.	Ship Code	Tow yr.	Date mo. day	Time (PST)	Vol. Water (cu. m)	Tow Depth (m)	Stand- ard Haul Factor	Percent Sorted	Total Larvae	Total Eggs
117.0	60.0	27 48.0	116 53.0	BD	65	09 16	1146	469	3.03	100.0	101	0	343
118.0	39.0	28 18.5	115 23.7	BD	65	09 17	0041	497	2.72	100.0	172	35	289
119.0	33.0	28 19.0	114 53.0	BD	65	09 17	2258	72	2.86	100.0	141	1265	195
120.0	22.5	28 28.0	114 04.0	BD	65	09 17	1504	13	1.57	100.0	1.88	359	174
120.0	23.0	28 26.9	114 06.7	BD	65	09 17	1539	20	104	2.38	100.0	284	1151.2
120.0	24.0	28 25.0	114 10.7	BD	65	09 17	1624	28	118	2.33	100.0	135	1170
120.0	25.0	28 22.5	114 15.0	BD	65	09 17	1704	46	198	2.98	100.0	452	1130
120.0	30.0	28 13.0	114 34.0	BD	65	09 17	2003	72	240	3.01	100.0	165	112
120.0	35.0	28 03.0	114 54.0	BD	65	09 18	0108	71	258	2.74	100.0	379	117
120.0	40.0	27 56.5	115 14.0	BD	65	09 18	0349	36	130	2.73	100.0	78	112
120.0	45.0	27 43.0	115 33.0	BD	65	09 19	1441	143	479	2.98	100.0	63	90
120.0	50.0	27 33.0	115 52.5	BD	65	09 19	1711	143	482	2.96	100.0	14	8
120.0	55.0	27 23.0	116 12.0	BD	65	09 19	1941	142	477	2.97	100.0	116	116
120.0	60.0	27 13.0	116 30.5	BD	65	09 19	2201	146	465	3.14	100.0	250	4
123.0	36.0	27 26.2	114 36.0	BD	65	09 20	1559	36	112	3.18	100.0	78	3293
123.0	37.0	27 24.0	114 40.0	BD	65	09 20	1448	66	222	2.99	100.0	14	8
123.0	42.0	27 20.1	114 59.0	BD	65	09 20	1211	144	480	3.01	100.0	116	116
123.0	45.0	27 12.5	115 11.0	BD	65	09 20	1021	145	478	3.03	100.0	3	23
123.0	50.0	27 03.0	115 30.0	BD	65	09 20	0736	146	473	3.08	100.0	51	10
123.0	55.0	26 51.0	115 51.5	BD	65	09 20	0456	142	459	3.10	100.0	27	8
123.0	60.0	26 41.5	116 10.5	BD	65	09 20	0221	145	469	3.10	100.0	164	30
123.5	35.6	27 24.2	114 31.8	BD	65	09 20	1645	10	72	1.43	100.0	325	256
127.0	32.6	26 58.3	114 00.6	BD	65	09 20	2130	13	149	0.87	100.0	173	405
127.0	33.0	26 57.5	114 02.2	BD	65	09 20	2159	34	133	2.60	100.0	34	199
127.0	34.0	26 55.0	114 06.5	BD	65	09 20	2248	73	241	3.01	100.0	19	37
127.0	40.0	26 43.0	114 29.5	BD	65	09 21	0156	144	457	3.14	100.0	23	19
127.0	45.0	26 32.8	114 51.0	BD	65	09 21	0426	142	459	3.10	100.0	58	10
127.0	50.0	26 23.0	115 08.0	BD	65	09 21	0646	141	467	3.02	100.0	29	44
127.0	55.0	26 15.0	115 27.0	BD	65	09 21	0856	145	457	3.18	100.0	17	315
127.0	60.0	26 06.0	115 45.0	BD	65	09 21	1121	146	432	3.37	100.0	25	249
130.0	25.5	26 38.0	113 11.1	BD	65	09 22	1225	14	73	1.92	100.0	233	69
130.0	26.0	26 37.0	113 31.1	BD	65	09 22	1154	28	120	2.32	100.0	263	103
130.0	28.0	26 33.0	113 21.0	BD	65	09 22	0944	48	204	2.37	100.0	71	52
130.0	30.0	26 29.0	113 29.0	BD	65	09 22	0808	67	260	2.58	100.0	77	41
130.0	35.0	26 19.0	113 47.5	BD	65	09 22	0546	141	461	3.05	100.0	29	10
130.0	40.0	26 07.0	114 06.5	BD	65	09 22	0236	141	469	3.00	100.0	226	14
130.0	45.0	25 57.8	114 26.0	BD	65	09 22	0001	140	461	3.03	100.0	129	77
130.0	50.0	25 49.0	114 45.0	BD	65	09 21	2126	147	442	3.32	100.0	96	340
130.0	55.0	25 41.0	115 05.5	BD	65	09 21	1836	143	460	3.10	100.0	421	179
130.0	60.0	25 32.0	115 23.0	BD	65	09 21	1611	144	459	3.13	100.0	28	177
133.0	19.5	26 13.6	112 25.8	BD	65	09 23	1750	11	93	1.21	100.0	237	155
133.0	21.0	26 12.6	112 32.0	BD	65	09 23	1848	46	201	2.29	100.0	476	210
133.0	23.0	26 08.5	112 40.0	BD	65	09 23	2008	61	12	1.61	100.0	121	161
133.0	24.5	26 04.5	112 48.0	BD	65	09 23	2113	72	243	2.94	100.0	63	39
133.0	30.0	25 54.8	113 07.0	BD	65	09 24	0006	141	492	2.86	100.0	113	0

TABLE 1. (cont.)

## CalCOFI Cruise 6509

Line	Station	Lat. (N) deg. min.	Long. (W) deg. min.	Ship Code	Tow Date yr. mo. day	Tow Time (PST)	Depth (m)	Strained Water (cu. m)	Stand- ard Haul Factor	Percent Sorted	Total Larvae	Total Eggs
133.0	35.0	25 45.0	113 27.5	BD	65 09 24	0251	141	475	2.96	100.0	82	52
133.0	40.0	25 36.5	113 47.5	BD	65 09 24	0530	145	476	3.04	100.0	28	617
133.0	45.0	25 25.0	114 05.0	BD	65 09 24	0811	139	494	2.82	100.0	2	63
133.0	50.0	25 13.6	114 25.0	BD	65 09 24	1051	140	474	2.95	100.0	5	35
133.0	50.0	25 40.1	112 07.2	BD	65 09 25	0835	14	72	1.88	100.0	65	111
137.0	20.0	25 38.1	112 11.0	BD	65 09 25	0749	20	93	2.16	100.0	192	987
137.0	21.0	25 36.1	112 14.8	BD	65 09 25	0714	30	146	2.04	100.0	151	453
137.0	22.0	25 34.0	112 19.0	BD	65 09 25	0613	68	244	2.76	100.0	98	270
137.0	23.0	25 20.0	112 46.0	BD	65 09 25	0236	143	469	3.05	100.0	87	8
137.0	30.0	25 10.0	113 04.5	BD	65 09 25	2341	142	461	3.09	100.0	50	168
137.0	35.0	25 00.0	113 23.5	BD	65 09 24	2111	138	462	2.99	100.0	66	200
137.0	40.0	25 48.8	113 44.3	BD	65 09 24	1801	145	477	3.04	100.0	97	23
137.0	45.0	24 39.5	114 03.0	BD	65 09 24	1516	141	469	3.00	100.0	13	55

TABLE 2. Pooled occurrences of fish larvae taken during CalCOFI cruises in 1965.

Rank	Taxon	Occurrences
1	<i>Engraulis mordax</i>	618
2	<i>Triphoturus mexicanus</i>	494
3	<i>Vinciguerra lucetia</i>	436
4	<i>Sebastes</i> spp.	387
5	<i>Citharichthys</i> spp.	342
6	Disintegrated fish larva	319
7	<i>Leuroglossus stilbius</i>	300
8	<i>Merluccius productus</i>	290
9	<i>Cyclothona</i> spp.	265
10	<i>Stenobrachius leucopsarus</i>	263
10	Unidentified fish larva	263
12	<i>Protomyctophum crockeri</i>	261
13	<i>Diogenichthys laternatus</i>	249
14	<i>Lampanyctus ritteri</i>	234
15	Myctophidae	222
16	<i>Bathylagus wesethi</i>	220
17	<i>Trachurus symmetricus</i>	214
18	<i>Lampanyctus</i> spp.	183
19	<i>Ceratoscopelus townsendi</i>	156
20	<i>Melamphaes</i> spp.	151
21	Sciaenidae	147
22	<i>Tarletonbeania crenularis</i>	132
23	<i>Bathylagus ochotensis</i>	127
24	<i>Diogenichthys atlanticus</i>	116
25	<i>Symbolophorus californiensis</i>	111
26	<i>Paralichthys californicus</i>	107
27	<i>Sardinops sagax</i>	104
27	Gobiidae	104
29	<i>Hygophum atratum</i>	103
30	<i>Stomias atriventer</i>	100
31	<i>Pleuronichthys verticalis</i>	88
32	<i>Gonichthys tenuiculus</i>	81
32	<i>Parophrys vetulus</i>	81
34	<i>Diaphus</i> spp.	80
35	<i>Sympodus</i> spp.	75
36	<i>Hypsoblennius</i> spp.	73
37	<i>Lestidiops ringens</i>	67
38	<i>Citharichthys stigmaeus</i>	65
39	Clinidae	64
40	Serranidae	59
40	Sternopychidae	59
42	<i>Chauliodus macouni</i>	57
43	<i>Icichthys lockingtoni</i>	53
44	<i>Argentina sialis</i>	49
45	<i>Peprius simillimus</i>	45
46	<i>Hippoglossina stomata</i>	44
46	<i>Notoscopelus resplendens</i>	44
48	<i>Synodus</i> spp.	42

TABLE 2. (cont.)

Rank	Taxon	Occurrences
49	<i>Ophidion scrippsae</i>	40
49	Chiasmodontidae	40
51	<i>Chromis punctipinnis</i>	39
52	Ophidiiformes	37
52	Cottidae	37
54	<i>Halichoeres</i> spp.	36
55	<i>Etrumeus acuminatus</i>	35
56	<i>Scomber japonicus</i>	34
57	<i>Lyopsetta exilis</i>	33
57	Scopelarchidae	33
57	<i>Idiacanthus antrostomus</i>	33
60	<i>Poromitra</i> spp.	32
60	<i>Lampadena urophaos</i>	32
62	<i>Oxyjulis californica</i>	31
62	<i>Tetragonurus cuvieri</i>	31
64	<i>Prionotus</i> spp.	30
65	<i>Chilara taylori</i>	29
66	Trichiuridae	27
66	<i>Semicossyphus pulcher</i>	27
66	<i>Hygophum reinhardtii</i>	27
69	Carangidae	25
69	<i>Sphyraena argentea</i>	25
69	<i>Scorpaena</i> spp.	25
72	Anguilliformes	24
73	<i>Nansenia crassa</i>	22
73	Agonidae	22
75	<i>Notolychnus valdiviae</i>	21
76	<i>Scorpaenichthys marmoratus</i>	20
76	<i>Sebastolobus</i> spp.	20
78	Trachipteridae	19
78	<i>Scopelosaurus</i> spp.	19
78	<i>Myctophum nitidulum</i>	19
81	Gonostomatidae	18
81	Ceratioidei	18
83	<i>Microstoma microstoma</i>	17
84	Cyclopteridae	16
84	<i>Microstomus pacificus</i>	16
86	Gerreidae	14
86	<i>Seriola lalandi</i>	14
88	<i>Scopelogadus bispinosus</i>	13
88	<i>Oxylebius pictus</i>	13
90	<i>Syngnathus</i> spp.	12
91	Haemulidae	11
91	<i>Diogenichthys</i> spp.	11
91	<i>Hygophum</i> spp.	11
94	<i>Hypsopsetta guttulata</i>	10
94	<i>Coryphaena hippurus</i>	10
96	<i>Pleuronichthys ritteri</i>	9
96	<i>Sarda chiliensis</i>	9

TABLE 2. (cont.)

Rank	Taxon	Occurrences
96	<i>Lampanyctus regalis</i>	9
96	<i>Psettichthys melanostictus</i>	9
96	<i>Nansenia candida</i>	9
96	<i>Brama</i> spp.	9
96	<i>Notolepis risso</i>	9
96	<i>Aristostomias scintillans</i>	9
96	<i>Cololabis saira</i>	9
96	Gobiesocidae	9
106	<i>Xystreurus liolepis</i>	8
106	<i>Auxis</i> spp.	8
106	Atherinidae	8
106	Gempylidae	8
106	<i>Loweina rara</i>	8
106	<i>Hypsypops rubicundus</i>	8
106	Pomacentridae	8
113	<i>Scopeloberyx robustus</i>	7
113	<i>Bathylagus</i> spp.	7
113	<i>Zaniolepis</i> spp.	7
113	<i>Brosmophycis marginata</i>	7
113	<i>Ichthyococcus</i> spp.	7
113	Labridae	7
113	<i>Macroramphosus gracilis</i>	7
113	<i>Caulolatilus princeps</i>	7
121	<i>Photonectes</i> spp.	6
121	<i>Stemonosudis macrura</i>	6
123	<i>Pleuronichthys coenosus</i>	5
123	<i>Scomberomorus</i> spp.	5
123	<i>Medialuna californiensis</i>	5
126	<i>Glyptocephalus zachirus</i>	4
126	<i>Howella brodiei</i>	4
126	<i>Bathophilus</i> spp.	4
126	Pleuronectiformes	4
126	Scombridae	4
126	<i>Pleuronichthys decurrens</i>	4
126	<i>Sudis atrox</i>	4
126	Blennioidei	4
134	<i>Vinciguerria poweriae</i>	3
134	Paralepididae	3
134	Tetraodontidae	3
134	<i>Girella nigricans</i>	3
134	Macrouridae	3
134	<i>Bathylagus pacificus</i>	3
134	<i>Physiculus</i> spp.	3
134	<i>Diplophos taenia</i>	3
142	<i>Tactostoma macropus</i>	2
142	<i>Lepidopsetta bilineata</i>	2
142	<i>Opisthonema</i> spp.	2
142	<i>Centrobranchus</i> spp.	2
142	Exocoetidae	2

TABLE 2. (cont.)

Rank	Taxon	Occurrences
142	<i>Electrona rissoi</i>	2
148	Nomeidae	1
148	<i>Pleuronichthys</i> spp.	1
148	Hexagrammidae	1
148	<i>Eustomias</i> spp.	1
148	<i>Caristius macropus</i>	1
148	Hemiramphidae	1
148	Soleidae	1
148	<i>Bathylagus milleri</i>	1
148	Stomiiformes	1
148	<i>Triphoturus nigrescens</i>	1
148	<i>Syacium ovale</i>	1
148	<i>Paralepis atlantica</i>	1
148	Evermannellidae	1
148	<i>Mugil</i> spp.	1

TABLE 3. Pooled numbers of fish larvae taken during CalCOFI cruises in 1965. Counts are adjusted for percent of sample sorted and standard haul factor (see text).

Rank	Taxon	Count
1	<i>Engraulis mordax</i>	321562
2	<i>Vinciguerria lucetia</i>	40034
3	<i>Sebastes</i> spp.	24848
4	<i>Merluccius productus</i>	21963
5	<i>Triphoturus mexicanus</i>	18173
6	<i>Leuroglossus stilbius</i>	17761
7	<i>Sardinops sagax</i>	16659
8	<i>Citharichthys</i> spp.	14756
9	<i>Stenobrachius leucopsarus</i>	12130
10	<i>Trachurus symmetricus</i>	8456
11	Sciaenidae	6466
12	<i>Diogenichthys laternatus</i>	6277
13	<i>Bathylagus wesethi</i>	3418
14	Serranidae	3214
15	Unidentified fish larva	2918
16	<i>Cyclothona</i> spp.	2748
17	<i>Ceratoscopelus townsendi</i>	2733
18	Disintegrated fish larva	2039
19	<i>Lampanyctus ritteri</i>	2021
20	Haemulidae	1974
21	<i>Syphurus</i> spp.	1794
22	<i>Etrumeus acuminatus</i>	1739
23	<i>Lampanyctus</i> spp.	1451
24	Myctophidae	1308
25	<i>Protomyctophum crockeri</i>	1247
26	<i>Tarletonbeania crenularis</i>	1213
27	<i>Bathylagus ochotensis</i>	1190
28	<i>Prionotus</i> spp.	957
29	<i>Diogenichthys atlanticus</i>	889
30	<i>Parophrys vetulus</i>	861
31	<i>Symbolophorus californiensis</i>	712
32	<i>Chromis punctipinnis</i>	710
33	<i>Hygophum atratum</i>	709
34	Carangidae	698
35	<i>Hypsypops rubicundus</i>	680
36	<i>Diaphus</i> spp.	670
37	Gobiidae	669
38	<i>Melamphaes</i> spp.	655
39	<i>Paralichthys californicus</i>	631
40	<i>Synodus</i> spp.	575
41	<i>Stomias atriventer</i>	518
42	<i>Ophidion scrippsae</i>	516
43	<i>Gonichthys tenuiculus</i>	501
44	<i>Hypsoblennius</i> spp.	448
45	<i>Halichoeres</i> spp.	439
46	Ophidiiformes	418
47	<i>Peprilus simillimus</i>	390

TABLE 3. (cont.)

Rank	TAXON	Count
48	Clinidae	349
49	<i>Sphyraena argentea</i>	344
50	<i>Pleuronichthys verticalis</i>	327
51	<i>Icichthys lockingtoni</i>	313
52	<i>Argentina sialis</i>	306
53	<i>Citharichthys stigmatus</i>	293
54	<i>Notoscopelus resplendens</i>	290
55	<i>Lestidiops ringens</i>	287
56	<i>Scomber japonicus</i>	286
57	<i>Hygophum reinhardtii</i>	259
58	Gerreidae	250
59	Sternoptychidae	236
60	<i>Oxyjulis californica</i>	227
61	<i>Tetragonurus cuvieri</i>	219
62	<i>Chauliodus macouni</i>	212
63	Anguilliformes	210
64	<i>Auxis</i> spp.	195
65	<i>Hippoglossina stomata</i>	190
66	Cottidae	181
66	<i>Scomberomorus</i> spp.	181
66	<i>Lampadena urophaos</i>	181
69	<i>Lyopsetta exilis</i>	176
70	<i>Idiacanthus antrostomus</i>	170
71	Chiasmodontidae	152
72	<i>Notolychnus valdiviae</i>	141
73	Pomacentridae	140
74	Gonostomatidae	137
75	Scopelarchidae	136
76	<i>Scorpaena</i> spp.	123
77	<i>Poromitra</i> spp.	118
78	Trichiuridae	109
79	<i>Chilara taylori</i>	104
80	<i>Nansenia crassa</i>	102
81	<i>Semicossyphus pulcher</i>	101
82	<i>Sarda chiliensis</i>	95
82	Atherinidae	95
84	<i>Seriola lalandi</i>	93
85	<i>Sebastolobus</i> spp.	87
86	<i>Myctophum nitidulum</i>	75
87	Agonidae	74
88	Ceratioidei	65
89	<i>Hygophum</i> spp.	64
90	<i>Scopelosaurus</i> spp.	62
90	<i>Microstomus pacificus</i>	62
92	<i>Scorpaenichthys marmoratus</i>	61
92	<i>Microstoma microstoma</i>	61
94	Trachipteridae	57
95	<i>Scopeloberyx robustus</i>	50
95	<i>Scopelogadus bispinosus</i>	50

TABLE 3. (cont.)

Rank	Taxon	Count
97	<i>Opisthonema</i> spp.	46
98	<i>Xystreurus liolepis</i>	44
99	<i>Diogenichthys</i> spp.	40
99	<i>Bathylagus</i> spp.	40
99	<i>Brama</i> spp.	40
102	<i>Nansenia candida</i>	38
102	<i>Lampanyctus regalis</i>	38
102	<i>Notolepis risso</i>	38
102	Labridae	38
106	<i>Oxylebius pictus</i>	36
106	Gobiesocidae	36
108	Cyclopteridae	35
109	<i>Aristostomias scintillans</i>	34
110	<i>Coryphaena hippurus</i>	33
111	<i>Cololabis saira</i>	32
112	<i>Pleuronichthys ritteri</i>	31
113	Gempylidae	28
113	<i>Syngnathus</i> spp.	28
115	<i>Photonectes</i> spp.	27
116	<i>Hypsopsetta guttulata</i>	26
117	<i>Vinciguerria poweriae</i>	25
118	<i>Macroramphosus gracilis</i>	24
118	<i>Psettichthys melanostictus</i>	24
120	<i>Ichthyococcus</i> spp.	23
121	<i>Loweina rara</i>	22
122	<i>Caulolatilus princeps</i>	21
123	Blennioidei	20
124	<i>Zaniolepis</i> spp.	19
125	<i>Brosmophycis marginata</i>	18
126	<i>Sudis atrox</i>	17
127	<i>Glyptocephalus zachirus</i>	16
127	<i>Stemonosudis macrura</i>	16
127	Scombridae	16
127	Paralepididae	16
131	<i>Medialuna californiensis</i>	15
132	<i>Bathophilus</i> spp.	14
133	<i>Pleuronichthys coenosus</i>	13
133	<i>Pleuronichthys decurrens</i>	13
135	Nomeidae	12
135	<i>Bathylagus pacificus</i>	12
137	<i>Howella brodiei</i>	11
138	<i>Syacium ovale</i>	9
138	<i>Physiculus</i> spp.	9
138	Macrouridae	9
138	<i>Lepidopsetta bilineata</i>	9
138	Pleuronectiformes	9
143	<i>Diplophos taenia</i>	8
143	<i>Centrobranchus</i> spp.	8
145	<i>Girella nigricans</i>	7

TABLE 3. (cont.)

Rank	TAXON	Count
146	Exocoetidae	6
146	<i>Electrona rissoii</i>	6
146	<i>Tactostoma macropus</i>	6
146	Stomiiformes	6
150	Tetraodontidae	5
151	<i>BathyLAGUS milleri</i>	3
151	<i>Paralepis atlantica</i>	3
151	<i>Caristius macropus</i>	3
151	<i>Mugil</i> spp.	3
151	Evermannellidae	3
151	<i>Eustomias</i> spp.	3
151	<i>Triphoturus nigrescens</i>	3
151	<i>Pleuronichthys</i> spp.	3
159	Soleidae	2
160	Hemiramphidae	1
160	Hexagrammidae	1
	Total	562034

TABLE 4. Numbers of fish larvae taken on stations occupied during CalCOFI cruises in 1965. Counts are adjusted for percent of sample sorted and standard haul factor (see text). Average number is given for stations occupied twice during a single month. Unoccupied stations are indicated by a dash.

Anguilliformes												
STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
60.0	200.0	0.0	-	-	-	-	-	-	-	-	-	-
87.0	333.0	0.0	-	-	0.0	-	0.0	-	-	2.6	-	-
113.0	35.0	-	0.0	-	0.0	-	0.0	-	-	3.0	-	-
120.0	65.0	-	2.8	-	0.0	-	-	0.0	0.0	0.0	-	-
127.0	40.0	-	1.4	-	0.0	-	-	0.0	0.0	0.0	-	-
130.0	25.5	-	0.0	-	0.0	-	-	-	0.8	0.0	-	-
130.0	30.0	-	0.0	-	0.0	-	-	-	0.0	0.0	2.6	-
130.0	35.0	-	0.0	-	3.1	-	-	-	0.0	0.0	6.1	-
130.0	40.0	-	3.1	-	0.0	-	-	-	0.0	0.0	11.4	-
133.0	30.0	-	0.0	-	0.0	-	-	0.0	-	8.9	-	-
133.0	35.0	-	2.9	-	3.1	-	-	0.0	-	3.0	-	-
133.0	40.0	-	0.0	-	0.0	-	-	0.0	-	2.8	-	-
133.0	45.0	-	0.0	-	3.0	-	-	0.0	-	-	-	-
133.0	60.0	-	2.9	-	0.0	-	-	0.0	-	13.2	-	-
137.0	20.0	-	0.0	-	0.0	-	-	-	0.0	47.5	-	-
137.0	21.0	-	0.0	-	0.0	-	-	-	0.0	28.6	-	-
137.0	22.0	-	0.0	-	0.0	-	-	-	0.0	46.9	-	-
137.0	23.0	-	0.0	-	0.0	-	-	-	0.0	3.0	-	-
137.0	30.0	-	0.0	-	0.0	-	-	-	0.0	3.1	-	-
137.0	35.0	-	0.0	-	0.0	-	-	-	0.0	-	-	-

Etrumeus acuminatus												
STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
87.0	34.0	0.0	-	0.0	-	0.0	-	-	-	5.3	-	-
117.0	25.0	-	0.0	-	0.0	-	0.0	-	0.0	15.4	-	-
117.0	26.0	-	0.0	-	0.0	-	0.0	-	0.0	5.3	-	-
118.0	39.0	-	0.0	-	0.0	-	0.0	-	0.0	114.2	-	-
120.0	22.5	-	0.0	-	0.0	-	0.0	-	2.9	22.6	-	-
120.0	23.0	-	0.0	-	0.0	-	0.0	-	0.0	7.1	-	-
120.0	24.0	-	0.0	-	0.0	-	0.0	-	0.0	4.7	-	-
120.0	25.0	-	0.0	-	0.0	-	0.0	-	2.7	9.0	-	-
120.0	30.0	-	0.0	-	0.0	-	0.0	-	2.6	2.7	-	-
120.0	35.0	-	0.0	-	0.0	-	0.0	-	0.0	199.3	-	-
120.0	40.0	-	0.0	-	0.0	-	0.0	-	0.0	35.0	-	-
123.0	36.0	-	0.0	-	0.0	-	0.0	-	0.0	6.0	-	-
123.0	37.0	-	0.0	-	0.0	-	0.0	-	0.0	82.9	-	-
123.5	35.6	-	0.0	-	0.0	-	0.0	-	0.0	106.9	-	-
127.0	32.6	-	0.0	-	0.0	-	0.0	-	0.0	9.9	-	-
127.0	34.0	-	0.0	-	0.0	-	0.0	-	0.0	5.8	-	-
130.0	25.5	-	0.0	-	0.0	-	0.0	-	0.0	-	134.6	-
130.0	26.0	-	0.0	-	0.0	-	0.0	-	0.0	0.8	-	-

TABLE 4. (cont.)

*Etrumeus acuminatus* (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
130.0	28.0	-	0.0	-	0.0	-	-	-	0.0	4.7	-	-
130.0	30.0	-	0.0	-	0.0	-	-	-	0.0	5.2	-	-
133.0	21.0	-	0.0	-	0.0	-	-	-	0.0	-	6.9	-
133.0	23.0	-	0.0	-	0.0	-	-	-	0.0	-	5.0	-
133.0	30.0	-	0.0	-	0.0	-	-	-	0.0	-	2.9	-
137.0	21.0	-	0.0	-	0.0	-	-	-	75.3	4.3	-	-
137.0	22.0	-	0.0	-	0.0	-	-	-	451.5	0.0	-	-
137.0	23.0	-	0.0	-	0.0	-	-	-	383.8	5.5	-	-
140.0	28.8	-	0.0	-	0.0	-	-	-	1.9	-	-	-
140.0	29.0	-	0.0	-	-	-	-	-	7.8	-	-	-

*Opisthonema* spp.

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
140.0	28.8	-	0.0	-	-	-	-	-	9.1	-	-	-
140.0	29.0	-	0.0	-	-	-	-	-	37.2	-	-	-

*Sardinops sagax*

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
60.0	49.5	0.9	-	-	-	-	-	-	-	-	-	-
60.0	50.0	1.3	-	-	-	-	-	-	0.0	-	-	-
67.0	47.2	-	-	-	-	-	-	-	2.2	-	-	-
70.0	80.0	0.0	-	-	2.7	-	-	-	0.0	-	-	-
73.0	53.0	0.0	-	-	3.0	-	-	-	0.0	-	-	-
77.0	47.6	0.0	-	-	0.6	-	-	-	0.0	-	-	-
83.0	43.0	0.0	-	-	0.0	-	-	-	3.2	-	-	-
87.0	33.0	0.0	-	-	2.6	-	-	-	5.4	-	-	-
87.0	40.0	0.0	-	-	0.0	-	-	-	3.3	-	-	-
90.0	28.0	5.6	-	-	0.0	-	-	-	0.0	-	-	-
90.0	32.0	2.7	-	-	0.0	-	-	-	2.5	-	-	-
93.0	26.5	12.0	-	-	-	-	-	-	32.8	-	-	-
93.0	27.0	0.0	-	-	0.0	-	-	-	17.2	-	-	-
93.0	28.0	0.0	-	-	0.0	-	-	-	3.2	-	-	-
93.0	30.0	0.0	-	-	0.0	-	-	-	0.0	-	-	-
97.0	29.0	5.6	-	-	0.0	-	-	-	0.0	-	-	-
103.0	28.5	1.3	-	-	-	-	-	-	12.5	-	-	-
103.0	29.0	0.0	-	-	7.2	-	-	-	6.7	-	-	-
107.0	30.0	-	0.0	-	-	-	-	-	1.6	-	-	-
107.0	31.0	-	0.0	-	-	-	-	-	3.1	-	-	-
107.0	32.0	-	0.0	-	-	-	-	-	8.5	-	-	-
110.0	32.0	-	0.0	-	-	-	-	-	12.2	-	-	-
113.0	29.0	-	2.2	-	-	-	-	-	2.4	-	0.0	0.9
113.0	30.0	-	-	-	-	-	-	-	0.0	-	3.6	0.0

TABLE 4. (cont.)

*Sardinops sagax* (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
113.0	50.0	-	0.0	-	0.0	-	3.2	-	-	0.0	-	-
117.0	24.8	-	0.0	-	2.2	-	-	1.9	-	3.5	-	-
117.0	25.0	-	5.2	-	2.8	-	0.0	-	-	7.7	-	-
117.0	26.0	-	0.0	-	0.0	-	5.3	-	-	2.6	-	-
117.0	35.0	-	0.0	-	0.0	-	0.0	-	-	14.6	-	-
117.0	40.0	-	0.0	-	0.0	-	0.0	-	-	15.5	-	-
117.0	45.0	-	0.0	-	0.0	-	0.0	-	-	135.8	-	-
118.0	39.0	-	0.0	-	0.0	-	0.0	-	-	323.7	-	-
119.0	33.0	-	0.0	-	0.0	-	-	8.7	-	22.9	-	-
120.0	22.5	-	0.0	-	0.0	-	13.4	-	-	141.3	-	-
120.0	23.0	-	43.8	-	0.0	-	27.9	-	-	203.0	-	-
120.0	24.0	-	137.8	-	0.0	-	1.2	-	-	273.7	-	-
120.0	25.0	-	22.0	-	0.0	-	4.0	-	-	28.0	-	-
120.0	30.0	-	0.0	-	0.0	-	0.0	-	-	848.8	-	-
120.0	35.0	-	0.0	-	0.0	-	0.0	-	-	238.4	-	-
120.0	40.0	-	0.0	-	0.0	-	0.0	-	-	581.5	-	-
120.0	60.0	-	0.0	-	0.0	-	-	0.0	-	3.1	-	-
123.0	36.0	-	0.0	-	0.0	-	0.0	-	-	143.1	-	-
123.0	42.0	-	24.7	-	0.0	-	0.0	-	-	3.2	-	-
123.5	35.6	-	8.9	-	0.0	-	0.0	-	-	264.5	-	-
127.0	32.6	-	0.0	-	1.7	-	2.7	-	-	13.1	-	-
127.0	33.0	-	0.0	-	0.0	-	0.0	-	-	10.4	-	-
127.0	34.0	-	2.6	-	0.0	-	2.4	-	-	0.0	-	-
127.0	45.0	-	0.0	-	0.0	-	0.0	-	-	0.0	-	-
130.0	25.5	-	0.0	-	0.0	-	58.1	-	-	991.4	-	-
130.0	26.0	-	1.9	-	0.0	-	-	-	-	47.3	-	-
130.0	28.0	-	0.0	-	0.0	-	-	0.0	-	167.0	-	-
130.0	30.0	-	0.0	-	0.0	-	-	0.0	-	7.1	-	-
133.0	19.5	-	0.0	-	0.0	-	-	-	-	12.9	-	-
133.0	21.0	-	40.0	-	0.0	-	-	0.8	-	0.0	-	-
133.0	23.0	-	5.3	-	0.0	-	20.2	-	-	18.3	-	-
133.0	30.0	-	0.0	-	0.0	-	0.0	-	-	5.0	-	-
137.0	20.0	-	0.5	-	0.0	-	5.9	-	-	5.7	-	-
137.0	21.0	-	49.1	-	0.0	-	-	-	-	1167.6	-	-
137.0	22.0	-	271.7	-	0.0	-	-	-	-	5384.1	-	-
137.0	23.0	-	18.5	-	0.0	-	-	-	-	3103.0	-	-
137.0	30.0	-	0.0	-	0.0	-	-	-	-	1130.5	-	-
137.0	50.0	-	0.0	-	0.0	-	-	-	-	8.3	-	-
140.0	28.8	-	15.0	-	0.0	-	-	-	-	2.6	-	-
140.0	29.0	-	60.8	-	0.0	-	-	-	-	0.0	-	-
140.0	30.0	-	219.5	-	-	-	-	-	-	0.0	-	-

TABLE 4. (cont.)

*Engraulis mordax*

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
60.0	49.5	4.3	-	-	-	-	-	-	-	-	-	-
60.0	50.0	2.6	-	-	-	-	-	-	-	-	-	-
60.0	51.0	4.9	-	-	-	-	-	-	-	-	-	-
60.0	52.0	9.0	-	-	-	-	-	-	-	-	-	-
63.0	49.7	2.9	-	-	-	-	-	-	-	-	-	-
63.0	50.0	5.7	-	-	-	-	-	-	-	-	-	-
63.0	51.0	5.3	-	-	-	-	-	-	-	-	-	-
63.0	55.0	3.2	-	-	-	-	-	-	-	-	-	-
63.0	60.0	0.0	-	-	-	-	-	-	-	-	-	-
67.0	47.2	-	-	-	-	-	-	-	-	-	-	-
67.0	47.3	0.0	-	-	-	-	-	-	-	-	-	-
67.0	48.0	0.0	-	-	-	-	-	-	-	-	-	-
67.0	49.0	0.0	-	-	-	-	-	-	-	-	-	-
67.0	50.0	0.0	-	-	-	-	-	-	-	-	-	-
67.0	55.0	0.0	-	-	-	-	-	-	-	-	-	-
67.0	60.0	0.0	-	-	-	-	-	-	-	-	-	-
67.0	70.0	-	-	-	-	-	-	-	-	-	-	-
70.0	50.0	2.9	-	-	-	-	-	-	-	-	-	-
70.0	51.0	0.0	-	-	-	-	-	-	-	-	-	-
70.0	52.0	6.3	-	-	-	-	-	-	-	-	-	-
70.0	53.0	5.9	-	-	-	-	-	-	-	-	-	-
70.0	60.0	0.0	-	-	-	-	-	-	-	-	-	-
70.0	65.0	-	-	-	-	-	-	-	-	-	-	-
70.0	70.0	0.0	-	-	-	-	-	-	-	-	-	-
73.0	49.4	0.0	-	-	-	-	-	-	-	-	-	-
73.0	50.0	7.0	-	-	-	-	-	-	-	-	-	-
73.0	51.0	21.6	-	-	-	-	-	-	-	-	-	-
73.0	53.0	0.0	-	-	-	-	-	-	-	-	-	-
73.0	60.0	0.0	-	-	-	-	-	-	-	-	-	-
77.0	47.6	0.3	-	-	-	-	-	-	-	-	-	-
77.0	48.0	1.2	-	-	-	-	-	-	-	-	-	-
77.0	49.0	-	-	-	-	-	-	-	-	-	-	-
77.0	51.0	0.0	-	-	-	-	-	-	-	-	-	-
77.0	55.0	0.0	-	-	-	-	-	-	-	-	-	-
77.0	57.0	0.0	-	-	-	-	-	-	-	-	-	-
77.0	60.0	-	-	-	-	-	-	-	-	-	-	-
77.0	65.0	-	-	-	-	-	-	-	-	-	-	-
80.0	50.2	2.9	-	-	-	-	-	-	-	-	-	-
80.0	51.0	9.5	-	-	-	-	-	-	-	-	-	-
80.0	52.0	9.9	-	-	-	-	-	-	-	-	-	-
80.0	55.0	6.2	-	-	-	-	-	-	-	-	-	-
80.0	60.0	0.0	-	-	-	-	-	-	-	-	-	-
80.0	65.0	0.0	-	-	-	-	-	-	-	-	-	-
80.0	70.0	0.0	-	-	-	-	-	-	-	-	-	-
80.0	75.0	0.0	-	-	-	-	-	-	-	-	-	-
											96.6	
											5.9	
											-	21.1

TABLE 4. (cont.)

*Engraulis mordax* (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
80.0	90.0	0.0	-	-	-	-	-	-	-	-	-	-
80.0	200.0	0.0	14.3	-	1748.4	-	155.0	-	-	-	-	-
82.0	47.0	-	-	-	-	-	88.9	-	-	-	-	-
83.0	39.0	0.0	-	-	-	127.8	147.3	-	-	-	25.1	-
83.0	40.0	206.3	-	-	-	2475.2	2343.6	-	-	-	61.4	-
83.0	43.0	44.4	-	-	-	1688.7	434.7	-	-	-	463.7	-
83.0	51.0	16.5	-	-	-	2177.2	162.7	-	-	-	69.4	-
83.0	55.0	168.2	-	-	-	54.0	683.1	-	-	-	474.3	-
83.0	60.0	6.1	-	-	-	2.9	210.6	-	-	-	180.9	-
83.0	65.0	5.7	-	-	-	66.2	515.6	-	-	-	2.8	-
83.0	70.0	25.5	-	-	-	-	-	-	-	-	0.0	-
83.0	75.0	-	-	-	-	-	-	-	-	-	2.9	-
83.0	80.0	0.0	-	-	-	169.3	-	16.2	-	-	0.0	-
87.0	32.5	665.2	-	-	-	322.7	-	316.7	-	-	107.9	-
87.0	33.0	4230.7	-	-	-	1404.4	-	1038.3	-	-	134.2	-
87.0	34.0	1313.6	-	-	-	1879.2	-	815.8	-	-	142.6	-
87.0	35.0	2622.5	-	-	-	1220.8	-	1252.6	-	-	329.2	-
87.0	40.0	1576.0	-	-	-	1208.9	-	1464.0	-	-	500.2	-
87.0	45.0	1009.8	-	-	-	547.2	-	368.5	-	-	691.2	-
87.0	50.0	421.6	-	-	-	644.5	-	1539.3	-	-	1075.9	-
87.0	55.0	1287.4	-	-	-	28.0	-	735.8	-	-	39.5	-
87.0	60.0	51.0	-	-	-	28.7	-	0.0	-	-	6.1	-
87.0	65.0	6.2	-	-	-	26.8	-	0.0	-	-	0.0	-
87.0	70.0	6.2	-	-	-	3.2	-	0.0	-	-	3.0	-
87.0	80.0	0.0	-	-	-	0.0	-	9.8	-	-	0.0	-
90.0	27.5	9.7	-	-	-	2159.3	-	20.5	-	-	26.0	-
90.0	28.0	404.5	-	-	-	2689.7	-	0.0	-	-	51.3	-
90.0	32.0	1044.9	-	-	-	536.8	-	-	-	-	563.5	-
90.0	37.0	2828.5	-	-	-	1171.1	-	-	-	-	4432.9	-
90.0	45.0	1856.7	-	-	-	3608.9	-	-	-	-	1354.2	-
90.0	53.0	390.7	-	-	-	640.8	-	-	-	-	45.9	-
90.0	60.0	553.2	-	-	-	310.6	-	-	-	-	3607.0	-
90.0	65.0	0.0	-	-	-	35.3	-	-	-	-	2.7	-
90.0	70.0	0.0	-	-	-	0.0	-	-	-	-	22.2	-
90.0	90.0	0.0	-	-	-	6.4	-	-	-	-	0.0	-
90.0	100.0	2.8	-	-	-	9.4	-	-	-	-	2.7	-
93.0	26.5	-	-	-	-	-	-	-	-	-	7.9	-
93.0	27.0	59.4	-	-	-	2174.0	-	-	-	-	2.2	-
93.0	28.0	407.5	-	-	-	1025.9	-	-	-	-	2.7	-
93.0	30.0	2346.9	-	-	-	889.7	-	-	-	-	1268.5	-
93.0	35.0	3450.6	-	-	-	680.9	-	-	-	-	3.7	-
93.0	40.0	593.0	-	-	-	-	-	-	-	-	0.0	-
93.0	45.0	1377.1	-	-	-	-	-	-	-	-	9.1	-
93.0	50.0	450.9	-	-	-	-	-	-	-	-	0.0	-
93.0	55.0	202.6	-	-	-	-	-	-	-	-	2.9	-
93.0	60.0	488.0	-	-	-	-	-	-	-	-	33.3	-

TABLE 4. (cont.)

*Engraulis mordax* (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
93.0	65.0	37.8	-	-	2050.2	-	0.0	-	-	0.0	-	-
93.0	70.0	0.0	-	-	6773.2	-	1165.1	-	-	0.0	-	-
93.0	75.0	-	-	-	-	557.9	-	6.3	-	5.7	-	-
93.0	80.0	0.0	-	-	-	6.7	-	0.0	-	0.0	-	-
93.0	90.0	3.2	-	-	-	-	-	0.0	-	15.2	-	-
96.8	28.7	705.6	-	14.8	-	-	-	25.6	-	31.7	-	-
97.0	29.0	2941.5	-	324.3	-	-	-	2.8	-	22.7	-	-
97.0	30.0	2505.6	-	533.8	-	-	-	3.4	-	6.2	-	-
97.0	32.0	5447.5	-	2511.0	-	-	-	3025.8	-	61.1	-	-
97.0	35.0	3027.0	-	1313.6	-	-	-	-	-	15.1	-	-
97.0	40.0	3046.1	-	245.6	-	-	-	-	-	12.2	-	-
97.0	45.0	6602.9	-	828.0	-	-	-	6.4	-	0.0	-	-
97.0	50.0	4347.0	-	489.4	-	-	-	3.3	-	0.0	-	-
97.0	55.0	546.0	-	-	-	-	-	0.0	-	0.0	-	-
97.0	60.0	896.9	-	-	-	-	-	0.0	-	0.0	-	-
97.0	65.0	1817.4	-	1373.1	-	-	-	0.0	-	0.0	-	-
97.0	70.0	1177.1	-	-	2596.8	-	0.0	-	-	0.0	-	-
97.0	75.0	-	-	-	-	32.2	-	-	-	6.5	-	-
97.0	80.0	6.7	-	-	-	282.3	-	-	-	2.9	-	-
97.0	85.0	-	-	-	-	606.7	-	-	-	0.0	-	-
100.0	29.0	1563.8	-	-	-	110.3	-	-	-	4.2	-	-
100.0	30.0	761.5	-	-	-	32.2	-	-	-	65.5	-	-
100.0	35.0	2867.4	-	-	-	31.8	-	-	-	8.3	-	-
100.0	40.0	1452.4	-	-	-	544.8	-	-	-	2.8	-	-
100.0	45.0	1480.4	-	-	-	359.4	-	-	-	0.0	-	-
100.0	50.0	957.0	-	-	-	134.0	-	-	-	0.0	-	-
100.0	55.0	1833.2	-	-	-	1209.0	-	-	-	2.9	-	-
100.0	60.0	524.5	-	-	-	2077.5	-	-	-	3.1	-	-
100.0	65.0	512.6	-	-	-	485.3	-	-	-	9.2	-	-
100.0	70.0	1059.5	-	-	-	-	-	-	-	4.7	-	-
100.0	75.0	-	-	-	-	-	-	-	-	0.0	-	-
100.0	80.0	40.7	-	-	-	-	-	-	-	7.0	-	-
103.0	28.5	206.3	-	-	-	-	-	-	-	34.2	-	-
103.0	29.0	75.6	-	-	-	-	-	-	-	95.8	-	-
103.0	30.0	327.2	-	-	-	-	-	-	-	0.0	-	-
103.0	35.0	3607.4	-	-	-	-	-	-	-	0.0	-	-
103.0	40.0	633.6	-	-	-	-	-	-	-	0.0	-	-
103.0	45.0	44.3	-	-	-	-	-	-	-	0.0	-	-
103.0	50.0	14.6	-	-	-	-	-	-	-	0.0	-	-
103.0	55.0	146.4	-	-	-	-	-	-	-	0.0	-	-
103.0	60.0	282.7	-	-	-	-	-	-	-	0.0	-	-
103.0	65.0	758.4	-	-	-	-	-	-	-	0.0	-	-
103.0	70.0	635.1	-	-	-	-	-	-	-	0.0	-	-
103.0	75.0	-	-	-	-	-	-	-	-	0.0	-	-
103.0	80.0	-	-	-	-	-	-	-	-	0.0	-	-
103.0	90.0	-	-	-	-	-	-	-	-	0.0	-	-

TABLE 4. (cont.)

*Engraulis mordax* (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
107.0	30.0	-	57.0	-	38.3	-	1.1	-	-	0.0	-	-
107.0	31.0	-	236.5	-	237.7	-	3.8	-	-	7.1	-	-
107.0	32.0	-	889.9	-	930.2	-	0.0	-	-	0.0	-	-
107.0	35.0	-	601.6	-	308.7	-	3.5	-	-	0.0	-	-
107.0	40.0	-	514.5	-	39.0	-	30.0	-	-	0.0	-	-
107.0	50.0	-	23.4	-	0.0	-	0.0	-	-	0.0	-	-
107.0	55.0	-	14.8	-	0.0	-	0.0	-	-	0.0	-	-
107.0	60.0	-	351.5	-	0.0	-	0.0	-	-	0.0	-	-
107.0	65.0	-	2.9	-	0.0	-	0.0	-	-	0.0	-	-
107.0	70.0	-	6.4	-	0.0	-	0.0	-	-	0.0	-	-
110.0	32.0	-	172.2	-	16.9	-	-	-	-	0.0	-	-
110.0	35.0	-	973.5	-	0.0	-	-	-	-	2.8	-	-
110.0	40.0	-	2182.8	-	11.4	-	-	-	-	0.0	-	-
110.0	45.0	-	2.8	-	3.2	-	-	-	-	0.0	-	-
110.0	55.0	-	2.7	-	0.0	-	-	-	-	5.7	-	-
110.0	60.0	-	0.0	-	0.0	-	-	-	-	2.8	-	-
110.0	65.0	-	5.3	-	0.0	-	-	-	-	0.0	-	-
110.0	70.0	-	14.0	-	0.0	-	-	-	-	0.0	-	-
113.0	28.5	-	34.3	-	2.2	-	12.9	-	-	5.6	-	-
113.0	29.0	-	92.5	-	3.7	-	7.2	-	-	112.1	-	-
113.0	30.0	-	54.6	-	5.6	-	56.6	-	-	427.4	-	-
113.0	35.0	-	6.0	-	0.0	-	3.3	-	-	8.0	-	-
113.0	40.0	-	11.7	-	0.0	-	0.0	-	-	3.0	-	-
113.0	45.0	-	271.9	-	0.0	-	-	-	-	0.0	-	-
113.0	50.0	-	27.4	-	0.0	-	-	-	-	0.0	-	-
113.0	60.0	-	35.3	-	0.0	-	-	-	-	0.0	-	-
113.0	65.0	-	3.1	-	0.0	-	-	-	-	-	-	-
113.0	70.0	-	2.7	-	24.6	-	0.0	-	-	-	-	-
113.0	75.0	-	-	-	3.3	-	-	-	-	-	-	-
113.0	80.0	-	0.0	-	15.4	-	0.0	-	-	10.4	-	-
115.0	35.0	-	507.3	-	-	-	-	-	-	9.4	-	-
117.0	24.8	-	16.8	-	30.0	-	-	-	-	466.0	-	-
117.0	25.0	-	237.5	-	399.0	-	-	-	-	899.2	-	-
117.0	26.0	-	918.4	-	484.6	-	-	-	-	245.6	-	-
117.0	30.0	-	869.9	-	97.8	-	-	-	-	35.3	-	-
117.0	35.0	-	147.6	-	780.0	-	-	-	-	0.0	-	-
117.0	40.0	-	198.3	-	128.5	-	-	-	-	0.0	-	-
117.0	45.0	-	719.3	-	2.9	-	-	-	-	33.6	-	-
117.0	50.0	-	371.2	-	247.4	-	-	-	-	0.0	-	-
117.0	55.0	-	0.0	-	32.0	-	-	-	-	72.6	-	-
117.0	60.0	-	0.0	-	0.0	-	-	-	-	0.0	-	-
117.0	75.0	-	-	-	9.1	-	-	-	-	-	-	-
117.0	90.0	-	0.0	-	0.0	-	-	-	-	3.3	-	-
118.0	39.0	-	235.6	-	-	-	-	-	-	42.0	-	-
119.0	33.0	-	1729.0	-	49.3	-	-	-	-	-	316.8	-
120.0	22.5	-	3.3	-	-	-	-	-	-	6.2	-	-

TABLE 4. (cont.)

*Engraulis mordax* (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
120.0	23.0	-	0.0	-	56.4	-	-	0.0	-	7.5	-	-
120.0	24.0	-	1635.1	-	470.1	-	-	158.9	-	83.3	-	-
120.0	25.0	-	757.6	-	405.5	-	-	910.0	-	51.3	-	-
120.0	30.0	-	1670.4	-	783.8	-	-	565.8	-	183.6	-	-
120.0	35.0	-	503.3	-	823.0	-	-	850.2	-	57.5	-	-
120.0	40.0	-	18.6	-	387.6	-	-	967.5	-	2.7	-	-
120.0	45.0	-	40.3	-	24.7	-	-	-	5.7	3.0	-	-
120.0	50.0	-	55.3	-	106.9	-	-	-	0.0	0.0	-	-
120.0	55.0	-	2.8	-	83.4	-	-	-	0.0	0.0	-	-
120.0	60.0	-	2.9	-	5.8	-	-	-	0.0	0.0	-	-
120.0	65.0	-	5.6	-	0.0	-	-	-	0.0	-	-	-
120.0	70.0	-	2.9	-	0.0	-	-	-	0.0	-	-	-
120.0	70.0	-	0.0	-	6.9	-	-	-	0.0	-	-	-
120.0	80.0	-	37.4	-	19.3	-	-	70.4	-	3.2	-	-
123.0	36.0	-	123.0	-	22.1	-	-	36.0	-	0.0	-	-
123.0	37.0	-	85.8	-	2230.8	-	-	0.0	110.3	0.0	-	-
123.0	42.0	-	1350.3	-	4877.2	-	-	16.6	-	0.0	-	-
123.0	45.0	-	756.6	-	0.0	-	-	0.0	-	0.0	-	-
123.0	50.0	-	8.8	-	0.0	-	-	5.7	-	0.0	-	-
123.0	55.0	-	3.0	-	0.0	-	-	153.7	-	1.4	-	-
123.5	35.6	-	257.5	-	25.6	-	-	6.8	-	57.4	-	-
127.0	32.6	-	618.3	-	43.2	-	-	8.9	-	5.2	-	-
127.0	33.0	-	4522.9	-	169.5	-	-	50.8	-	0.0	-	-
127.0	34.0	-	1964.6	-	65.9	-	-	39.8	-	2.7	-	-
127.0	40.0	-	619.8	-	1093.0	-	-	67.8	-	0.0	-	-
127.0	45.0	-	0.0	-	0.0	-	-	-	1601.4	255.4	-	-
130.0	25.5	-	34.0	-	12.7	-	-	-	-	171.0	171.7	-
130.0	26.0	-	66.2	-	98.8	-	-	-	-	267.8	4.7	-
130.0	28.0	-	466.5	-	365.9	-	-	-	-	117.2	18.1	-
130.0	30.0	-	209.0	-	14.3	-	-	-	-	198.9	0.0	-
130.0	35.0	-	855.8	-	1088.9	-	-	-	-	0.0	0.0	-
130.0	40.0	-	3.1	-	5131.4	-	-	-	-	0.0	0.0	-
130.0	45.0	-	58.6	-	82.1	-	-	-	-	0.0	0.0	-
130.0	50.0	-	167.0	-	167.0	-	-	-	-	0.0	0.0	-
130.0	55.0	-	0.0	-	2.9	-	-	-	-	0.0	0.0	-
130.0	70.0	-	0.0	-	3.1	-	-	-	-	0.0	0.0	-
133.0	19.5	-	6.8	-	29.9	-	-	3.3	-	258.9	-	-
133.0	21.0	-	232.0	-	194.0	-	-	4.0	-	297.7	-	-
133.0	23.0	-	5777.0	-	1322.9	-	-	2.5	-	47.5	-	-
133.0	25.0	-	5375.7	-	577.6	-	-	24.2	-	11.8	-	-
133.0	30.0	-	2240.9	-	353.4	-	-	352.2	-	11.4	-	-
133.0	35.0	-	858.2	-	9.3	-	-	53.3	-	0.0	-	-
133.0	40.0	-	795.7	-	9.2	-	-	93.1	-	0.0	-	-
133.0	45.0	-	4523.9	-	6.0	-	-	2.7	-	0.0	-	-
133.0	50.0	-	662.2	-	0.0	-	-	5.2	-	0.0	-	-
133.0	55.0	-	11.0	-	0.0	-	-	0.0	-	426.3	56.4	-
	20.0	-	6.2	-	-	-	-	-	-	534.5	-	-

TABLE 4. (cont.)

*Engraulis mordax* (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
137.0	21.0	-	42.2	-	244.2	-	-	-	1715.2	0.0	-	-
137.0	22.0	-	1587.3	-	1296.2	-	-	-	436.6	0.0	-	-
137.0	23.0	-	124.0	-	1267.2	-	-	-	195.7	0.0	-	-
137.0	30.0	-	268.0	-	183.5	-	-	-	140.8	3.0	-	-
137.0	35.0	-	37.3	-	12.0	-	-	-	0.0	0.0	-	-
137.0	40.0	-	227.9	-	15.0	-	-	-	0.0	0.0	-	-
137.0	45.0	-	17.4	-	3.0	-	-	-	0.0	3.0	-	-
137.0	55.0	-	0.0	-	0.0	-	-	-	2.8	-	-	-
140.0	28.8	-	230.5	-	-	-	-	-	2.4	-	-	-
140.0	29.0	-	704.3	-	-	-	-	-	9.8	-	-	-
140.0	30.0	-	1512.2	-	-	-	-	-	2.8	-	-	-
140.0	35.0	-	60.6	-	-	-	-	-	0.0	-	-	-

*Argentina sialis*

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
83.0	43.0	5.9	-	-	0.0	-	-	9.5	-	-	0.0	-
83.0	55.0	0.0	-	-	0.0	-	-	10.0	-	-	0.0	-
87.0	35.0	2.9	-	-	0.0	-	-	0.0	-	-	0.0	-
90.0	28.0	13.9	-	-	0.0	-	-	-	0.0	-	0.0	-
90.0	32.0	5.4	-	-	0.0	-	-	-	0.0	-	0.0	-
93.0	27.0	0.0	-	-	0.0	-	-	-	8.6	-	0.0	-
93.0	28.0	15.4	-	-	0.0	-	-	-	3.2	-	0.0	-
93.0	30.0	3.0	-	-	0.0	-	-	-	3.1	-	0.0	-
93.0	40.0	0.0	-	-	-	3.3	-	-	6.3	-	0.0	-
97.0	29.0	0.0	-	-	0.0	-	-	-	2.6	-	0.0	-
97.0	30.0	2.2	-	-	0.0	-	-	-	5.6	-	0.0	-
97.0	32.0	0.0	-	-	0.0	-	-	-	10.1	-	0.0	-
97.0	40.0	3.2	-	-	0.0	-	-	-	0.0	-	0.0	-
100.0	55.0	3.0	-	-	0.0	-	-	-	0.0	-	0.0	-
103.0	29.0	0.0	-	-	-	1.6	-	-	-	-	0.0	-
107.0	31.0	-	-	-	-	2.2	-	-	-	-	0.0	-
107.0	32.0	-	-	-	-	27.8	-	-	-	-	3.4	-
113.0	30.0	-	-	-	0.0	-	-	-	0.0	-	1.6	-
113.0	45.0	-	-	-	0.0	-	-	-	9.2	-	0.0	-
113.0	60.0	-	-	-	0.0	-	-	-	3.4	-	0.0	-
115.0	35.0	-	-	-	-	2.8	-	-	-	-	2.6	-
117.0	25.0	-	-	-	-	2.6	-	-	-	-	0.0	-
117.0	30.0	-	-	-	-	14.5	-	-	-	-	2.9	-
117.0	35.0	-	-	-	-	3.1	-	-	-	-	2.9	-
117.0	40.0	-	-	-	-	6.1	-	-	-	-	2.6	-
117.0	45.0	-	-	-	-	3.0	-	-	-	-	3.4	-
117.0	50.0	-	-	-	-	3.2	-	-	-	-	0.0	-
118.0	39.0	-	-	-	-	0.0	-	-	-	-	22.4	-
119.0	33.0	-	-	-	-	-	-	-	-	-	2.2	-

TABLE 4. (cont.)

*Argentina sialis* (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
127.0	50.0	-	2.9	-	0.0	-	-	0.0	-	0.0	-	-
133.0	30.0	-	0.0	-	3.1	-	-	0.0	-	0.0	-	-
133.0	45.0	-	2.9	-	0.0	-	-	0.0	-	0.0	-	-

*Microstoma microstoma*

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
70.0	90.0	0.0	-	-	-	-	-	2.8	-	-	-	-
80.0	65.0	0.0	-	-	0.0	-	-	3.0	3.0	-	-	-
80.0	75.0	-	-	-	-	-	-	-	-	-	-	-
80.0	80.0	3.0	-	-	0.0	-	-	0.0	-	2.7	-	-
87.0	70.0	3.1	-	-	0.0	-	-	0.0	-	0.0	-	-
90.0	70.0	0.0	-	-	0.0	-	-	0.0	-	0.0	-	-
90.0	80.0	0.0	-	-	6.4	-	-	0.0	-	8.9	-	-
90.0	100.0	0.0	-	-	3.1	-	-	0.0	-	0.0	-	-
90.0	100.0	0.0	-	-	0.0	-	-	-	-	-	-	-
93.0	30.0	0.0	-	0.0	-	-	-	3.1	-	0.0	-	-
97.0	40.0	0.0	-	0.0	-	-	-	3.3	-	0.0	-	-
100.0	50.0	0.0	-	-	3.0	-	-	0.0	-	0.0	-	-
100.0	70.0	3.1	-	-	6.0	-	-	0.0	-	0.0	-	-
100.0	90.0	3.0	-	-	0.0	-	-	0.0	-	0.0	-	-
103.0	65.0	0.0	-	-	3.2	-	-	0.0	-	0.0	-	-
103.0	80.0	-	3.0	-	0.0	-	-	0.0	-	0.0	-	-
107.0	70.0	-	3.2	-	0.0	-	-	0.0	-	0.0	-	-

*Nansenia candida*

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
70.0	60.0	0.0	-	-	11.4	-	-	0.0	-	-	-	-
70.0	90.0	0.0	-	-	-	-	-	2.8	-	-	-	-
70.0	100.0	0.0	-	-	-	-	-	5.6	-	-	-	-
73.0	65.0	-	-	-	2.9	-	-	-	-	-	-	-
80.0	65.0	3.1	-	-	0.0	-	-	0.0	0.0	0.0	-	-
80.0	80.0	0.0	-	-	3.2	-	-	0.0	-	0.0	-	-
83.0	80.0	0.0	-	-	2.9	-	-	0.0	-	0.0	-	-
87.0	70.0	0.0	-	-	0.0	-	-	3.0	-	0.0	-	-
93.0	70.0	0.0	-	-	0.0	-	-	3.3	-	0.0	-	-

*Nansenia crassa*

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
100.0	35.0	0.0	-	-	0.0	-	-	0.0	-	3.0	-	-
107.0	35.0	-	0.0	-	3.2	-	-	0.0	-	0.0	-	-
107.0	75.0	-	-	-	3.1	-	-	-	-	-	-	-

TABLE 4. (cont.)

*Nansenia crassa* (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
107.0	80.0	-	2.9	-	0.0	-	0.0	-	-	-	-	-
113.0	70.0	-	0.0	-	3.1	-	0.0	-	-	-	-	-
117.0	60.0	-	0.0	-	0.0	-	3.8	-	-	-	-	-
117.0	80.0	-	0.0	-	3.1	-	0.0	-	-	-	-	-
117.0	85.0	-	-	-	2.9	-	-	-	-	-	-	-
123.0	42.0	-	3.3	-	0.0	-	-	0.0	0.0	0.0	-	-
127.0	40.0	-	1.4	-	0.0	-	-	0.0	0.0	0.0	-	-
127.0	45.0	-	6.0	-	3.1	-	-	0.0	-	0.0	-	-
127.0	50.0	-	5.9	-	0.0	-	-	0.0	-	0.0	-	-
130.0	40.0	-	6.1	-	0.0	-	-	-	2.6	0.0	-	-
130.0	45.0	-	5.6	-	0.0	-	-	-	0.0	-	-	-
130.0	50.0	-	20.2	-	0.0	-	-	-	0.0	3.3	-	-
130.0	55.0	-	0.0	-	0.0	-	-	-	2.7	0.0	-	-
133.0	45.0	-	5.9	-	0.0	-	-	0.0	-	0.0	-	-
133.0	60.0	-	0.0	-	0.0	-	-	-	2.8	-	0.0	-
137.0	45.0	-	2.9	-	0.0	-	-	-	0.0	-	-	-

*Bathylagus* spp.

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
67.0	55.0	10.0	-	-	0.0	-	-	0.0	-	-	-	-
77.0	57.0	3.0	-	-	0.0	-	-	0.0	-	-	-	-
80.0	60.0	7.7	-	-	0.0	-	-	0.0	0.0	-	-	-
83.0	43.0	0.0	-	-	2.8	-	-	0.0	-	0.0	-	-
93.0	30.0	0.0	-	8.6	-	-	-	0.0	-	0.0	-	-
118.0	39.0	-	0.0	-	0.0	-	-	0.0	-	2.7	-	-
123.0	50.0	-	0.0	-	0.0	-	-	5.7	-	0.0	-	-

*Bathylagus milleri*

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
67.0	55.0	3.3	-	-	0.0	-	-	0.0	-	-	-	-

*Bathylagus ochotensis*

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
60.0	60.0	0.0	-	-	-	-	-	-	-	-	-	-
60.0	70.0	3.2	-	-	-	-	-	-	-	2.8	-	-
60.0	80.0	5.1	-	-	-	-	-	-	-	0.0	-	-
60.0	100.0	0.0	-	-	-	-	-	-	-	2.6	-	-
63.0	55.0	22.2	-	-	-	-	-	-	-	2.8	-	-
63.0	60.0	13.0	-	-	-	-	-	-	-	0.0	-	-
63.0	65.0	-	-	-	-	-	-	-	-	17.5	-	-

TABLE 4. (cont.)

*Bathyergus ochotensis* (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
63.0	70.0	-	-	-	25.0	-	-	-	-	-	-	-
63.0	80.0	-	-	-	38.1	-	-	-	-	-	-	-
67.0	49.0	0.0	-	-	2.8	-	-	-	-	-	-	-
67.0	50.0	0.0	-	-	2.7	-	-	-	-	-	-	-
67.0	55.0	0.0	-	-	13.5	-	-	-	-	-	-	-
67.0	60.0	3.0	-	-	11.0	-	-	-	-	-	-	-
67.0	65.0	-	-	-	34.9	-	-	-	-	-	-	-
67.0	80.0	-	-	-	71.6	-	-	-	-	-	-	-
70.0	50.0	2.9	-	-	4.5	-	-	-	-	-	-	-
70.0	51.0	3.1	-	-	5.5	-	-	-	-	-	-	-
70.0	52.0	31.5	-	-	28.4	-	-	-	-	-	-	-
70.0	53.0	11.8	-	-	45.9	-	-	-	-	-	-	-
70.0	55.0	2.9	-	-	-	-	-	-	-	-	-	-
70.0	60.0	2.9	-	-	37.2	-	-	-	-	-	-	-
70.0	65.0	-	-	-	16.1	-	-	-	-	-	-	-
70.0	70.0	13.6	-	-	14.5	-	-	-	-	-	-	-
70.0	120.0	3.0	-	-	-	-	-	-	-	-	-	-
73.0	50.0	2.3	-	-	2.8	-	-	-	-	-	-	-
73.0	51.0	15.4	-	-	2.8	-	-	-	-	-	-	-
73.0	53.0	41.0	-	-	9.0	-	-	-	-	-	-	-
73.0	60.0	13.9	-	-	2.8	-	-	-	-	-	-	-
73.0	65.0	-	-	-	2.9	-	-	-	-	-	-	-
73.0	70.0	-	-	-	2.8	-	-	-	-	-	-	-
73.0	80.0	-	-	-	5.2	-	-	-	-	-	-	-
77.0	51.0	-	-	-	0.0	-	-	-	-	-	-	-
77.0	55.0	5.9	-	-	3.6	-	-	-	-	-	-	-
77.0	57.0	0.0	-	-	7.4	-	-	-	-	-	-	-
77.0	65.0	-	-	-	15.3	-	-	-	-	-	-	-
77.0	70.0	-	-	-	9.2	-	-	-	-	-	-	-
80.0	50.2	0.7	-	-	0.0	-	-	-	-	-	-	-
80.0	52.0	3.3	-	-	0.0	-	-	-	-	-	-	-
80.0	55.0	12.4	-	-	0.0	-	-	-	-	-	-	-
80.0	60.0	25.7	-	-	0.0	-	-	-	-	-	-	-
80.0	65.0	3.1	-	-	20.2	-	-	-	-	-	-	-
80.0	70.0	0.0	-	-	30.6	-	-	-	-	-	-	-
80.0	80.0	0.0	-	-	9.7	-	-	-	-	-	-	-
80.0	120.0	3.0	-	-	-	-	-	-	-	-	-	-
83.0	55.0	5.8	-	-	0.0	-	-	-	-	-	-	-
83.0	60.0	3.0	-	-	0.0	-	-	-	-	-	-	-
83.0	65.0	2.9	-	-	0.0	-	-	-	-	-	-	-
83.0	70.0	-	-	-	0.0	-	-	-	-	-	-	-
83.0	80.0	3.2	-	-	0.0	-	-	-	-	-	-	-
83.0	80.0	7.1	-	-	0.0	-	-	-	-	-	-	-
87.0	32.5	1.6	-	-	0.0	-	-	-	-	-	-	-
87.0	33.0	8.0	-	-	0.0	-	-	-	-	-	-	-
87.0	34.0	2.7	-	-	0.0	-	-	-	-	-	-	-
87.0	35.0	41.2	-	-	0.0	-	-	-	-	-	-	-

TABLE 4. (cont.)

*Bathyergus ochotensis* (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
87.0	40.0	14.3	-	-	0.0	-	-	-	-	-	0.0	-
87.0	55.0	6.0	-	-	0.0	-	-	-	-	-	0.0	-
87.0	70.0	3.1	-	-	0.0	-	-	-	-	-	0.0	-
87.0	80.0	12.1	-	-	3.0	-	-	-	-	-	0.0	-
87.0	90.0	0.0	-	-	2.8	-	-	-	-	-	0.0	-
90.0	28.0	2.8	-	-	0.0	-	-	-	-	-	0.0	-
90.0	32.0	2.7	-	-	0.0	-	-	-	-	-	0.0	-
90.0	37.0	3.0	-	-	0.0	-	-	-	-	-	0.0	-
90.0	53.0	16.1	-	-	0.0	-	-	-	-	-	0.0	-
90.0	60.0	0.0	-	-	3.3	-	-	-	-	-	0.0	-
90.0	65.0	0.0	-	-	2.9	-	-	-	-	-	0.0	-
90.0	70.0	3.1	-	-	0.0	-	-	-	-	-	0.0	-
90.0	80.0	2.7	-	-	0.0	-	-	-	-	-	0.0	-
93.0	30.0	12.2	-	-	0.0	-	-	-	-	-	0.0	-
93.0	35.0	21.0	-	-	0.0	-	-	-	-	-	0.0	-
93.0	40.0	12.0	-	-	0.0	-	-	-	-	-	3.2	-
93.0	45.0	6.1	-	-	0.0	-	-	-	-	-	0.0	-
93.0	50.0	3.1	-	-	0.0	-	-	-	-	-	0.0	-
93.0	55.0	0.0	-	-	4.2	-	-	-	-	-	3.3	-
93.0	70.0	0.0	-	-	0.0	-	-	-	-	-	6.6	-
97.0	29.0	1.9	-	-	0.0	-	-	-	-	-	0.0	-
97.0	30.0	2.2	-	-	0.0	-	-	-	-	-	0.0	-
97.0	32.0	6.2	-	-	0.0	-	-	-	-	-	0.0	-
97.0	40.0	0.0	-	-	0.0	-	-	-	-	-	0.0	-
97.0	45.0	6.1	-	-	0.0	-	-	-	-	-	0.0	-
97.0	50.0	15.8	-	-	0.0	-	-	-	-	-	0.0	-
97.0	55.0	3.0	-	-	0.0	-	-	-	-	-	0.0	-
97.0	65.0	15.3	-	-	0.0	-	-	-	-	-	0.0	-
97.0	70.0	15.9	-	-	-	-	-	-	-	-	0.0	-
100.0	55.0	3.0	-	-	-	-	-	-	-	-	0.0	-
100.0	60.0	3.0	-	-	-	-	-	-	-	-	0.0	-
100.0	65.0	3.0	-	-	-	-	-	-	-	-	0.0	-
100.0	70.0	27.7	-	-	-	-	-	-	-	-	0.0	-
103.0	55.0	5.7	-	-	-	-	-	-	-	-	0.0	-
103.0	70.0	3.0	-	-	-	-	-	-	-	-	0.0	-
103.0	85.0	-	-	-	-	-	-	-	-	-	2.9	-
107.0	31.0	-	-	-	-	-	-	-	-	-	0.0	-
107.0	32.0	-	-	-	-	-	-	-	-	-	0.0	-
107.0	35.0	-	-	-	-	-	-	-	-	-	3.2	-
107.0	40.0	-	-	-	-	-	-	-	-	-	0.0	-
107.0	60.0	-	-	-	-	-	-	-	-	-	0.0	-
110.0	65.0	-	-	-	-	-	-	-	-	-	6.1	-
												2.7

TABLE 4. (cont.)

*Bathylagus pacificus*

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
60.0	70.0	3.2	-	-	-	-	-	0.0	-	-	-	-
63.0	80.0	-	-	-	5.9	-	-	-	-	-	-	-
80.0	60.0	2.6	-	-	0.0	-	-	0.0	0.0	-	-	-

*Bathylagus wesethi*

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
60.0	140.0	0.0	-	-	-	-	-	28.7	-	-	-	-
60.0	160.0	0.0	-	-	-	-	-	11.4	-	-	-	-
60.0	180.0	0.0	-	-	0.0	-	-	8.1	-	-	-	-
70.0	53.0	0.0	-	-	-	-	-	2.9	-	-	-	-
70.0	90.0	0.0	-	-	-	-	-	11.0	-	-	-	-
70.0	100.0	0.0	-	-	-	-	-	13.9	-	-	-	-
70.0	120.0	0.0	-	-	-	-	-	11.0	-	-	-	-
70.0	160.0	3.0	-	-	-	-	-	-	-	-	-	-
73.0	60.0	0.0	-	-	0.0	-	-	5.6	-	-	-	-
80.0	60.0	0.0	-	-	0.0	-	-	16.8	0.0	-	-	-
80.0	65.0	0.0	-	-	0.0	-	-	8.9	0.0	-	-	-
80.0	70.0	0.0	-	-	0.0	-	-	12.3	0.0	-	-	-
80.0	80.0	0.0	-	-	0.0	-	-	5.9	-	0.0	-	-
80.0	90.0	0.0	-	-	-	-	-	11.2	-	-	-	-
80.0	100.0	0.0	-	-	-	-	-	3.0	-	-	-	-
80.0	120.0	0.0	-	-	-	-	-	26.6	-	-	-	-
83.0	90.0	0.0	-	-	-	-	-	-	-	-	-	-
87.0	55.0	0.0	-	-	0.0	-	-	22.5	-	-	-	-
87.0	60.0	0.0	-	-	0.0	-	-	3.3	-	0.0	-	-
87.0	65.0	0.0	-	-	0.0	-	-	32.0	-	0.0	-	-
87.0	70.0	0.0	-	-	28.9	-	-	30.4	-	0.0	-	-
87.0	80.0	0.0	-	-	9.0	-	-	88.0	-	0.0	-	-
87.0	90.0	0.0	-	-	1.1.3	-	-	37.4	-	-	-	-
87.0	100.0	-	-	-	31.2	-	-	-	-	-	-	-
90.0	60.0	0.0	-	-	3.3	-	-	-	-	-	-	-
90.0	65.0	0.0	-	-	11.8	-	-	0.0	-	-	-	-
90.0	70.0	0.0	-	-	65.3	-	-	2.7	-	-	-	-
90.0	80.0	0.0	-	-	80.5	-	-	5.2	-	-	-	-
90.0	90.0	0.0	-	-	19.2	-	-	2.3	-	-	-	-
90.0	100.0	0.0	-	-	0.0	-	-	24.7	-	-	-	-
90.0	120.0	0.0	-	-	-	-	-	2.7	-	-	-	-
90.0	160.0	0.0	-	-	-	-	-	5.2	-	-	-	-
93.0	50.0	0.0	-	-	0.0	-	-	6.6	-	0.0	-	-
93.0	55.0	0.0	-	-	0.0	-	-	3.3	-	0.0	-	-
93.0	60.0	0.0	-	-	0.0	-	-	3.3	-	0.0	-	-
93.0	65.0	0.0	-	-	0.0	-	-	6.5	-	0.0	-	-
93.0	70.0	0.0	-	-	0.0	-	-	6.6	-	0.0	-	-
93.0	80.0	0.0	-	-	50.7	-	-	15.8	-	32.3	-	-

TABLE 4. (cont.)

*Bathylagus wesethi* (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
93.0	85.0	-	-	-	-	-	-	-	-	-	-	-
93.0	90.0	0.0	-	-	-	-	-	-	-	-	-	-
93.0	100.0	-	-	-	-	-	-	-	-	-	-	-
97.0	29.0	0.0	0.0	-	-	-	-	-	-	-	-	-
97.0	35.0	0.0	0.0	-	-	-	-	-	-	-	-	-
97.0	40.0	0.0	0.0	-	-	-	-	-	-	-	-	-
97.0	45.0	0.0	0.0	-	-	-	-	-	-	-	-	-
97.0	50.0	0.0	0.0	-	-	-	-	-	-	-	-	-
97.0	55.0	0.0	0.0	-	-	-	-	-	-	-	-	-
97.0	60.0	0.0	0.0	-	-	-	-	-	-	-	-	-
97.0	65.0	0.0	0.0	-	-	-	-	-	-	-	-	-
97.0	70.0	0.0	0.0	-	-	-	-	-	-	-	-	-
97.0	75.0	-	-	-	-	-	-	-	-	-	-	-
97.0	80.0	3.3	-	-	-	-	-	-	-	-	-	-
97.0	85.0	-	-	-	-	-	-	-	-	-	-	-
97.0	90.0	0.0	-	-	-	-	-	-	-	-	-	-
97.0	95.0	-	-	-	-	-	-	-	-	-	-	-
97.0	100.0	-	-	-	-	-	-	-	-	-	-	-
100.0	30.0	0.0	0.0	-	-	-	-	-	-	-	-	-
100.0	35.0	0.0	0.0	-	-	-	-	-	-	-	-	-
100.0	40.0	0.0	0.0	-	-	-	-	-	-	-	-	-
100.0	45.0	0.0	0.0	-	-	-	-	-	-	-	-	-
100.0	50.0	0.0	0.0	-	-	-	-	-	-	-	-	-
100.0	55.0	0.0	0.0	-	-	-	-	-	-	-	-	-
100.0	60.0	0.0	0.0	-	-	-	-	-	-	-	-	-
100.0	65.0	0.0	0.0	-	-	-	-	-	-	-	-	-
100.0	70.0	0.0	0.0	-	-	-	-	-	-	-	-	-
100.0	75.0	-	-	-	-	-	-	-	-	-	-	-
100.0	80.0	0.0	0.0	-	-	-	-	-	-	-	-	-
100.0	85.0	-	-	-	-	-	-	-	-	-	-	-
100.0	90.0	0.0	0.0	-	-	-	-	-	-	-	-	-
100.0	100.0	-	-	-	-	-	-	-	-	-	-	-
103.0	45.0	0.0	0.0	-	-	-	-	-	-	-	-	-
103.0	50.0	0.0	0.0	-	-	-	-	-	-	-	-	-
103.0	55.0	0.0	0.0	-	-	-	-	-	-	-	-	-
103.0	60.0	0.0	0.0	-	-	-	-	-	-	-	-	-
103.0	65.0	0.0	0.0	-	-	-	-	-	-	-	-	-
103.0	70.0	0.0	0.0	-	-	-	-	-	-	-	-	-
103.0	75.0	-	-	-	-	-	-	-	-	-	-	-
103.0	80.0	-	-	-	-	-	-	-	-	-	-	-
103.0	85.0	-	-	-	-	-	-	-	-	-	-	-
103.0	90.0	-	-	-	-	-	-	-	-	-	-	-
105.0	140.0	-	-	-	-	-	-	-	-	-	-	-
107.0	45.0	-	-	-	-	-	-	-	-	-	-	-
107.0	50.0	-	-	-	-	-	-	-	-	-	-	-
107.0	55.0	-	-	-	-	-	-	-	-	-	-	-

TABLE 4. (cont.)

*Bathyergus wesethi* (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
107.0	60.0	-	0.0	-	0.0	-	0.0	-	-	3.0	-	-
107.0	65.0	-	0.0	-	3.1	-	15.9	-	-	23.0	-	-
107.0	70.0	-	0.0	-	3.1	-	3.1	-	-	33.3	-	-
107.0	75.0	-	-	-	9.2	-	-	-	-	-	-	-
107.0	80.0	-	2.9	-	0.0	-	35.4	-	-	-	-	-
107.0	85.0	-	-	-	2.9	-	-	-	-	-	-	-
107.0	90.0	-	0.0	-	58.1	0.0	-	-	1.5	0.0	-	-
110.0	32.0	-	0.0	-	0.0	-	-	-	22.8	0.0	-	-
110.0	35.0	-	0.0	-	0.0	-	-	-	47.9	0.0	-	-
110.0	40.0	-	0.0	-	0.0	-	-	-	5.1	6.7	-	-
110.0	45.0	-	0.0	-	0.0	-	-	-	0.0	0.0	-	-
110.0	50.0	-	0.0	-	0.0	-	-	-	0.0	0.0	-	-
110.0	55.0	-	0.0	-	0.0	-	-	-	0.0	15.3	-	-
110.0	60.0	-	0.0	-	0.0	-	-	-	0.0	3.0	-	-
110.0	65.0	-	0.0	-	0.0	-	-	-	0.0	3.1	-	-
110.0	75.0	-	-	-	3.2	-	-	-	-	-	-	-
110.0	80.0	-	0.0	-	5.6	-	-	-	-	-	-	-
110.0	90.0	-	0.0	-	3.0	-	-	-	-	-	-	-
113.0	40.0	-	0.0	-	0.0	-	-	-	-	-	-	-
113.0	45.0	-	0.0	-	0.0	-	-	-	-	-	-	-
113.0	50.0	-	0.0	-	0.0	-	-	-	-	-	-	-
113.0	55.0	-	0.0	-	0.0	-	-	-	-	-	-	-
113.0	60.0	-	0.0	-	0.0	-	-	-	-	-	-	-
113.0	65.0	-	0.0	-	0.0	-	-	-	-	-	-	-
113.0	70.0	-	0.0	-	0.0	-	-	-	-	-	-	-
113.0	80.0	-	0.0	-	0.0	-	-	-	-	-	-	-
113.0	85.0	-	-	-	3.6	-	-	-	-	-	-	-
113.0	90.0	-	0.0	-	0.0	-	-	-	-	-	-	-
117.0	30.0	-	0.0	-	3.1	0.0	-	-	-	-	-	-
117.0	40.0	-	0.0	-	0.0	-	-	-	-	-	-	-
117.0	70.0	-	0.0	-	0.0	-	-	-	-	-	-	-
117.0	45.0	-	0.0	-	0.0	-	-	-	-	-	-	-
117.0	50.0	-	0.0	-	0.0	-	-	-	-	-	-	-
117.0	55.0	-	0.0	-	0.0	-	-	-	-	-	-	-
117.0	60.0	-	0.0	-	0.0	-	-	-	-	-	-	-
120.0	45.0	-	0.0	-	0.0	-	-	-	-	-	-	-
120.0	50.0	-	0.0	-	0.0	-	-	-	-	-	-	-
120.0	55.0	-	0.0	-	0.0	-	-	-	-	-	-	-
120.0	60.0	-	0.0	-	0.0	-	-	-	-	-	-	-
120.0	70.0	-	0.0	-	0.0	-	-	-	-	-	-	-
120.0	75.0	-	0.0	-	0.0	-	-	-	-	-	-	-
120.0	80.0	-	0.0	-	0.0	-	-	-	-	-	-	-
123.0	42.0	-	-	-	-	-	-	-	-	0.0	3.2	0.0

TABLE 4. (cont.)

*Bathyergus wesethi* (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
123.0	45.0	-	0.0	-	0.0	-	-	3.3	-	0.0	-	-
123.0	50.0	-	2.9	-	0.0	-	-	0.0	-	0.0	-	-
123.0	55.0	-	0.0	-	0.0	-	-	5.7	-	0.0	-	-
123.0	60.0	-	0.0	-	0.0	-	-	5.5	-	9.3	-	-
123.0	65.0	-	0.0	-	0.0	-	-	5.6	-	-	-	-
123.0	70.0	-	0.0	-	0.0	-	-	5.5	-	-	-	-
123.0	80.0	-	0.0	-	0.0	-	-	16.9	-	-	-	-
127.0	40.0	-	0.0	-	11.9	-	-	0.0	0.0	0.0	-	-
127.0	45.0	-	0.0	-	3.1	-	-	3.2	-	0.0	-	-
127.0	50.0	-	0.0	-	0.0	-	-	2.9	-	0.0	-	-
127.0	60.0	-	2.7	-	0.0	-	-	2.9	-	0.0	-	-
127.0	65.0	-	2.9	-	0.0	-	-	2.9	-	-	-	-
130.0	35.0	-	3.3	-	3.1	-	-	0.0	0.0	0.0	-	-
130.0	40.0	-	0.0	-	6.3	-	-	0.0	0.0	0.0	-	-
130.0	45.0	-	0.0	-	3.0	-	-	0.0	0.0	0.0	-	-
130.0	50.0	-	0.0	-	0.0	-	-	0.0	0.0	0.0	-	-
130.0	55.0	-	0.0	-	0.0	-	-	13.7	0.0	0.0	-	-
130.0	60.0	-	0.0	-	0.0	-	-	67.4	0.0	0.0	-	-
133.0	50.0	-	0.0	-	3.0	-	-	0.0	-	0.0	-	-
137.0	60.0	-	3.1	-	-	-	-	0.0	-	-	-	-

*Leuroglossus stibbius*

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
60.0	60.0	13.3	-	-	-	-	-	0.0	-	-	-	-
63.0	55.0	76.1	-	-	-	-	-	0.0	-	-	-	-
63.0	60.0	32.6	-	-	-	-	-	2.8	-	-	-	-
63.0	65.0	-	-	-	-	-	-	-	-	-	-	-
63.0	70.0	-	-	-	-	-	-	-	-	-	-	-
63.0	80.0	-	-	-	-	-	-	-	-	-	-	-
67.0	50.0	0.0	-	-	-	-	-	-	-	-	-	-
67.0	55.0	6.6	-	-	-	-	-	-	-	-	-	-
67.0	60.0	60.2	-	-	-	-	-	-	-	-	-	-
67.0	65.0	-	-	-	-	-	-	-	-	-	-	-
67.0	70.0	-	-	-	-	-	-	-	-	-	-	-
67.0	80.0	-	-	-	-	-	-	-	-	-	-	-
70.0	50.0	0.0	-	-	-	-	-	-	-	-	-	-
70.0	51.0	70.6	-	-	-	-	-	-	-	-	-	-
70.0	52.0	132.3	-	-	-	-	-	-	-	-	-	-
70.0	53.0	73.8	-	-	-	-	-	-	-	-	-	-
70.0	55.0	60.9	-	-	-	-	-	-	-	-	-	-
70.0	60.0	0.0	-	-	-	-	-	-	-	-	-	-
70.0	65.0	-	-	-	-	-	-	-	-	-	-	-
70.0	70.0	8.1	-	-	-	-	-	-	-	-	-	-
70.0	80.0	8.7	-	-	-	-	-	-	-	-	-	-
70.0	80.0	2.7	-	-	-	-	-	-	-	-	-	-

TABLE 4. (cont.)

*Leuroglossus stilius* (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
73.0	51.0	9.3	-	-	5.6	-	-	3.2	-	-	-	-
73.0	53.0	384.9	-	-	6.0	-	-	5.6	-	-	-	-
73.0	60.0	5.6	-	-	14.1	-	-	2.8	-	-	-	-
73.0	80.0	-	-	-	7.8	-	-	-	-	-	-	-
77.0	47.6	2.8	-	-	0.0	-	-	0.0	-	-	-	-
77.0	51.0	133.6	-	-	0.0	-	-	8.6	-	-	-	-
77.0	55.0	28.8	-	-	7.2	-	-	3.1	-	-	-	-
77.0	57.0	98.7	-	-	2.5	-	-	2.8	-	-	-	-
77.0	60.0	-	-	-	15.7	-	-	-	-	-	-	-
77.0	65.0	-	-	-	71.4	-	-	-	-	-	-	-
77.0	70.0	-	-	-	2.3	-	-	-	-	-	-	-
77.0	80.0	-	-	-	5.7	-	-	-	-	-	-	-
80.0	50.2	2.5	-	-	0.8	-	-	0.0	-	-	-	-
80.0	51.0	322.3	-	-	0.0	-	-	0.0	-	-	-	-
80.0	52.0	177.7	-	-	3.3	-	-	2.2	-	-	-	-
80.0	55.0	267.5	-	-	35.2	-	-	0.0	-	-	-	-
80.0	60.0	43.7	-	-	74.6	-	-	2.8	-	-	-	-
80.0	65.0	0.0	-	-	40.4	-	-	3.0	-	-	-	-
80.0	70.0	0.0	-	-	85.0	-	-	0.0	-	-	-	-
80.0	80.0	0.0	-	-	29.0	-	-	0.0	-	-	-	-
82.0	47.0	368.9	-	-	24.8	-	-	0.0	-	-	-	-
82.0	49.0	74.0	-	-	33.6	-	-	22.1	-	-	-	-
83.0	51.0	111.8	-	-	2.6	-	-	3.2	-	-	-	-
83.0	55.0	176.9	-	-	0.0	-	-	6.6	-	-	-	-
83.0	60.0	179.4	-	-	12.0	-	-	51.8	-	-	-	-
83.0	65.0	232.5	-	-	12.9	-	-	13.2	-	-	-	-
83.0	70.0	16.0	-	-	3.2	-	-	27.0	-	-	-	-
83.0	80.0	7.1	-	-	23.0	-	-	8.1	-	-	-	-
83.0	90.0	0.0	-	-	-	-	-	3.2	-	-	-	-
87.0	32.5	15.8	-	-	0.0	-	-	0.0	-	-	-	-
87.0	33.0	19.9	-	-	0.0	-	-	0.0	-	-	-	-
87.0	34.0	285.7	-	-	0.0	-	-	2.6	-	-	-	-
87.0	35.0	1129.0	-	-	20.5	-	-	9.2	-	-	-	-
87.0	40.0	892.0	-	-	91.1	-	-	6.6	-	-	-	-
87.0	45.0	541.6	-	-	17.1	-	-	0.0	-	-	-	-
87.0	50.0	98.5	-	-	9.2	-	-	13.6	-	-	-	-
87.0	55.0	759.9	-	-	5.6	-	-	42.5	-	-	-	-
87.0	60.0	0.0	-	-	23.9	-	-	6.5	-	-	-	-
87.0	65.0	0.0	-	-	2.7	-	-	0.0	-	-	-	-
87.0	80.0	3.0	-	-	6.0	-	-	0.0	-	-	-	-
87.0	90.0	3.0	-	-	2.8	-	-	0.0	-	-	-	-
90.0	27.5	-	-	-	3.3	-	-	0.0	-	-	-	-
90.0	28.0	440.8	-	-	12.2	-	-	0.0	-	-	-	-
90.0	32.0	348.3	-	-	24.4	-	-	0.0	-	-	-	-
90.0	37.0	624.9	-	-	64.9	-	-	0.0	-	-	-	-
90.0	45.0	-	-	-	332.7	-	-	0.0	-	-	-	-

TABLE 4. (cont.)

*Leuroglossus stilibius* (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
90.0	53.0	232.6	-	-	39.8	-	-	0.0	-	0.0	-	-
90.0	60.0	6.0	-	-	0.0	-	-	0.0	-	0.0	-	-
90.0	65.0	0.0	-	-	5.9	-	-	0.0	-	0.0	-	-
90.0	80.0	0.0	-	-	3.2	-	-	0.0	-	0.0	-	-
90.0	90.0	0.0	-	-	3.2	-	-	0.0	-	0.0	-	-
90.0	100.0	0.0	-	-	6.2	-	-	0.0	-	0.0	-	-
93.0	27.0	0.0	-	-	11.9	-	-	11.5	-	0.0	-	-
93.0	28.0	52.5	-	-	23.2	-	-	19.3	-	0.0	-	-
93.0	30.0	653.6	-	-	37.3	-	-	9.2	-	0.0	-	-
93.0	35.0	432.0	-	-	11.6	-	-	6.1	-	0.0	-	-
93.0	40.0	57.2	-	-	52.6	-	-	6.3	-	0.0	-	-
93.0	45.0	483.4	-	-	22.3	-	-	3.3	-	0.0	-	-
93.0	50.0	186.6	-	-	17.3	-	-	6.6	-	0.0	-	-
93.0	55.0	128.9	-	-	70.7	-	-	3.3	-	0.0	-	-
93.0	60.0	3.0	-	-	47.3	-	-	6.7	-	0.0	-	-
93.0	65.0	3.2	-	-	113.2	-	-	0.0	-	0.0	-	-
93.0	70.0	0.0	-	-	63.3	-	-	13.2	-	0.0	-	-
93.0	80.0	0.0	-	-	9.5	-	-	3.2	-	0.0	-	-
96.8	28.7	31.9	-	-	0.0	-	-	0.0	-	0.0	-	-
97.0	29.0	48.6	-	-	0.0	-	-	0.0	-	0.0	-	-
97.0	30.0	118.8	-	-	0.0	-	-	20.2	-	0.0	-	-
97.0	32.0	259.0	-	-	22.1	-	-	3.3	-	0.0	-	-
97.0	35.0	86.0	-	-	9.6	-	-	29.4	-	0.0	-	-
97.0	40.0	167.0	-	-	10.4	-	-	9.5	-	0.0	-	-
97.0	45.0	212.8	-	-	41.9	-	-	0.0	-	0.0	-	-
97.0	50.0	182.7	-	-	-	-	-	0.0	-	0.0	-	-
97.0	55.0	39.0	-	-	-	-	-	0.0	-	0.0	-	-
97.0	60.0	14.8	-	-	-	-	-	0.0	-	0.0	-	-
97.0	65.0	95.2	-	-	79.3	-	-	3.0	-	0.0	-	-
97.0	70.0	41.3	-	-	-	-	-	0.0	-	0.0	-	-
97.0	75.0	-	-	-	-	-	-	-	-	0.0	-	-
97.0	85.0	-	-	-	-	-	-	-	-	0.0	-	-
100.0	29.0	56.3	-	-	8.7	-	-	2.5	-	0.0	-	-
100.0	30.0	30.9	-	-	32.1	-	-	0.0	-	0.0	-	-
100.0	35.0	11.7	-	-	9.5	-	-	0.0	-	0.0	-	-
100.0	40.0	5.9	-	-	6.4	-	-	0.0	-	0.0	-	-
100.0	45.0	49.8	-	-	20.4	-	-	0.0	-	0.0	-	-
100.0	50.0	97.7	-	-	60.2	-	-	0.0	-	0.0	-	-
100.0	55.0	36.4	-	-	18.1	-	-	0.0	-	0.0	-	-
100.0	60.0	44.7	-	-	16.8	-	-	0.0	-	0.0	-	-
100.0	65.0	80.5	-	-	35.1	-	-	0.0	-	0.0	-	-
100.0	70.0	123.2	-	-	3.3	-	-	0.0	-	0.0	-	-
100.0	75.0	-	-	-	10.1	-	-	0.0	-	0.0	-	-
100.0	80.0	5.8	-	-	-	-	-	0.0	-	0.0	-	-
103.0	35.0	41.2	-	-	-	-	-	0.0	-	0.0	-	-
103.0	40.0	20.4	-	-	-	-	-	0.0	-	0.0	-	-
		84.0	-	-	-	-	-	6.7	-	0.0	-	-
		85.7	-	-	-	-	-	-	-	0.0	-	-

TABLE 4. (cont.)

*Leuroglossus stibius* (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
103.0	45.0	5.9	-	-	9.1	-	3.4	-	0.0	-	-	-
103.0	50.0	0.0	-	-	13.2	-	0.0	-	0.0	-	-	-
103.0	55.0	51.7	-	-	0.0	-	0.0	-	0.0	-	-	-
103.0	60.0	12.2	-	-	0.0	-	0.0	-	0.0	-	-	-
103.0	65.0	2.8	-	-	9.5	-	0.0	-	0.0	-	-	-
103.0	70.0	12.0	-	-	24.9	-	0.0	-	0.0	-	-	-
103.0	75.0	-	-	-	3.2	-	-	-	-	-	-	-
103.0	80.0	-	-	-	3.0	-	0.0	-	0.0	-	-	-
103.0	85.0	-	-	-	11.6	-	-	-	0.0	-	-	-
103.0	90.0	-	-	-	3.2	-	-	-	0.0	-	-	-
107.0	30.0	-	-	-	0.0	-	0.0	-	0.0	-	-	-
107.0	31.0	-	-	-	33.0	-	0.0	-	0.0	-	-	-
107.0	32.0	-	-	-	167.7	-	0.0	-	0.0	-	-	-
107.0	35.0	-	-	-	750.9	-	17.0	-	0.0	-	-	-
107.0	40.0	-	-	-	131.2	-	122.4	-	0.0	-	-	-
107.0	50.0	-	-	-	161.7	-	94.5	-	0.0	-	-	-
107.0	55.0	-	-	-	5.9	-	15.0	-	0.0	-	-	-
107.0	60.0	-	-	-	3.0	-	0.0	-	0.0	-	-	-
110.0	32.0	-	-	-	12.1	-	0.0	-	0.0	-	-	-
110.0	35.0	-	-	-	0.7	-	0.0	-	0.0	-	-	-
110.0	40.0	-	-	-	8.5	-	3.1	-	0.0	-	-	-
110.0	45.0	-	-	-	119.1	-	8.6	-	0.0	-	-	-
110.0	50.0	-	-	-	0.0	-	3.2	-	0.0	-	-	-
110.0	55.0	-	-	-	2.8	-	0.0	-	0.0	-	-	-
110.0	65.0	-	-	-	0.0	-	3.0	-	0.0	-	-	-
113.0	35.0	-	-	-	2.7	-	0.0	-	0.0	-	-	-
113.0	40.0	-	-	-	0.0	-	0.0	-	0.0	-	-	-
113.0	45.0	-	-	-	26.4	-	26.4	-	0.0	-	-	-
113.0	50.0	-	-	-	40.2	-	0.0	-	0.0	-	-	-
113.0	55.0	-	-	-	9.1	-	0.0	-	0.0	-	-	-
113.0	65.0	-	-	-	3.0	-	0.0	-	0.0	-	-	-
115.0	35.0	-	-	-	3.1	-	0.0	-	0.0	-	-	-
115.0	45.0	-	-	-	11.4	-	14.6	-	0.0	-	-	-
117.0	25.0	-	-	-	2.6	-	0.0	-	0.0	-	-	-
117.0	30.0	-	-	-	5.8	-	0.0	-	0.0	-	-	-
117.0	35.0	-	-	-	22.0	-	6.5	-	0.0	-	-	-
117.0	45.0	-	-	-	91.8	-	0.0	-	0.0	-	-	-
117.0	50.0	-	-	-	41.6	-	14.6	-	0.0	-	-	-
118.0	39.0	-	-	-	6.2	-	9.3	-	0.0	-	-	-
120.0	22.5	-	-	-	1.6	-	0.0	-	0.0	-	-	-
120.0	45.0	-	-	-	8.6	-	0.0	-	0.0	-	-	-
120.0	50.0	-	-	-	0.0	-	5.9	-	0.0	-	-	-
123.0	37.0	-	-	-	0.0	-	0.0	-	0.0	-	-	-
123.0	42.0	-	-	-	4.4	-	0.0	-	0.0	-	-	-
123.0	45.0	-	-	-	0.0	-	5.8	-	0.0	-	-	-
127.0	32.6	-	-	-	1.5	-	0.0	-	0.0	-	-	-
127.0	40.0	-	-	-	2.9	-	3.0	-	0.0	-	-	-

TABLE 4. (cont.)

*Leuroglossus stilbius* (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
127.0	45.0	-	0.0	-	0.0	-	-	-	3.2	-	0.0	-
130.0	25.5	-	0.6	-	0.0	-	-	-	0.0	0.0	0.0	-
130.0	35.0	-	6.7	-	0.0	-	-	-	0.0	0.0	0.0	-
130.0	50.0	-	20.2	-	0.0	-	-	-	0.0	0.0	0.0	-
133.0	30.0	-	0.0	-	15.5	-	-	-	5.9	-	0.0	-
133.0	35.0	-	5.8	-	3.1	-	-	-	0.0	-	0.0	-
133.0	45.0	-	82.0	-	0.0	-	-	-	0.0	-	0.0	-
137.0	30.0	-	3.1	-	6.2	-	-	-	2.8	0.0	0.0	-
137.0	35.0	-	174.2	-	15.1	-	-	-	0.0	0.0	0.0	-
137.0	40.0	-	0.0	-	3.0	-	-	-	0.0	0.0	0.0	-
137.0	45.0	-	0.0	-	3.0	-	-	-	0.0	0.0	0.0	-
137.0	50.0	-	0.0	-	2.7	-	-	-	0.0	0.0	0.0	-
140.0	30.0	-	2.7	-	-	-	-	-	0.0	-	-	-
140.0	40.0	-	8.9	-	-	-	-	-	0.0	-	-	-

## Stomiiformes

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
70.0	100.0	0.0	-	-	-	-	-	-	5.6	-	-	-

## Gonostomatidae

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
60.0	200.0	3.1	-	-	-	-	-	-	0.0	-	-	-
80.0	200.0	0.0	-	-	-	-	-	-	8.4	-	-	-
83.0	90.0	0.0	-	-	-	-	-	-	3.2	-	-	-
93.0	85.0	-	-	-	-	-	-	-	0.0	-	-	-
93.0	90.0	0.0	-	-	-	-	-	-	0.0	-	-	-
100.0	50.0	0.0	-	-	-	-	-	-	0.0	-	-	-
105.0	140.0	-	0.0	-	-	-	-	-	2.8	-	-	-
113.0	65.0	-	0.0	-	-	-	-	-	3.0	-	-	-
117.0	90.0	-	0.0	-	-	-	-	-	9.8	-	-	-
120.0	45.0	-	0.0	-	-	-	-	-	0.0	-	-	-
120.0	50.0	-	0.0	-	-	-	-	-	0.0	-	-	-
120.0	80.0	-	0.0	-	-	-	-	-	0.0	-	-	-
120.0	120.0	-	0.0	-	-	-	-	-	0.0	-	-	-
123.0	55.0	-	0.0	-	-	-	-	-	0.0	-	-	-
127.0	70.0	-	0.0	-	-	-	-	-	2.8	-	-	-
130.0	45.0	-	0.0	-	-	-	-	-	42.6	-	-	-
137.0	30.0	-	0.0	-	-	-	-	-	0.0	-	-	-
137.0	80.0	-	0.0	-	-	-	-	-	5.5	-	-	-
										17.4	-	-

TABLE 4. (cont.)

*Cyclothone spp.*

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
60.0	100.0	6.2	-	-	-	-	-	0.0	-	-	-	-
60.0	140.0	0.0	-	-	-	-	-	31.3	-	-	-	-
60.0	160.0	12.9	-	-	-	-	-	2.9	-	-	-	-
60.0	180.0	3.0	-	-	-	-	-	5.4	-	-	-	-
60.0	200.0	12.4	-	-	-	-	-	71.2	-	-	-	-
70.0	80.0	15.6	-	0.0	-	-	-	0.0	-	-	-	-
70.0	90.0	2.9	-	-	-	-	-	8.3	-	-	-	-
70.0	100.0	0.0	-	-	-	-	-	2.8	-	-	-	-
70.0	120.0	29.8	-	-	-	-	-	2.7	-	-	-	-
70.0	160.0	23.8	-	-	-	-	-	-	-	-	-	-
70.0	200.0	6.1	-	0.0	-	-	-	26.5	-	-	-	-
80.0	60.0	0.0	-	0.0	-	-	-	2.8	0.0	-	-	-
80.0	80.0	3.0	-	0.0	-	-	-	0.0	-	-	-	-
80.0	90.0	2.9	-	-	-	-	-	0.0	-	-	-	-
80.0	100.0	3.0	-	-	-	-	-	5.9	-	-	-	-
80.0	120.0	0.0	-	-	-	-	-	-	-	-	-	-
80.0	160.0	17.5	-	-	-	-	-	3.2	-	-	-	-
80.0	180.0	8.2	-	-	-	-	-	3.3	-	-	-	-
80.0	200.0	5.7	-	-	-	-	-	0.0	-	-	-	-
83.0	90.0	0.0	-	-	0.0	-	-	9.6	-	-	-	-
87.0	60.0	3.0	-	-	-	-	-	9.0	-	-	-	-
87.0	70.0	0.0	-	-	-	-	-	9.0	-	-	-	-
87.0	80.0	0.0	-	-	-	-	-	0.0	-	-	-	-
87.0	90.0	0.0	-	-	-	-	-	21.8	-	-	-	-
90.0	70.0	0.0	-	-	-	-	-	2.8	-	-	-	-
90.0	80.0	0.0	-	-	-	-	-	28.8	-	-	-	-
90.0	85.0	-	-	-	-	-	-	-	-	-	-	-
90.0	90.0	0.0	-	-	-	-	-	3.2	-	-	-	-
90.0	100.0	8.3	-	-	-	-	-	0.0	-	-	-	-
90.0	120.0	5.4	-	-	-	-	-	34.8	-	-	-	-
90.0	140.0	2.6	-	-	-	-	-	10.7	-	-	-	-
90.0	160.0	15.6	-	-	-	-	-	59.6	-	-	-	-
90.0	180.0	0.0	-	-	-	-	-	72.5	-	-	-	-
90.0	200.0	2.8	-	-	-	-	-	11.0	-	-	-	-
93.0	30.0	0.0	-	-	-	-	-	0.0	-	-	-	-
93.0	65.0	3.2	-	-	-	-	-	-	-	-	-	-
93.0	70.0	29.4	-	-	-	-	-	-	-	-	-	-
93.0	80.0	6.0	-	-	-	-	-	-	-	-	-	-
93.0	85.0	-	-	-	-	-	-	-	-	-	-	-
93.0	90.0	0.0	-	-	-	-	-	3.4	-	-	-	-
93.0	100.0	-	-	-	-	-	-	8.5	-	-	-	-
97.0	40.0	0.0	-	-	-	-	-	3.2	-	-	-	-
97.0	45.0	0.0	-	-	-	-	-	3.5	-	-	-	-
97.0	65.0	0.0	-	-	-	-	-	0.0	-	-	-	-
97.0	80.0	0.0	-	-	-	-	-	-	-	-	-	-
97.0	85.0	-	-	-	-	-	-	-	-	-	-	-
								14.3	-	-	-	-

TABLE 4. (cont.)

*Cyclothona* spp. (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
97.0	90.0	3.0	-	-	-	0.0	-	-	-	-	-	-
100.0	40.0	0.0	-	-	0.0	-	-	8.3	-	0.0	-	-
100.0	45.0	0.0	-	-	0.0	-	-	0.0	-	3.1	-	-
100.0	55.0	0.0	-	-	3.0	-	-	0.0	-	0.0	-	-
100.0	60.0	0.0	-	-	0.0	-	-	0.0	-	2.9	-	-
100.0	65.0	0.0	-	-	0.0	-	-	0.0	-	9.1	-	-
100.0	70.0	0.0	-	-	0.0	-	-	0.0	-	15.4	-	-
100.0	80.0	5.8	-	-	3.3	-	-	2.6	-	-	-	-
100.0	85.0	-	-	-	17.3	-	-	-	-	-	-	-
100.0	90.0	15.1	-	-	43.0	-	-	0.0	-	-	-	-
100.0	100.0	-	53.1	-	-	-	-	0.0	-	-	-	-
100.0	120.0	-	5.9	-	-	-	-	0.0	-	-	-	-
103.0	28.5	0.0	-	-	0.0	-	-	0.0	-	1.0	-	-
103.0	35.0	0.0	-	-	0.0	-	-	3.2	-	0.0	-	-
103.0	40.0	0.0	-	-	0.0	-	-	0.0	-	15.7	-	-
103.0	45.0	0.0	-	-	3.0	-	-	0.0	-	0.0	-	-
103.0	50.0	0.0	-	-	3.3	-	-	0.0	-	0.0	-	-
103.0	55.0	0.0	-	-	6.7	-	-	0.0	-	3.1	-	-
103.0	60.0	3.0	-	-	0.0	-	-	6.6	-	27.8	-	-
103.0	65.0	-	-	-	0.0	-	-	3.3	-	9.6	-	-
103.0	70.0	9.0	-	-	6.2	-	-	16.8	-	5.8	-	-
103.0	75.0	-	-	-	16.0	-	-	-	-	-	-	-
103.0	80.0	-	-	-	17.8	-	-	3.0	-	-	-	-
103.0	85.0	-	-	-	15.0	-	-	-	-	-	-	-
103.0	90.0	-	-	-	15.0	-	-	-	-	-	-	-
105.0	140.0	-	-	-	9.1	-	-	0.0	-	5.6	-	-
107.0	32.0	-	-	-	27.8	-	-	-	-	-	-	-
107.0	45.0	-	-	-	0.0	-	-	3.1	-	-	0.0	-
107.0	50.0	-	-	-	0.0	-	-	5.9	-	0.0	-	-
107.0	60.0	-	-	-	2.9	-	-	0.0	-	0.0	-	-
107.0	65.0	-	-	-	9.1	-	-	3.2	-	3.0	-	-
107.0	70.0	-	-	-	2.9	-	-	3.1	-	26.2	-	-
107.0	75.0	-	-	-	3.2	-	-	12.2	-	18.2	-	-
107.0	80.0	-	-	-	1.0	-	-	0.0	-	-	-	-
107.0	90.0	-	-	-	0.0	-	-	3.1	-	-	-	-
110.0	35.0	-	-	-	0.0	-	-	0.0	-	5.7	-	-
110.0	40.0	-	-	-	0.0	-	-	0.0	-	47.9	-	-
110.0	45.0	-	-	-	0.0	-	-	0.0	-	25.7	-	-
110.0	55.0	-	-	-	0.0	-	-	0.0	-	3.1	-	-
110.0	60.0	-	-	-	0.0	-	-	3.0	-	0.0	-	-
110.0	65.0	-	-	-	0.0	-	-	6.1	-	8.9	-	-
110.0	70.0	-	-	-	0.0	-	-	2.9	-	0.0	-	-
110.0	75.0	-	-	-	2.8	-	-	0.0	-	0.0	-	-
110.0	80.0	-	-	-	10.3	-	-	12.8	-	6.1	-	-
110.0	85.0	-	-	-	10.6	-	-	6.1	-	6.1	-	-
110.0	90.0	-	-	-	-	-	-	-	-	0.0	-	2.8

TABLE 4. (cont.)

## Cyclothona spp. (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
110.0	100.0	-	41.6	-	-	-	2.1	-	-	-	-	-
110.0	120.0	-	28.4	-	-	-	16.3	-	-	-	-	-
110.0	140.0	-	21.6	-	-	-	22.0	-	-	-	-	-
110.0	160.0	-	5.6	-	-	-	58.4	-	-	-	-	-
113.0	35.0	-	0.0	3.5	-	-	3.3	-	-	-	-	-
113.0	40.0	-	0.0	0.0	0.0	-	3.2	-	-	-	-	-
113.0	45.0	-	0.0	0.0	6.2	-	3.1	-	-	-	-	-
113.0	50.0	-	0.0	0.0	6.2	0.0	0.0	-	-	-	-	-
113.0	55.0	-	0.0	0.0	0.0	-	5.9	-	-	-	-	-
113.0	65.0	-	0.0	0.0	3.3	0.0	0.0	-	-	-	-	-
113.0	70.0	-	2.7	0.0	0.0	-	0.0	-	-	-	-	-
113.0	85.0	-	-	3.0	9.8	-	-	-	-	-	-	-
113.0	90.0	-	2.8	10.2	-	-	-	-	-	-	-	-
117.0	30.0	-	0.0	0.0	-	-	0.0	-	-	-	-	-
117.0	40.0	-	30.5	3.1	-	-	0.0	-	-	-	-	-
117.0	45.0	-	0.0	2.9	-	-	0.0	-	-	-	-	-
117.0	50.0	-	0.0	2.9	-	-	0.0	-	-	-	-	-
117.0	55.0	-	0.0	0.0	0.0	-	0.0	-	-	-	-	-
117.0	60.0	-	0.0	0.0	2.9	-	0.0	-	-	-	-	-
117.0	65.0	-	0.0	0.0	0.0	-	13.3	-	-	-	-	-
117.0	70.0	-	0.0	0.0	0.0	-	6.2	-	-	-	-	-
117.0	75.0	-	3.0	3.0	-	-	-	-	-	-	-	-
117.0	80.0	-	3.0	15.3	3.3	-	-	-	-	-	-	-
117.0	85.0	-	-	2.9	-	-	-	-	-	-	-	-
117.0	90.0	-	17.6	5.6	9.8	-	2.8	-	-	-	-	-
118.0	39.0	-	0.0	0.0	0.0	-	-	-	-	-	-	-
120.0	45.0	-	0.0	0.0	6.2	-	0.0	-	-	-	-	-
120.0	55.0	-	0.0	0.0	0.0	-	-	-	-	-	-	-
120.0	60.0	-	5.8	0.0	0.0	-	-	-	-	-	-	-
120.0	65.0	-	0.0	0.0	6.1	-	-	-	-	-	-	-
120.0	70.0	-	0.0	0.0	6.6	-	-	-	-	-	-	-
120.0	80.0	-	0.0	0.0	0.0	-	-	-	-	-	-	-
120.0	100.0	-	20.9	-	-	-	-	-	-	-	-	-
120.0	120.0	-	14.5	-	-	-	-	-	-	-	-	-
123.0	42.0	-	1.5	0.0	0.0	-	-	-	-	-	-	-
123.0	45.0	-	2.9	0.0	0.0	-	-	-	-	-	-	-
123.0	50.0	-	0.0	0.0	0.0	-	-	-	-	-	-	-
123.0	60.0	-	0.0	0.0	3.1	-	-	-	-	-	-	-
123.0	65.0	-	0.0	0.0	6.2	0.0	0.0	-	-	-	-	-
123.0	70.0	-	0.0	0.0	3.1	0.0	0.0	-	-	-	-	-
123.0	80.0	-	0.0	0.0	3.1	0.0	0.0	-	-	-	-	-
127.0	40.0	-	1.5	0.0	0.0	-	-	-	-	-	-	-
127.0	45.0	-	0.0	0.0	9.3	-	-	-	-	-	-	-
127.0	50.0	-	0.0	0.0	3.0	0.0	0.0	-	-	-	-	-
127.0	55.0	-	0.0	0.0	5.9	0.0	0.0	-	-	-	-	-
127.0	60.0	-	0.0	0.0	2.7	0.0	0.0	-	-	-	-	-

TABLE 4. (cont.)

## Cyclothonae spp. (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
127.0	65.0	-	0.0	-	2.9	-	-	5.8	-	-	-	-
127.0	70.0	-	0.0	-	0.0	-	-	14.2	-	-	-	-
127.0	80.0	-	8.6	-	-	-	-	16.4	-	-	-	-
130.0	35.0	-	0.0	-	6.2	-	-	0.0	-	-	-	-
130.0	40.0	-	-	-	3.1	-	-	7.9	-	-	-	-
130.0	45.0	-	-	-	0.0	-	-	0.0	-	-	-	-
130.0	50.0	-	-	-	0.0	-	-	19.1	-	-	-	-
130.0	55.0	-	-	-	0.0	-	-	62.8	-	-	-	-
130.0	60.0	-	-	-	0.0	-	-	39.3	-	-	-	-
130.0	70.0	-	-	-	5.7	-	-	2.5	-	-	-	-
130.0	90.0	-	-	-	2.7	-	-	2.7	-	-	-	-
130.0	100.0	-	-	-	16.6	-	-	0.0	-	-	-	-
130.0	120.0	-	-	-	63.5	-	-	5.4	-	-	-	-
133.0	40.0	-	-	-	3.0	-	-	5.8	-	-	-	-
133.0	45.0	-	-	-	5.9	-	-	8.0	-	-	-	-
133.0	50.0	-	-	-	0.0	-	-	0.0	-	-	-	-
133.0	55.0	-	-	-	2.7	-	-	2.7	-	-	-	-
133.0	65.0	-	-	-	0.0	-	-	7.9	-	-	-	-
133.0	70.0	-	-	-	0.0	-	-	18.8	-	-	-	-
137.0	35.0	-	-	-	3.1	-	-	-	-	-	-	-
137.0	45.0	-	-	-	0.0	-	-	0.0	-	-	-	-
137.0	65.0	-	-	-	0.0	-	-	0.0	-	-	-	-
137.0	80.0	-	-	-	5.9	-	-	-	-	-	-	-
140.0	40.0	-	-	-	3.0	-	-	-	-	-	-	-
140.0	50.0	-	-	-	3.0	-	-	2.3	-	-	-	-

## Diplophos taenia

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
110.0	140.0	-	0.0	-	-	-	-	2.8	-	-	-	-
127.0	65.0	-	0.0	-	0.0	-	-	2.9	-	-	-	-
130.0	50.0	-	0.0	-	0.0	-	-	2.7	0.0	-	-	-

## Ichthyococcus spp.

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
110.0	120.0	-	2.6	-	-	-	-	2.7	-	-	-	-
113.0	50.0	-	0.0	-	0.0	-	-	0.0	-	-	-	-
117.0	55.0	-	0.0	-	0.0	-	-	0.0	-	-	-	-
127.0	55.0	-	2.9	-	0.0	-	-	0.0	-	-	-	-
127.0	60.0	-	2.7	-	0.0	-	-	0.0	-	-	-	-
130.0	55.0	-	0.0	-	0.0	-	-	2.7	0.0	-	-	-

TABLE 4. (cont.)

*Vinciguerria lucetia*

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
60.0	160.0	6.5	-	-	-	-	0.0	-	-	-	-	-
60.0	180.0	3.0	-	-	-	-	0.0	-	-	-	-	-
60.0	200.0	21.8	-	-	0.0	-	98.6	-	-	-	-	-
70.0	80.0	8.3	-	-	-	-	0.0	-	-	-	-	-
70.0	120.0	8.9	-	-	-	-	2.7	-	-	-	-	-
70.0	160.0	8.9	-	-	-	-	-	-	-	-	-	-
70.0	180.0	14.9	-	-	-	-	-	-	-	-	-	-
70.0	200.0	9.2	-	-	-	-	31.8	-	-	-	-	-
73.0	60.0	0.0	-	-	2.8	-	0.0	-	-	-	-	-
80.0	90.0	0.0	-	-	-	-	2.8	-	-	-	-	-
80.0	140.0	2.7	-	-	-	-	-	-	-	-	-	-
80.0	160.0	55.5	-	-	-	-	-	-	-	-	-	-
80.0	180.0	21.9	-	-	-	-	-	-	-	-	-	-
80.0	200.0	0.0	-	-	-	-	-	-	-	-	-	-
87.0	65.0	0.0	-	-	0.0	-	0.0	-	-	-	-	-
87.0	80.0	0.0	-	-	-	-	3.0	-	-	-	-	-
87.0	90.0	0.0	-	-	-	-	40.6	-	-	-	-	-
90.0	45.0	0.0	-	-	-	-	0.0	-	-	-	-	-
90.0	53.0	0.0	-	-	-	-	3.3	-	-	-	-	-
90.0	70.0	0.0	-	-	-	-	40.6	-	-	-	-	-
90.0	75.0	-	-	-	-	-	0.0	-	-	-	-	-
90.0	80.0	0.0	-	-	-	-	0.0	-	-	-	-	-
90.0	85.0	-	-	-	-	-	-	-	-	-	-	-
90.0	90.0	0.0	-	-	-	-	-	-	-	-	-	-
90.0	100.0	0.0	-	-	-	-	-	-	-	-	-	-
90.0	120.0	8.1	-	-	-	-	-	-	-	-	-	-
90.0	140.0	0.0	-	-	-	-	-	-	-	-	-	-
90.0	160.0	7.8	-	-	-	-	-	-	-	-	-	-
90.0	180.0	5.5	-	-	-	-	-	-	-	-	-	-
90.0	200.0	14.1	-	-	-	-	-	-	-	-	-	-
93.0	35.0	0.0	-	-	0.0	-	0.0	-	-	-	-	-
93.0	40.0	0.0	-	-	-	-	-	-	-	-	-	-
93.0	50.0	0.0	-	-	-	-	-	-	-	-	-	-
93.0	65.0	0.0	-	-	-	-	-	-	-	-	-	-
93.0	70.0	5.9	-	-	-	-	-	-	-	-	-	-
93.0	75.0	-	-	-	-	-	-	-	-	-	-	-
93.0	80.0	0.0	-	-	-	-	-	-	-	-	-	-
93.0	85.0	-	-	-	-	-	-	-	-	-	-	-
93.0	90.0	0.0	-	-	-	-	-	-	-	-	-	-
93.0	100.0	-	-	-	-	-	-	-	-	-	-	-
97.0	35.0	0.0	-	-	-	-	-	-	-	-	-	-
97.0	40.0	0.0	-	-	-	-	-	-	-	-	-	-
97.0	45.0	0.0	-	-	-	-	-	-	-	-	-	-
97.0	60.0	0.0	-	-	-	-	-	-	-	-	-	-
97.0	65.0	0.0	-	-	-	-	-	-	-	-	-	-
97.0	70.0	0.0	-	-	-	-	-	-	-	-	-	-

TABLE 4. (cont.)

*Vinciguerria lucetia* (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
97.0	75.0	-	-	-	0.0	-	-	-	-	16.1	-	-
97.0	80.0	0.0	-	-	-	-	3.2	-	-	-	-	-
97.0	85.0	-	-	-	53.7	-	6.2	-	-	-	-	-
97.0	90.0	0.0	-	-	-	-	-	-	-	-	-	-
100.0	30.0	0.0	-	-	0.0	-	-	0.0	-	2.8	-	-
100.0	35.0	0.0	-	-	0.0	-	-	0.0	-	6.1	-	-
100.0	40.0	0.0	-	-	0.0	-	-	30.4	-	3.1	-	-
100.0	45.0	0.0	-	-	0.0	-	-	2.9	-	24.8	-	-
100.0	50.0	0.0	-	-	6.0	-	-	0.0	-	39.3	-	-
100.0	60.0	0.0	-	-	6.7	-	-	0.0	-	40.0	-	-
100.0	65.0	0.0	-	-	0.0	-	-	0.0	-	120.8	-	-
100.0	70.0	3.1	-	-	0.0	-	7.0	-	-	262.6	-	-
100.0	75.0	-	-	-	3.4	-	-	-	-	-	-	-
100.0	80.0	0.0	-	-	0.0	-	-	5.1	-	-	-	-
100.0	85.0	-	-	-	113.8	-	-	-	-	-	-	-
100.0	90.0	6.1	-	-	9.9	-	-	35.2	-	-	-	-
100.0	100.0	-	3.0	-	-	-	-	263.1	-	-	-	-
100.0	120.0	-	8.9	-	-	-	-	45.0	-	-	-	-
103.0	28.5	0.0	-	-	0.0	-	-	-	-	1.0	-	-
103.0	29.0	0.0	-	-	0.0	-	-	-	-	5.1	-	-
103.0	30.0	0.0	-	-	3.1	-	-	-	-	2.6	-	-
103.0	40.0	0.0	-	-	0.0	-	-	-	-	62.6	-	-
103.0	50.0	0.0	-	-	6.6	-	-	33.0	-	18.8	-	-
103.0	55.0	0.0	-	-	3.3	-	-	10.1	-	12.4	-	-
103.0	60.0	0.0	-	-	0.0	-	-	6.6	-	423.3	-	-
103.0	65.0	0.0	-	-	0.0	-	-	29.7	-	261.6	-	-
103.0	70.0	0.0	-	-	0.0	-	57.1	-	-	229.1	-	-
103.0	75.0	-	-	-	22.3	-	-	-	-	-	-	-
103.0	80.0	-	-	-	62.4	-	45.2	-	-	-	-	-
103.0	85.0	-	-	-	15.0	-	-	26.6	-	-	-	-
103.0	90.0	-	-	-	18.2	-	3.2	-	-	94.9	-	-
105.0	140.0	-	-	-	30.6	-	-	-	-	-	-	-
107.0	31.0	0.0	-	-	0.0	-	-	0.0	-	9.5	-	-
107.0	32.0	0.0	-	-	0.0	-	-	0.0	-	9.0	-	-
107.0	40.0	0.0	-	-	3.0	-	-	0.0	-	2.9	-	-
107.0	45.0	0.0	-	-	3.0	-	-	3.5	-	19.7	-	-
107.0	50.0	0.0	-	-	3.0	-	-	30.0	-	3.2	-	-
107.0	55.0	0.0	-	-	0.0	-	-	19.9	-	6.1	-	-
107.0	60.0	3.0	-	-	0.0	-	-	23.6	-	32.5	-	-
107.0	65.0	8.8	-	-	3.1	-	-	6.3	-	370.6	-	-
107.0	70.0	6.1	-	-	6.1	-	-	119.7	-	187.9	-	-
107.0	75.0	-	-	-	31.8	-	-	18.5	-	-	-	-
107.0	80.0	31.9	-	-	3.0	-	-	103.0	-	-	-	-
107.0	85.0	-	-	-	2.9	-	-	-	-	-	-	-
107.0	90.0	53.3	-	-	15.3	-	-	197.1	-	-	-	-
110.0	32.0	0.0	-	-	0.0	-	-	-	-	1.5	0.0	-

TABLE 4. (cont.)

*Vinciguerria lucetia* (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
110.0	35.0	-	0.0	-	0.0	-	-	-	57.0	3.0	-	-
110.0	40.0	-	2.8	-	0.0	-	-	-	552.7	0.0	-	-
110.0	45.0	-	0.0	-	0.0	-	-	-	128.5	10.1	-	-
110.0	50.0	-	2.8	-	0.0	-	-	-	11.4	6.2	-	-
110.0	55.0	-	0.0	-	3.0	-	-	-	0.0	88.7	-	-
110.0	60.0	-	2.7	-	42.7	-	-	-	2.8	187.7	-	-
110.0	65.0	-	2.7	-	2.9	-	-	-	3.0	43.5	-	-
110.0	70.0	-	0.0	-	32.0	-	-	-	3.0	-	-	-
110.0	75.0	-	-	-	-	-	-	-	-	-	-	-
110.0	80.0	-	-	-	-	-	-	-	-	-	-	-
110.0	85.0	-	-	-	-	-	-	-	-	-	-	-
110.0	90.0	-	-	-	-	-	-	-	-	-	-	-
110.0	100.0	-	-	-	-	-	-	-	-	-	-	-
110.0	120.0	-	-	-	-	-	-	-	-	-	-	-
110.0	140.0	-	-	-	-	-	-	-	-	-	-	-
110.0	160.0	-	-	-	-	-	-	-	-	-	-	-
113.0	28.5	-	-	-	-	-	-	-	-	-	-	-
113.0	29.0	-	-	-	-	-	-	-	-	-	-	-
113.0	30.0	-	-	-	-	-	-	-	-	-	-	-
113.0	35.0	-	-	-	-	-	-	-	-	-	-	-
113.0	40.0	-	-	-	-	-	-	-	-	-	-	-
113.0	45.0	-	-	-	-	-	-	-	-	-	-	-
113.0	50.0	-	-	-	-	-	-	-	-	-	-	-
113.0	55.0	-	-	-	-	-	-	-	-	-	-	-
113.0	60.0	-	-	-	-	-	-	-	-	-	-	-
113.0	65.0	-	-	-	-	-	-	-	-	-	-	-
113.0	70.0	-	-	-	-	-	-	-	-	-	-	-
113.0	75.0	-	-	-	-	-	-	-	-	-	-	-
113.0	80.0	-	-	-	-	-	-	-	-	-	-	-
113.0	85.0	-	-	-	-	-	-	-	-	-	-	-
113.0	90.0	-	-	-	-	-	-	-	-	-	-	-
115.0	35.0	-	0.0	-	-	-	-	-	0.0	6.3	-	-
117.0	35.0	-	0.0	-	-	-	-	-	17.2	2.9	-	-
117.0	40.0	-	0.0	-	-	-	-	-	18.1	9.3	-	-
117.0	45.0	-	3.0	-	-	-	-	-	0.0	43.4	-	-
117.0	50.0	-	0.0	-	-	-	-	-	29.9	106.4	-	-
117.0	55.0	-	0.0	-	-	-	-	-	13.2	357.9	-	-
117.0	60.0	-	0.0	-	-	-	-	-	-	15.4	-	-
117.0	65.0	-	0.0	-	-	-	-	-	-	43.3	-	-
117.0	70.0	-	70.8	-	10.2	-	-	-	17.0	83.7	-	-
117.0	75.0	-	0.0	-	-	-	-	-	-	-	-	-
117.0	80.0	-	-	-	-	-	-	-	-	-	-	-
117.0	85.0	-	-	-	-	-	-	-	-	-	-	-
117.0	90.0	-	-	-	-	-	-	-	-	-	-	-
118.0	39.0	-	-	-	-	-	-	-	-	-	-	-
120.0	25.0	-	-	-	-	-	-	-	-	-	-	-

TABLE 4. (cont.)

*Vinciguerria lucetia* (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
120.0	40.0	-	0.0	-	0.0	-	-	0.0	-	2.7	-	-
120.0	45.0	-	0.0	-	0.0	-	-	-	5.7	3.0	-	-
120.0	50.0	-	0.0	-	0.0	-	-	-	12.0	0.0	-	-
120.0	55.0	-	0.0	-	0.0	-	-	-	250.9	154.4	-	-
120.0	60.0	-	2.9	-	5.8	-	-	-	169.2	533.8	-	-
120.0	65.0	-	2.8	-	12.2	-	-	-	98.1	-	-	-
120.0	70.0	-	8.8	-	19.8	-	-	-	169.8	-	-	-
120.0	75.0	-	-	-	3.1	-	-	-	-	-	-	-
120.0	80.0	-	5.6	-	17.1	-	-	-	1205.1	-	-	-
120.0	90.0	-	2.9	-	-	-	-	-	210.6	-	-	-
120.0	100.0	-	107.6	-	-	-	-	-	176.0	-	-	-
120.0	120.0	-	203.0	-	-	-	-	-	-	-	-	-
123.0	42.0	-	4.4	-	0.0	-	-	-	28.4	9.5	6.0	-
123.0	45.0	-	5.8	-	2.9	-	-	-	219.1	3.0	-	-
123.0	50.0	-	17.6	-	37.1	-	-	-	221.5	-	67.8	-
123.0	55.0	-	6.0	-	40.0	-	-	-	11.4	9.3	-	-
123.0	60.0	-	2.9	-	-	-	-	-	8.2	306.9	-	-
123.0	65.0	-	2.9	-	43.7	-	-	-	168.6	-	-	-
123.0	70.0	-	5.6	-	70.8	-	-	-	133.0	-	-	-
123.0	75.0	-	-	-	40.6	-	-	-	-	-	-	-
123.0	80.0	-	0.0	-	92.4	-	-	-	341.2	-	-	-
127.0	34.0	-	0.0	-	0.0	-	-	-	0.0	9.0	-	-
127.0	40.0	-	18.5	-	23.8	-	-	-	0.0	2.7	40.8	-
127.0	45.0	-	18.0	-	230.1	-	-	-	29.1	-	136.4	-
127.0	50.0	-	2.9	-	127.3	-	-	-	17.5	-	33.2	-
127.0	55.0	-	2.9	-	273.2	-	-	-	38.4	-	12.7	-
127.0	60.0	-	0.0	-	86.2	-	-	-	8.8	-	57.3	-
127.0	65.0	-	17.3	-	88.2	-	-	-	60.5	-	-	-
127.0	70.0	-	55.9	-	89.3	-	-	-	514.0	-	-	-
127.0	80.0	-	83.5	-	-	-	-	-	1367.7	-	-	-
130.0	25.5	-	0.0	-	1.8	-	-	-	0.0	0.0	-	-
130.0	35.0	-	0.0	-	49.9	-	-	-	5.1	0.0	-	-
130.0	40.0	-	18.4	-	72.5	-	-	-	170.3	438.0	-	-
130.0	45.0	-	2.8	-	39.5	-	-	-	150.0	284.8	-	-
130.0	50.0	-	69.1	-	121.0	-	-	-	368.5	268.9	-	-
130.0	55.0	-	48.8	-	96.7	-	-	-	600.6	1069.5	-	-
130.0	60.0	-	36.3	-	43.3	-	-	-	1337.6	59.5	-	-
130.0	65.0	-	-	-	21.6	-	-	-	-	-	-	-
130.0	70.0	-	43.0	-	98.6	-	-	-	-	-	181.8	-
130.0	80.0	-	83.4	-	-	-	-	-	-	-	120.1	-
130.0	90.0	-	13.4	-	-	-	-	-	-	-	582.7	-
130.0	100.0	-	138.5	-	-	-	-	-	-	-	687.3	-
130.0	120.0	-	419.5	-	-	-	-	-	-	-	21.7	2.5
133.0	23.0	-	-	-	0.0	-	-	-	-	-	0.0	54.3
133.0	30.0	-	0.0	-	3.1	-	-	-	-	-	0.0	-
133.0	35.0	-	43.2	-	-	-	-	-	-	-	0.0	124.3

TABLE 4. (cont.)

*Vinciguerria lucetiae* (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
133.0	40.0	-	29.8	-	203.3	-	-	509.3	-	48.6	-	-
133.0	45.0	-	61.5	-	42.3	-	-	162.3	-	2.8	-	-
133.0	50.0	-	42.5	-	32.6	-	-	60.3	-	0.0	-	-
133.0	55.0	-	11.0	-	59.2	-	-	55.9	-	-	-	-
133.0	60.0	-	8.6	-	33.2	-	-	36.4	-	-	-	-
133.0	65.0	-	0.0	-	219.4	-	-	34.3	-	-	-	-
133.0	70.0	-	5.7	-	166.3	-	-	380.6	-	-	-	-
133.0	80.0	-	46.2	-	-	-	-	160.6	-	-	-	-
137.0	21.0	-	0.0	-	0.0	-	-	0.0	-	0.0	-	-
137.0	22.0	-	1.3	-	0.0	-	-	-	-	10.7	2.2	-
137.0	23.0	-	0.0	-	0.0	-	-	-	-	7.6	0.0	-
137.0	30.0	-	46.2	-	6.2	-	-	-	-	0.0	24.4	-
137.0	35.0	-	62.2	-	0.0	-	-	-	-	2.7	77.3	-
137.0	40.0	-	163.2	-	18.0	-	-	-	-	0.0	125.6	-
137.0	45.0	-	81.2	-	9.1	-	-	-	-	354.3	188.5	-
137.0	50.0	-	26.5	-	10.9	-	-	-	-	304.2	33.0	-
137.0	55.0	-	3.1	-	31.9	-	-	-	-	-	182.2	-
137.0	60.0	-	-	-	52.2	-	-	-	-	-	436.3	-
137.0	65.0	-	-	-	119.5	-	-	-	-	-	-	-
137.0	70.0	-	15.0	-	107.4	-	-	-	-	-	29.3	-
137.0	80.0	-	59.0	-	-	-	-	-	-	-	199.2	-
140.0	28.8	-	0.0	-	-	-	-	-	-	-	0.5	-
140.0	29.0	-	0.0	-	-	-	-	-	-	-	2.0	-
140.0	30.0	-	2.7	-	-	-	-	-	-	-	39.9	-
140.0	35.0	-	9.1	-	-	-	-	-	-	-	34.6	-
140.0	40.0	-	106.2	-	-	-	-	-	-	-	53.4	-
140.0	45.0	-	24.5	-	-	-	-	-	-	-	278.8	-
140.0	50.0	-	26.6	-	-	-	-	-	-	-	593.7	-

*Vinciguerria poweriae*

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
80.0	160.0	8.8	-	-	-	-	-	-	-	-	-	-
80.0	180.0	8.2	-	-	-	-	-	-	-	-	-	-
90.0	200.0	8.4	-	-	-	-	-	-	-	-	-	-

*Sternopychidae*

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
60.0	120.0	2.5	-	-	-	-	-	-	-	-	-	-
60.0	140.0	3.5	-	-	-	-	-	-	-	-	-	-
60.0	180.0	8.9	-	-	-	-	-	-	-	-	-	-
60.0	200.0	6.2	-	-	-	-	-	-	-	-	-	-
70.0	120.0	3.0	-	-	-	-	-	-	-	-	-	-

TABLE 4. (cont.)

## Sternopychidae (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
70.0	180.0	3.0	-	-	3.0	-	-	-	-	-	-	-
73.0	53.0	0.0	-	-	-	-	-	0.0	-	-	-	-
80.0	160.0	2.9	-	-	-	-	-	-	-	-	-	-
80.0	180.0	2.7	-	-	-	-	-	-	-	-	-	-
80.0	200.0	2.8	-	-	-	-	-	0.0	-	-	-	-
87.0	65.0	0.0	-	0.0	-	-	3.2	-	-	-	0.0	-
87.0	90.0	3.0	-	0.0	-	0.0	-	0.0	-	-	-	-
87.0	90.0	3.0	-	0.0	-	-	-	-	-	-	-	-
90.0	80.0	0.0	-	-	0.0	-	-	-	-	-	-	-
90.0	85.0	-	-	-	-	-	-	-	-	-	-	-
90.0	90.0	0.0	-	-	0.0	-	-	-	-	-	-	-
90.0	120.0	0.0	-	-	-	-	-	-	-	-	-	-
93.0	30.0	0.0	-	0.0	-	-	0.0	-	-	-	-	-
93.0	45.0	3.0	-	-	0.0	-	0.0	-	-	-	-	-
93.0	60.0	0.0	-	-	0.0	-	3.3	-	-	-	0.0	-
93.0	90.0	0.0	-	-	6.7	0.0	-	-	-	-	0.0	-
93.0	100.0	-	-	-	2.8	-	-	-	-	-	-	-
97.0	65.0	0.0	-	0.0	-	-	3.0	-	-	-	0.0	-
97.0	90.0	0.0	-	-	-	-	3.1	-	-	-	6.1	-
100.0	35.0	0.0	-	-	0.0	-	-	0.0	-	-	-	-
100.0	90.0	0.0	-	-	-	-	3.3	-	-	-	0.0	-
103.0	50.0	0.0	-	-	0.0	-	3.7	-	-	-	0.0	-
103.0	65.0	0.0	-	-	0.0	-	3.3	-	-	-	0.0	-
103.0	75.0	-	-	-	-	-	3.2	-	-	-	-	-
103.0	80.0	-	-	-	0.0	-	3.0	-	-	-	-	-
103.0	90.0	-	-	-	0.0	-	3.2	-	-	-	-	-
107.0	32.0	-	-	0.0	-	0.0	3.4	-	-	-	0.0	-
107.0	80.0	5.8	-	-	0.0	-	0.0	-	-	-	0.0	-
110.0	45.0	0.0	-	-	-	-	-	-	-	-	-	3.3
110.0	100.0	2.6	-	-	-	-	-	-	-	-	-	-
113.0	80.0	0.0	-	-	0.0	-	3.4	-	-	-	-	-
113.0	90.0	2.8	-	-	0.0	-	0.0	-	-	-	-	-
117.0	50.0	0.0	-	-	2.9	-	0.0	-	-	-	0.0	-
117.0	55.0	0.0	-	-	-	-	6.6	-	-	-	0.0	-
118.0	39.0	0.0	-	-	0.0	-	-	2.8	-	-	0.0	-
120.0	45.0	0.0	-	-	0.0	-	-	-	-	-	6.5	-
120.0	100.0	12.0	-	-	0.0	-	-	-	-	-	0.0	-
123.0	42.0	0.0	-	-	-	-	-	0.0	-	-	2.7	-
123.0	60.0	0.0	-	-	-	-	-	-	-	-	0.0	-
127.0	45.0	0.0	-	-	0.0	-	-	-	-	-	0.0	-
127.0	50.0	8.8	-	-	-	-	-	-	-	-	0.0	-
127.0	60.0	0.0	-	-	0.0	-	-	-	-	-	0.0	-
130.0	40.0	3.1	-	-	-	-	-	-	-	-	3.4	-
130.0	50.0	5.8	-	-	-	-	-	-	-	-	0.0	-
130.0	60.0	0.0	-	-	0.0	-	-	-	-	-	0.0	-
130.0	70.0	2.9	-	-	-	-	-	-	-	-	0.0	-
130.0	100.0	0.0	-	-	-	-	-	-	-	-	0.0	-

TABLE 4. (cont.)

## Sternopychidae (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
133.0	40.0	-	3.0	-	0.0	-	-	0.0	-	6.1	-	-
137.0	40.0	-	3.1	-	3.0	-	-	-	0.0	3.0	-	-
137.0	55.0	-	0.0	-	0.0	-	-	-	5.5	-	-	-

*Chauliodus macouni*

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
60.0	60.0	0.0	-	-	-	-	-	-	2.8	-	-	-
60.0	70.0	3.2	-	-	-	-	-	-	2.8	-	-	-
60.0	100.0	3.1	-	-	-	-	-	-	0.0	-	-	-
60.0	120.0	2.5	-	-	-	-	-	-	0.0	-	-	-
60.0	160.0	3.2	-	-	-	-	-	-	0.0	-	-	-
63.0	55.0	0.0	-	-	0.0	-	-	-	2.8	-	-	-
67.0	60.0	0.0	-	-	0.0	-	-	-	3.0	-	-	-
70.0	53.0	0.0	-	-	2.7	-	-	-	5.8	-	-	-
70.0	60.0	0.0	-	-	0.0	-	-	-	5.3	-	-	-
70.0	65.0	-	-	-	2.7	-	-	-	-	-	-	-
70.0	80.0	0.0	-	-	0.0	-	-	-	3.2	-	-	-
70.0	90.0	0.0	-	-	-	-	-	-	2.8	-	-	-
73.0	53.0	0.0	-	-	0.0	-	-	-	8.3	-	-	-
73.0	60.0	5.6	-	-	0.0	-	-	-	5.6	-	-	-
73.0	80.0	-	-	-	2.6	-	-	-	-	-	-	-
77.0	57.0	0.0	-	-	0.0	-	-	-	2.8	-	-	-
77.0	80.0	-	-	-	2.8	-	-	-	-	-	-	-
80.0	55.0	0.0	-	-	0.0	-	-	-	2.4	0.0	-	-
80.0	60.0	0.0	-	-	0.0	-	-	-	2.8	0.0	-	-
80.0	65.0	0.0	-	-	3.4	-	-	-	0.0	3.0	-	-
80.0	70.0	2.8	-	-	0.0	-	-	-	0.0	0.0	-	-
80.0	80.0	0.0	-	-	0.0	-	-	-	0.0	0.0	-	-
83.0	70.0	6.4	-	-	0.0	-	-	-	-	0.0	-	-
83.0	90.0	6.1	-	-	-	-	-	-	0.0	-	-	-
87.0	35.0	0.0	-	-	3.4	-	-	-	0.0	-	-	-
87.0	55.0	0.0	-	-	0.0	-	-	-	3.3	-	-	-
87.0	60.0	0.0	-	-	0.0	-	-	-	0.0	3.3	-	-
87.0	65.0	3.1	-	-	0.0	-	-	-	0.0	0.0	-	-
87.0	70.0	3.1	-	-	0.0	-	-	-	0.0	3.0	-	-
87.0	80.0	0.0	-	-	3.0	-	-	-	0.0	0.0	-	-
87.0	100.0	-	-	-	9.4	-	-	-	-	0.0	-	-
90.0	45.0	0.0	-	-	3.1	-	-	-	0.0	-	-	-
90.0	60.0	3.0	-	-	0.0	-	-	-	0.0	-	-	-
90.0	65.0	6.5	-	-	0.0	-	-	-	0.0	-	-	-
90.0	100.0	2.8	-	-	3.1	-	-	-	0.0	-	-	-
93.0	60.0	0.0	-	-	3.4	-	-	-	0.0	-	-	-
93.0	90.0	6.4	-	-	0.0	-	-	-	0.0	-	-	-
97.0	35.0	-	-	-	3.2	-	-	-	-	-	-	-

TABLE 4. (cont.)

*Chauliodus macouni* (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
97.0	65.0	0.0	-	0.0	-	-	5.9	-	-	0.0	-	-
97.0	75.0	-	-	-	3.1	-	-	-	-	0.0	-	-
100.0	35.0	2.9	-	0.0	-	-	-	0.0	-	0.0	-	-
100.0	45.0	0.0	-	0.0	-	-	-	0.0	-	3.1	-	-
103.0	35.0	2.9	-	0.0	-	-	0.0	-	-	0.0	-	-
103.0	40.0	2.9	-	0.0	-	-	0.0	-	-	0.0	-	-
103.0	50.0	5.8	-	0.0	-	-	0.0	-	-	0.0	-	-
107.0	35.0	-	3.2	-	3.2	-	-	-	-	0.0	-	-
110.0	70.0	-	2.8	-	0.0	-	-	-	-	0.0	-	-
120.0	60.0	-	0.0	-	-	-	-	-	-	3.1	-	-

*Idiacanthus antrostomus*

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
60.0	140.0	0.0	-	-	-	-	-	10.4	-	-	-	-
60.0	160.0	0.0	-	-	-	-	-	2.9	-	-	-	-
60.0	180.0	5.9	-	-	-	-	-	0.0	-	-	-	-
60.0	200.0	0.0	-	-	-	-	-	2.7	-	-	-	-
70.0	120.0	0.0	-	-	-	-	-	2.7	-	-	-	-
70.0	160.0	3.0	-	-	-	-	-	-	-	-	-	-
80.0	65.0	0.0	-	0.0	-	-	-	3.0	0.0	-	-	-
80.0	160.0	2.9	-	-	-	-	-	-	-	-	-	-
83.0	65.0	0.0	-	0.0	-	-	-	0.0	-	-	-	-
87.0	80.0	0.0	-	0.0	-	-	-	9.8	-	-	-	-
90.0	70.0	0.0	-	0.0	-	-	-	-	-	-	-	-
90.0	80.0	0.0	-	0.0	-	-	-	-	-	-	-	-
90.0	90.0	0.0	-	0.0	-	-	-	-	-	-	-	-
90.0	100.0	0.0	-	0.0	-	-	-	-	-	-	-	-
90.0	120.0	0.0	-	0.0	-	-	-	-	-	-	-	-
90.0	140.0	0.0	-	-	-	-	-	-	-	-	-	-
90.0	160.0	0.0	-	-	-	-	-	-	-	-	-	-
93.0	85.0	-	-	-	-	-	-	-	-	-	-	-
93.0	90.0	0.0	-	0.0	-	-	-	3.1	-	-	-	-
97.0	80.0	0.0	-	0.0	-	-	-	0.0	-	-	-	-
100.0	50.0	0.0	-	0.0	-	-	-	6.6	-	-	-	-
100.0	90.0	0.0	-	0.0	-	-	-	-	-	2.9	-	-
100.0	100.0	-	3.0	-	-	-	-	-	-	0.0	-	-
103.0	65.0	2.8	-	-	-	-	-	-	-	3.3	-	-
107.0	40.0	-	0.0	-	-	-	-	-	-	3.4	-	-
107.0	60.0	-	0.0	-	-	-	-	-	-	2.8	-	-
123.0	50.0	-	0.0	-	-	-	-	-	-	2.7	-	-
127.0	80.0	-	0.0	-	-	-	-	-	-	0.0	-	-
130.0	60.0	-	0.0	-	-	-	-	-	-	0.0	-	-
130.0	120.0	-	0.0	-	-	-	-	-	-	0.0	-	-
133.0	70.0	-	0.0	-	-	-	-	-	-	2.7	-	-

TABLE 4. (cont.)

*Aristostomias scintillans*

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
70.0	200.0	3.1	-	-	-	-	-	0.0	-	-	-	-
80.0	160.0	2.9	-	-	-	-	-	-	-	-	-	-
87.0	90.0	0.0	-	-	0.0	-	6.2	-	-	-	-	-
90.0	120.0	5.4	-	-	-	-	-	2.7	-	-	-	-
93.0	100.0	-	-	-	2.8	-	-	-	-	-	-	-
100.0	90.0	0.0	-	-	3.3	-	-	0.0	-	-	-	-
110.0	100.0	-	2.6	-	-	-	-	0.0	-	-	-	-
110.0	120.0	-	5.2	-	-	-	-	0.0	-	-	-	-

*Bathophilus* spp.

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
90.0	120.0	5.4	-	-	-	-	-	0.0	-	-	-	-
90.0	160.0	2.6	-	-	-	-	-	0.0	-	-	-	-
90.0	180.0	2.7	-	-	-	-	-	0.0	-	-	-	-
100.0	90.0	0.0	-	-	3.3	-	-	0.0	-	-	-	-

*Eustomias* spp.

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
80.0	200.0	0.0	-	-	-	-	-	2.8	-	-	-	-

*Photonectes* spp.

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
60.0	180.0	0.0	-	-	-	-	-	2.7	-	-	-	-
70.0	200.0	0.0	-	-	-	-	-	2.7	-	-	-	-
80.0	160.0	2.9	-	-	-	-	-	-	-	-	-	-
80.0	200.0	0.0	-	-	-	-	-	5.6	-	-	-	-
90.0	120.0	0.0	-	-	-	-	-	10.7	-	-	-	-
90.0	180.0	0.0	-	-	-	-	-	2.8	-	-	-	-

*Tactostoma macropus*

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
60.0	140.0	0.0	-	-	-	-	-	2.6	-	-	-	-
90.0	90.0	0.0	-	-	0.0	-	-	0.0	-	3.1	-	-

TABLE 4. (cont.)

<i>Stomias atriventris</i>												
STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
60.0	100.0	0.0	-	-	-	-	-	2.6	-	-	-	-
80.0	120.0	0.0	-	-	-	-	-	3.0	-	-	-	-
80.0	200.0	2.8	-	-	3.2	-	0.0	0.0	-	-	-	-
87.0	70.0	0.0	-	-	0.0	-	3.1	-	0.0	-	-	-
87.0	90.0	0.0	-	-	9.4	-	-	0.0	-	-	-	-
87.0	100.0	-	-	-	3.1	-	-	0.0	-	-	-	-
90.0	70.0	0.0	-	-	0.0	-	0.0	0.0	-	-	-	-
93.0	70.0	5.9	-	-	6.3	-	0.0	0.0	-	-	-	-
93.0	80.0	0.0	-	-	-	-	-	-	-	-	-	-
93.0	85.0	-	-	-	-	-	-	-	-	-	-	-
93.0	90.0	0.0	-	-	6.7	-	3.1	-	-	-	-	-
93.0	100.0	-	-	-	8.5	-	-	-	-	-	-	-
97.0	25.0	0.0	-	3.2	-	-	0.0	-	-	-	-	-
97.0	50.0	3.2	-	0.2	-	-	0.0	-	-	-	-	-
97.0	60.0	3.0	-	-	-	-	0.0	-	-	-	-	-
97.0	70.0	3.0	-	-	-	-	0.0	-	-	-	-	-
100.0	35.0	0.0	-	-	3.2	-	0.0	-	-	-	-	-
100.0	50.0	0.0	-	-	3.0	-	0.0	-	-	-	-	-
100.0	70.0	3.1	-	-	-	-	0.0	-	-	-	-	-
100.0	80.0	0.0	-	-	-	-	0.0	-	-	-	-	-
100.0	85.0	-	-	-	-	-	0.0	-	-	-	-	-
100.0	100.0	-	-	-	3.0	-	-	-	-	-	-	-
103.0	155.0	2.9	-	-	-	0.0	0.0	-	-	-	-	-
103.0	65.0	0.0	-	-	-	6.3	-	-	-	-	-	-
107.0	32.0	-	-	-	3.1	-	3.1	-	-	-	-	-
107.0	40.0	-	-	-	2.9	-	0.0	-	-	-	-	-
107.0	60.0	-	-	-	0.0	-	3.2	-	-	-	-	-
107.0	70.0	-	-	-	3.2	-	0.0	-	-	-	-	-
107.0	80.0	-	-	-	8.7	-	0.0	-	-	-	-	-
107.0	90.0	-	-	-	14.8	-	3.1	-	-	-	-	-
110.0	40.0	-	-	-	5.5	-	0.0	-	-	-	-	-
110.0	60.0	-	-	-	0.0	-	0.0	-	-	-	-	-
110.0	45.0	-	-	-	25.7	-	0.0	-	-	-	-	-
110.0	80.0	-	-	-	10.6	-	0.0	-	-	-	-	-
110.0	90.0	-	-	-	5.2	-	-	-	-	-	-	-
110.0	100.0	-	-	-	-	-	-	-	-	-	-	-
110.0	120.0	-	-	-	2.6	-	-	-	-	-	-	-
113.0	45.0	-	-	-	0.0	-	6.2	-	-	-	-	-
113.0	50.0	-	-	-	0.0	-	6.2	-	-	-	-	-
113.0	55.0	-	-	-	0.0	-	3.6	-	-	-	-	-
113.0	75.0	-	-	-	-	-	3.3	-	-	-	-	-
113.0	90.0	-	-	-	2.8	-	0.0	-	-	-	-	-
117.0	60.0	-	-	-	-	-	2.9	-	-	-	-	-
117.0	75.0	-	-	-	-	-	3.0	-	-	-	-	-
117.0	80.0	-	-	-	11.9	-	0.0	-	-	-	-	-
117.0	90.0	-	-	-	20.6	-	0.0	-	-	-	-	-
120.0	60.0	-	-	-	8.6	-	0.0	-	-	-	-	-

TABLE 4. (cont.)

*Stomias atriventris* (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
120.0	65.0	-	8.4	-	6.1	-	-	0.0	-	-	-	-
120.0	70.0	-	5.8	-	0.0	-	-	2.8	-	-	-	-
120.0	90.0	-	8.7	-	-	-	-	0.0	0.0	0.0	0.0	-
123.0	42.0	-	1.6	-	0.0	-	-	0.0	-	-	-	-
123.0	55.0	-	3.0	-	0.0	-	-	0.0	-	-	-	-
123.0	60.0	-	0.0	-	0.0	-	-	0.0	-	-	-	-
123.0	70.0	-	5.6	-	0.0	-	-	0.0	-	-	-	-
123.0	75.0	-	-	-	3.1	-	-	-	-	-	-	-
123.0	80.0	-	2.9	-	0.0	-	-	0.0	-	-	-	-
127.0	40.0	-	1.4	-	0.0	-	-	0.0	-	-	-	-
127.0	45.0	-	9.0	-	0.0	-	-	0.0	-	-	-	-
127.0	50.0	-	8.8	-	0.0	-	-	0.0	-	-	-	-
127.0	55.0	-	0.0	-	0.0	-	-	2.7	-	-	-	-
127.0	60.0	-	5.5	-	0.0	-	-	0.0	-	-	-	-
127.0	65.0	-	0.0	-	0.0	-	-	0.0	-	-	-	-
127.0	70.0	-	5.9	-	0.0	-	-	2.9	-	-	-	-
127.0	80.0	-	5.8	-	0.0	-	-	0.0	-	-	-	-
130.0	35.0	-	0.0	-	0.0	-	-	0.0	-	-	-	-
130.0	40.0	-	3.1	-	15.8	-	-	5.2	-	-	-	-
130.0	45.0	-	0.0	-	3.0	-	-	2.8	-	-	-	-
130.0	50.0	-	5.8	-	0.0	-	-	2.7	-	-	-	-
130.0	55.0	-	5.7	-	0.0	-	-	0.0	-	-	-	-
130.0	60.0	-	0.0	-	0.0	-	-	0.0	-	-	-	-
130.0	70.0	-	11.5	-	0.0	-	-	2.5	-	-	-	-
130.0	90.0	-	22.7	-	-	-	-	0.0	-	-	-	-
130.0	100.0	-	22.2	-	-	-	-	0.0	-	-	-	-
133.0	35.0	-	2.9	-	0.0	-	-	0.0	-	-	-	-
133.0	50.0	-	0.0	-	0.0	-	-	2.6	-	-	-	-
133.0	55.0	-	2.7	-	3.0	-	-	0.0	-	-	-	-
133.0	65.0	-	2.9	-	3.1	-	-	0.0	-	-	-	-
137.0	35.0	-	6.2	-	3.0	-	-	0.0	-	-	-	-
137.0	45.0	-	0.0	-	3.0	-	-	0.0	-	-	-	-
137.0	50.0	-	0.0	-	0.0	-	-	0.0	-	-	-	-
137.0	60.0	-	-	-	0.0	-	-	4.3	-	-	-	-
137.0	65.0	-	-	-	3.2	-	-	-	-	-	-	-
137.0	70.0	-	0.0	-	0.0	-	-	2.4	-	-	-	-
137.0	80.0	-	3.0	-	-	-	-	0.0	-	-	-	-
140.0	45.0	-	0.0	-	-	-	-	2.8	-	-	-	-

## Evermannellidae

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
130.0	80.0	-	2.8	-	-	-	-	-	-	-	-	-

TABLE 4. (cont.)

Paralepididae												
STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
60.0	160.0	9.7	-	-	-	-	-	0.0	-	-	-	-
103.0	65.0	0.0	-	-	0.0	-	-	0.0	-	-	-	-
110.0	75.0	-	-	-	3.2	-	-	-	-	-	-	-
<i>Lestidiops ringens</i>												
STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
60.0	200.0	0.0	-	-	-	-	-	-	-	-	-	-
63.0	55.0	0.0	-	-	1.9	-	-	-	-	-	-	-
63.0	60.0	9.8	-	-	0.0	-	-	-	-	-	-	-
67.0	55.0	0.0	-	-	2.7	-	-	-	-	-	-	-
67.0	80.0	-	-	-	5.3	-	-	-	-	-	-	-
70.0	60.0	5.7	-	-	0.0	-	-	-	-	-	-	-
70.0	70.0	5.4	-	-	5.8	-	-	-	-	-	-	-
70.0	90.0	0.0	-	-	-	-	-	-	-	-	-	-
70.0	100.0	0.0	-	-	-	-	-	-	-	-	-	-
73.0	53.0	2.7	-	-	0.0	-	-	-	-	-	-	-
73.0	60.0	2.8	-	-	0.0	-	-	-	-	-	-	-
77.0	55.0	0.0	-	-	3.6	-	-	-	-	-	-	-
80.0	60.0	0.0	-	-	0.0	-	-	-	-	-	-	-
80.0	65.0	0.0	-	-	0.0	-	-	-	-	-	-	-
80.0	80.0	3.0	-	-	0.0	-	-	-	-	-	-	-
80.0	90.0	0.0	-	-	0.0	-	-	-	-	-	-	-
80.0	120.0	5.9	-	-	-	-	-	-	-	-	-	-
83.0	80.0	0.0	-	-	2.9	-	-	-	-	-	-	-
83.0	90.0	3.0	-	-	-	-	-	-	-	-	-	-
87.0	34.0	2.7	-	-	0.0	-	-	-	-	-	-	-
87.0	60.0	0.0	-	-	0.0	-	-	-	-	-	-	-
87.0	65.0	0.0	-	-	0.0	-	-	-	-	-	-	-
87.0	80.0	0.0	-	-	0.0	-	-	-	-	-	-	-
87.0	90.0	0.0	-	-	0.0	-	-	-	-	-	-	-
90.0	90.0	0.0	-	-	0.0	-	-	-	-	-	-	-
90.0	120.0	0.0	-	-	6.4	-	-	-	-	-	-	-
90.0	200.0	2.8	-	-	-	-	-	-	-	-	-	-
93.0	75.0	-	-	-	-	-	-	-	-	-	-	-
93.0	80.0	3.0	-	-	-	-	-	-	-	-	-	-
93.0	85.0	-	-	-	-	-	-	-	-	-	-	-
93.0	90.0	0.0	-	-	-	-	-	-	-	-	-	-
97.0	40.0	0.0	-	-	0.0	-	-	-	-	-	-	-
97.0	65.0	3.1	-	-	0.0	-	-	-	-	-	-	-
97.0	75.0	-	-	-	-	-	-	-	-	-	-	-
97.0	80.0	0.0	-	-	-	-	-	-	-	-	-	-
100.0	40.0	0.0	-	-	-	-	-	-	-	-	-	-
100.0	45.0	0.0	-	-	-	-	-	-	-	-	-	-

TABLE 4. (cont.)

*Lestidiops ringens* (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
100.0	50.0	0.0	-	-	3.0	-	-	0.0	-	0.0	-	-
100.0	55.0	0.0	-	-	0.0	-	-	0.0	-	3.0	-	-
100.0	65.0	0.0	-	-	0.0	-	-	0.0	-	6.0	-	-
100.0	70.0	0.0	-	-	0.0	-	-	0.0	-	3.1	-	-
103.0	45.0	0.0	-	-	0.0	-	-	0.0	-	3.2	-	-
103.0	55.0	0.0	-	-	0.0	-	-	0.0	-	6.2	-	-
103.0	60.0	0.0	-	-	0.0	-	-	0.0	-	6.2	-	-
103.0	65.0	2.8	-	-	0.0	-	-	0.0	-	0.0	-	-
103.0	70.0	0.0	-	-	0.0	-	-	0.0	-	5.8	-	-
103.0	85.0	-	-	-	5.8	-	-	-	-	-	-	-
107.0	50.0	-	0.0	-	0.0	-	-	0.0	-	3.2	-	-
107.0	55.0	-	0.0	-	0.0	-	-	0.0	-	3.1	-	-
107.0	70.0	-	3.2	-	3.1	-	-	0.0	-	0.0	-	-
110.0	35.0	-	0.0	-	0.0	-	-	-	-	2.8	0.0	-
110.0	40.0	-	0.0	-	0.0	-	-	-	-	5.6	0.0	-
110.0	60.0	-	2.7	-	3.0	-	-	-	-	0.0	0.0	-
110.0	80.0	-	0.0	-	3.0	-	-	0.0	-	-	-	-
113.0	75.0	-	-	-	3.3	-	-	-	-	-	-	-
117.0	50.0	-	3.2	-	0.0	-	-	0.0	-	0.0	-	-
117.0	70.0	-	3.1	-	0.0	-	-	-	-	-	-	-
120.0	90.0	-	5.8	-	-	-	-	-	-	-	-	-
130.0	80.0	-	2.8	-	-	-	-	-	-	0.0	-	-

*Notolepis risso*

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
60.0	140.0	14.2	-	-	-	-	-	0.0	-	-	-	-
60.0	180.0	0.0	-	-	-	-	-	2.7	-	-	-	-
60.0	200.0	3.1	-	-	-	-	-	0.0	-	-	-	-
87.0	60.0	0.0	-	-	2.4	-	-	0.0	-	0.0	-	-
90.0	160.0	0.0	-	-	-	-	-	2.6	-	-	-	-
93.0	60.0	0.0	-	-	0.0	-	-	3.3	-	0.0	-	-
93.0	65.0	0.0	-	-	0.0	-	-	3.3	-	0.0	-	-
103.0	70.0	3.0	-	-	0.0	-	-	0.0	-	0.0	-	-
107.0	70.0	-	3.2	-	0.0	-	-	0.0	-	0.0	-	-

*Paralepis atlantica*

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
80.0	200.0	2.8	-	-	-	-	-	0.0	-	-	-	-

TABLE 4. (cont.)

*Stemonosudis macrura*

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
90.0	160.0	0.0	-	-	-	-	-	2.6	-	-	-	-
90.0	180.0	0.0	-	-	-	-	-	2.8	-	-	-	-
127.0	65.0	-	0.0	-	0.0	-	-	2.9	-	-	-	-
127.0	70.0	-	0.0	-	0.0	-	-	2.8	-	-	-	-
133.0	65.0	-	0.0	-	0.0	-	-	2.6	-	-	-	-
133.0	70.0	-	0.0	-	0.0	-	-	2.7	-	-	-	-

*Sudis atrox*

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
80.0	200.0	0.0	-	-	-	-	-	5.6	-	-	-	-
90.0	180.0	0.0	-	-	-	-	-	5.6	-	-	-	-
110.0	160.0	-	2.8	-	-	-	-	2.9	-	-	-	-

*Scopelosaurus spp.*

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
60.0	140.0	0.0	-	-	-	-	-	2.6	-	-	-	-
60.0	200.0	0.0	-	-	-	-	-	2.7	-	-	-	-
70.0	100.0	0.0	-	-	-	-	-	5.6	-	-	-	-
70.0	120.0	0.0	-	-	-	-	-	2.7	-	-	-	-
73.0	60.0	0.0	-	-	0.0	-	-	2.8	-	-	-	-
80.0	80.0	3.0	-	-	0.0	-	-	0.0	-	-	0.0	-
80.0	120.0	0.0	-	-	-	-	-	5.9	-	-	-	-
87.0	80.0	0.0	-	-	0.0	-	-	0.0	-	-	3.0	-
90.0	100.0	2.8	-	-	0.0	-	-	0.0	-	-	-	-
93.0	80.0	0.0	-	-	3.2	-	-	0.0	-	-	0.0	-
93.0	90.0	0.0	-	-	3.4	-	-	0.0	-	-	0.0	-
93.0	100.0	-	-	-	2.8	-	-	-	-	-	-	-
100.0	70.0	0.0	-	-	0.0	-	-	2.3	-	-	0.0	-
100.0	85.0	-	-	-	3.5	-	-	-	-	-	-	-
100.0	90.0	0.0	-	-	3.3	-	-	0.0	-	-	0.0	-
100.0	120.0	-	3.0	-	-	-	-	0.0	-	-	-	-
103.0	80.0	-	0.0	-	3.0	-	-	0.0	-	-	-	-
103.0	90.0	-	0.0	-	3.2	-	-	0.0	-	-	-	-
107.0	80.0	-	0.0	-	0.0	-	-	3.2	-	-	-	-

*Scopelarchidae*

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
70.0	200.0	6.1	-	-	-	-	-	0.0	-	-	-	-
80.0	200.0	0.0	-	-	-	-	-	5.6	-	-	-	-
83.0	43.0	0.0	-	-	2.8	-	-	0.0	-	-	0.0	-

TABLE 4. (cont.)

## Scopelarchidae (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
83.0	90.0	0.0	-	-	-	3.2	-	-	-	-	-	-
90.0	65.0	0.0	-	-	-	2.9	-	0.0	-	0.0	-	-
90.0	160.0	2.6	-	-	-	0.0	-	0.0	-	2.9	-	-
93.0	80.0	0.0	-	-	-	0.0	-	3.3	-	0.0	-	-
97.0	50.0	0.0	-	-	-	0.0	-	3.2	-	2.9	-	-
97.0	80.0	0.0	-	-	-	0.0	-	0.0	-	2.9	-	-
100.0	60.0	0.0	-	-	-	0.0	-	0.0	-	2.9	-	-
100.0	70.0	0.0	-	-	-	0.0	-	0.0	-	3.1	-	-
103.0	70.0	0.0	-	-	-	0.0	-	3.4	-	0.0	-	-
107.0	90.0	-	-	-	-	0.0	-	0.0	-	-	-	-
110.0	40.0	-	-	-	-	0.0	-	-	-	2.8	0.0	-
110.0	120.0	-	-	-	-	5.2	-	-	-	5.4	-	-
110.0	140.0	-	-	-	-	8.1	-	-	-	0.0	-	-
117.0	90.0	-	-	-	-	2.9	-	0.0	-	-	-	-
123.0	50.0	-	-	-	-	0.0	-	-	-	5.7	0.0	-
127.0	45.0	-	-	-	-	6.0	-	-	-	0.0	3.1	-
127.0	50.0	-	-	-	-	0.0	-	-	-	0.0	3.0	-
127.0	70.0	-	-	-	-	0.0	-	-	-	2.8	-	-
127.0	80.0	-	-	-	-	0.0	-	-	-	8.2	-	-
130.0	60.0	-	-	-	-	0.0	-	-	-	2.8	0.0	-
130.0	70.0	-	-	-	-	2.9	-	-	-	0.0	-	-
130.0	100.0	-	-	-	-	5.5	-	-	-	0.0	-	-
130.0	120.0	-	-	-	-	2.8	-	-	-	2.7	-	-
133.0	70.0	-	-	-	-	0.0	-	-	-	0.0	-	-
133.0	80.0	-	-	-	-	3.0	-	-	-	5.1	-	-

## Myctophidae

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
60.0	51.0	0.0	-	-	-	-	-	-	2.0	-	-	-
60.0	100.0	6.2	-	-	-	-	-	-	0.0	-	-	-
60.0	140.0	0.0	-	-	-	-	-	-	5.2	-	-	-
60.0	200.0	0.0	-	-	-	-	-	-	27.4	-	-	-
63.0	55.0	0.0	-	-	-	-	-	-	2.8	-	-	-
67.0	49.0	3.2	-	-	-	-	-	-	0.0	-	-	-
70.0	70.0	2.7	-	-	-	-	-	-	0.0	-	-	-
70.0	90.0	0.0	-	-	-	-	-	-	8.3	-	-	-
70.0	100.0	0.0	-	-	-	-	-	-	2.8	-	-	-
70.0	120.0	3.0	-	-	-	-	-	-	2.7	-	-	-
70.0	200.0	0.0	-	-	-	-	-	-	5.3	-	-	-
73.0	60.0	2.8	-	-	-	-	-	-	0.0	-	-	-
73.0	65.0	-	-	-	-	-	-	-	2.9	-	-	-
77.0	70.0	-	-	-	-	-	-	-	2.3	-	-	-
77.0	80.0	-	-	-	-	-	-	-	2.8	-	-	-
80.0	51.0	0.0	-	-	-	-	-	-	0.0	-	-	1.6

TABLE 4. (cont.)

## Myctophidae (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
80.0	65.0	0.0	-	-	0.0	-	-	5.9	0.0	-	-	-
80.0	90.0	5.8	-	-	-	-	-	0.0	-	-	-	-
80.0	100.0	3.0	-	-	-	-	-	3.0	-	-	-	-
80.0	120.0	5.9	-	-	-	-	-	0.0	-	-	-	-
80.0	160.0	5.8	-	-	-	-	-	-	-	-	-	-
80.0	200.0	8.5	-	-	-	-	-	8.4	-	-	-	-
83.0	60.0	0.0	-	-	0.0	-	-	-	-	-	-	-
83.0	65.0	0.0	-	-	2.9	-	-	-	-	-	-	-
83.0	70.0	0.0	-	-	3.2	-	-	-	-	-	-	-
83.0	80.0	0.0	-	-	2.9	-	-	-	-	-	-	-
83.0	90.0	0.0	-	-	0.0	-	-	-	-	-	-	-
87.0	34.0	0.0	-	-	0.0	-	-	2.6	-	-	-	-
87.0	35.0	0.0	-	-	0.0	-	-	12.3	-	-	-	-
87.0	60.0	3.0	-	-	0.0	-	-	13.3	-	-	-	-
87.0	65.0	3.1	-	-	0.0	-	-	0.0	-	-	-	-
87.0	80.0	0.0	-	-	3.0	-	-	3.3	-	-	-	-
87.0	90.0	3.0	-	-	0.0	-	-	6.2	-	-	-	-
87.0	100.0	-	-	-	6.2	-	-	-	-	-	-	-
90.0	53.0	0.0	-	-	6.6	-	-	0.0	-	-	-	-
90.0	65.0	6.5	-	-	5.9	-	-	0.0	-	-	-	-
90.0	70.0	0.0	-	-	0.0	-	-	5.5	-	-	-	-
90.0	80.0	0.0	-	-	6.4	-	-	0.0	-	-	-	-
90.0	90.0	0.0	-	-	9.6	-	-	0.0	-	-	-	-
90.0	100.0	5.5	-	-	0.0	-	-	5.5	-	-	-	-
90.0	160.0	0.0	-	-	-	-	-	-	-	-	-	-
90.0	180.0	0.0	-	-	-	-	-	-	-	-	-	-
90.0	200.0	2.8	-	-	-	-	-	-	-	-	-	-
93.0	28.0	0.0	-	-	3.3	-	-	0.0	-	-	-	-
93.0	35.0	0.0	-	-	2.9	-	-	0.0	-	-	-	-
93.0	40.0	0.0	-	-	-	-	-	0.0	-	-	-	-
93.0	55.0	0.0	-	-	-	-	-	6.8	-	-	-	-
93.0	60.0	0.0	-	-	-	-	-	0.0	-	-	-	-
93.0	70.0	0.0	-	-	-	-	-	3.3	-	-	-	-
93.0	80.0	0.0	-	-	-	-	-	3.2	-	-	-	-
93.0	90.0	0.0	-	-	-	-	-	3.4	-	-	-	-
93.0	100.0	-	-	-	-	-	-	2.8	-	-	-	-
97.0	32.0	0.0	-	-	-	-	-	-	-	-	-	-
97.0	35.0	0.0	-	-	3.2	-	-	0.0	-	-	-	-
97.0	40.0	0.0	-	-	3.2	-	-	0.0	-	-	-	-
97.0	45.0	3.0	-	-	10.4	-	-	0.0	-	-	-	-
97.0	50.0	6.3	-	-	3.2	-	-	0.0	-	-	-	-
97.0	55.0	3.0	-	-	-	-	-	-	-	-	-	-
97.0	80.0	3.3	-	-	-	-	-	-	-	-	-	-
97.0	85.0	-	-	-	-	-	-	-	-	-	-	-
97.0	100.0	35.0	0.0	-	-	-	-	-	-	-	-	-

TABLE 4. (cont.)

## Myctophidae (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
100.0	45.0	0.0	-	-	0.0	-	-	2.9	-	0.0	-	-
100.0	50.0	0.0	-	-	3.0	-	-	0.0	-	3.0	-	-
100.0	65.0	0.0	-	-	0.0	-	-	3.1	-	0.0	-	-
100.0	70.0	3.1	-	-	0.0	-	-	0.0	-	3.1	-	-
100.0	75.0	-	-	-	3.4	-	-	-	-	-	-	-
100.0	80.0	5.8	-	-	3.3	-	-	-	-	-	-	-
100.0	85.0	-	-	-	6.9	-	-	-	-	-	-	-
100.0	90.0	0.0	-	-	9.9	-	-	0.0	-	0.0	-	-
100.0	100.0	-	5.9	-	-	-	-	0.0	-	3.1	-	-
103.0	45.0	0.0	-	-	9.1	-	-	-	-	2.9	-	-
103.0	60.0	0.0	-	-	0.0	-	-	-	-	-	-	-
103.0	70.0	0.0	-	-	3.1	-	-	-	-	-	-	-
103.0	80.0	-	3.0	-	0.0	-	-	-	-	-	-	-
103.0	90.0	-	0.0	-	6.3	-	-	-	-	-	-	-
105.0	140.0	-	2.8	-	-	-	-	0.0	-	-	-	-
107.0	31.0	-	0.0	-	5.7	-	-	-	-	-	-	-
107.0	32.0	-	0.0	-	3.1	-	-	-	-	-	-	-
107.0	45.0	-	0.0	-	3.0	-	-	-	-	-	-	-
107.0	50.0	-	0.0	-	0.0	-	-	-	-	-	-	-
107.0	60.0	-	3.0	-	3.2	-	-	-	-	-	-	-
107.0	65.0	-	0.0	-	0.0	-	-	-	-	-	-	-
107.0	70.0	-	9.5	-	0.0	-	-	-	-	-	-	-
107.0	80.0	-	0.0	-	3.0	-	-	-	-	-	-	-
107.0	90.0	-	5.9	-	3.1	-	-	-	-	-	-	-
110.0	35.0	-	0.0	-	0.0	-	-	-	-	-	-	-
110.0	40.0	-	0.0	-	0.0	-	-	-	-	-	-	-
110.0	50.0	-	2.8	-	0.0	-	-	-	-	-	-	-
110.0	55.0	-	0.0	-	3.0	-	-	-	-	-	-	-
110.0	60.0	-	0.0	-	6.1	-	-	-	-	-	-	-
110.0	70.0	-	5.6	-	0.0	-	-	-	-	-	-	-
110.0	90.0	-	0.0	-	0.0	-	-	-	-	-	-	-
110.0	100.0	-	0.0	-	-	-	-	-	-	-	-	-
110.0	120.0	-	2.6	-	-	-	-	-	-	-	-	-
110.0	160.0	-	14.0	-	-	-	-	-	-	-	-	-
113.0	40.0	-	0.0	-	0.0	-	-	-	-	-	-	-
113.0	45.0	-	0.0	-	0.0	-	-	-	-	-	-	-
113.0	50.0	-	0.0	-	3.1	-	-	-	-	-	-	-
113.0	55.0	-	0.0	-	0.0	-	-	-	-	-	-	-
113.0	60.0	-	0.0	-	0.0	-	-	-	-	-	-	-
113.0	65.0	-	0.0	-	0.0	-	-	-	-	-	-	-
113.0	70.0	-	0.0	-	3.1	-	-	-	-	-	-	-
113.0	80.0	-	0.0	-	6.2	-	-	-	-	-	-	-
113.0	90.0	-	0.0	-	0.0	-	-	-	-	-	-	-
117.0	40.0	-	3.0	-	-	-	-	-	-	-	-	-
117.0	45.0	-	0.0	-	2.9	-	-	-	-	-	-	-
117.0	50.0	-	0.0	-	0.0	-	-	-	-	-	-	-

TABLE 4. (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.	Myctophidae (cont.)												
													117.0	117.0	117.0	117.0	117.0	117.0	117.0	117.0	117.0	117.0	117.0	117.0	117.0
117.0	55.0	-	0.0	-	0.0	-	-	-	6.6	-	5.8	-	-	-	-	-	-	-	-	-	-	-	-	-	-
117.0	75.0	-	0.0	-	3.0	-	6.1	0.0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
117.0	80.0	-	0.0	-	8.7	-	5.6	0.0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
117.0	85.0	-	0.0	-	0.0	-	0.0	0.0	-	-	6.0	6.0	-	-	-	-	-	-	-	-	-	-	-	-	-
117.0	90.0	-	0.0	-	0.0	-	0.0	0.0	-	-	5.1	5.1	-	-	-	-	-	-	-	-	-	-	-	-	-
120.0	50.0	-	0.0	-	0.0	-	0.0	0.0	-	-	0.0	0.0	-	-	-	-	-	-	-	-	-	-	-	-	-
120.0	55.0	-	0.0	-	0.0	-	0.0	0.0	-	-	8.0	8.0	-	-	-	-	-	-	-	-	-	-	-	-	-
120.0	60.0	-	0.0	-	0.0	-	0.0	0.0	-	-	10.8	10.8	-	-	-	-	-	-	-	-	-	-	-	-	-
120.0	65.0	-	0.0	-	0.0	-	0.0	0.0	-	-	5.7	5.7	-	-	-	-	-	-	-	-	-	-	-	-	-
120.0	75.0	-	0.0	-	0.0	-	0.0	0.0	-	-	0.0	0.0	-	-	-	-	-	-	-	-	-	-	-	-	-
120.0	80.0	-	0.0	-	0.0	-	0.0	0.0	-	-	0.0	0.0	-	-	-	-	-	-	-	-	-	-	-	-	-
123.0	50.0	-	0.0	-	0.0	-	0.0	0.0	-	-	0.0	0.0	-	-	-	-	-	-	-	-	-	-	-	-	-
123.0	55.0	-	0.0	-	0.0	-	0.0	0.0	-	-	3.1	3.1	-	-	-	-	-	-	-	-	-	-	-	-	-
123.0	65.0	-	0.0	-	0.0	-	0.0	0.0	-	-	0.0	0.0	-	-	-	-	-	-	-	-	-	-	-	-	-
123.0	70.0	-	0.0	-	0.0	-	0.0	0.0	-	-	18.5	18.5	-	-	-	-	-	-	-	-	-	-	-	-	-
123.0	80.0	-	0.0	-	0.0	-	0.0	0.0	-	-	3.1	3.1	-	-	-	-	-	-	-	-	-	-	-	-	-
127.0	33.0	-	2.8	-	2.7	-	0.0	0.0	-	-	0.0	0.0	-	-	-	-	-	-	-	-	-	-	-	-	-
127.0	34.0	-	2.6	-	2.6	-	0.0	0.0	-	-	0.0	0.0	-	-	-	-	-	-	-	-	-	-	-	-	-
127.0	40.0	-	7.2	-	7.2	-	5.9	5.9	-	-	0.0	0.0	-	-	-	-	-	-	-	-	-	-	-	-	-
127.0	45.0	-	0.0	-	0.0	-	6.2	6.2	-	-	0.0	0.0	-	-	-	-	-	-	-	-	-	-	-	-	-
127.0	50.0	-	0.0	-	0.0	-	3.0	3.0	-	-	0.0	0.0	-	-	-	-	-	-	-	-	-	-	-	-	-
127.0	55.0	-	0.0	-	0.0	-	8.9	8.9	-	-	0.0	0.0	-	-	-	-	-	-	-	-	-	-	-	-	-
127.0	65.0	-	0.0	-	0.0	-	2.9	2.9	-	-	11.4	11.4	-	-	-	-	-	-	-	-	-	-	-	-	-
127.0	70.0	-	0.0	-	0.0	-	2.9	2.9	-	-	5.5	5.5	-	-	-	-	-	-	-	-	-	-	-	-	-
127.0	80.0	-	0.0	-	0.0	-	0.0	0.0	-	-	0.0	0.0	-	-	-	-	-	-	-	-	-	-	-	-	-
130.0	25.5	-	0.0	-	0.0	-	9.5	9.5	-	-	0.0	0.0	-	-	-	-	-	-	-	-	-	-	-	-	-
130.0	40.0	-	0.0	-	8.4	-	0.0	0.0	-	-	16.3	16.3	-	-	-	-	-	-	-	-	-	-	-	-	-
130.0	45.0	-	0.0	-	0.0	-	5.9	5.9	-	-	16.3	16.3	-	-	-	-	-	-	-	-	-	-	-	-	-
130.0	50.0	-	0.0	-	0.0	-	8.8	8.8	-	-	16.3	16.3	-	-	-	-	-	-	-	-	-	-	-	-	-
130.0	55.0	-	0.0	-	0.0	-	0.0	0.0	-	-	12.4	12.4	-	-	-	-	-	-	-	-	-	-	-	-	-
130.0	60.0	-	0.0	-	0.0	-	3.1	3.1	-	-	0.0	0.0	-	-	-	-	-	-	-	-	-	-	-	-	-
130.0	80.0	-	0.0	-	6.2	-	9.1	9.1	-	-	0.0	0.0	-	-	-	-	-	-	-	-	-	-	-	-	-
130.0	90.0	-	0.0	-	11.3	-	0.0	0.0	-	-	0.0	0.0	-	-	-	-	-	-	-	-	-	-	-	-	-
130.0	120.0	-	0.0	-	16.6	-	12.4	12.4	-	-	0.0	0.0	-	-	-	-	-	-	-	-	-	-	-	-	-
133.0	30.0	-	0.0	-	0.0	-	0.0	0.0	-	-	0.0	0.0	-	-	-	-	-	-	-	-	-	-	-	-	-
133.0	40.0	-	0.0	-	0.0	-	6.2	6.2	-	-	0.0	0.0	-	-	-	-	-	-	-	-	-	-	-	-	-
133.0	45.0	-	0.0	-	0.0	-	9.1	9.1	-	-	0.0	0.0	-	-	-	-	-	-	-	-	-	-	-	-	-
133.0	50.0	-	0.0	-	11.3	-	0.0	0.0	-	-	0.0	0.0	-	-	-	-	-	-	-	-	-	-	-	-	-
133.0	55.0	-	0.0	-	0.0	-	0.0	0.0	-	-	0.0	0.0	-	-	-	-	-	-	-	-	-	-	-	-	-
133.0	65.0	-	0.0	-	8.6	-	12.4	12.4	-	-	0.0	0.0	-	-	-	-	-	-	-	-	-	-	-	-	-
133.0	70.0	-	0.0	-	2.8	-	0.0	0.0	-	-	0.0	0.0	-	-	-	-	-	-	-	-	-	-	-	-	-
133.0	80.0	-	0.0	-	0.0	-	0.0	0.0	-	-	0.0	0.0	-	-	-	-	-	-	-	-	-	-	-	-	-
133.0	35.0	-	0.0	-	3.0	-	0.0	0.0	-	-	0.0	0.0	-	-	-	-	-	-	-	-	-	-	-	-	-
137.0	45.0	-	0.0	-	0.0	-	0.0	0.0	-	-	0.0	0.0	-	-	-	-	-	-	-	-	-	-	-	-	-
137.0	50.0	-	0.0	-	2.7	-	0.0	0.0	-	-	0.0	0.0	-	-	-	-	-	-	-	-	-	-	-	-	-
137.0	55.0	-	0.0	-	0.0	-	0.0	0.0	-	-	0.0	0.0	-	-	-	-	-	-	-	-	-	-	-	-	-

TABLE 4. (cont.)

## Myctophidae (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
137.0	60.0	-	-	-	0.0	-	-	-	2.2	-	-	-
137.0	65.0	-	-	-	12.9	-	-	-	-	-	-	-
137.0	70.0	-	6.0	-	0.0	-	-	-	0.0	-	-	-
137.0	80.0	-	3.0	-	-	-	-	-	5.0	-	-	-
140.0	28.8	-	0.7	-	-	-	-	-	0.0	-	-	-
140.0	50.0	-	11.8	-	-	-	-	-	9.2	-	-	-

*Ceratoscopelus townsendi*

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
60.0	140.0	3.5	-	-	-	-	-	-	33.9	-	-	-
60.0	160.0	6.5	-	-	-	-	-	-	2.9	-	-	-
60.0	180.0	11.8	-	-	-	-	-	-	10.8	-	-	-
60.0	200.0	9.3	-	-	-	-	-	-	142.5	-	-	-
70.0	51.0	0.0	-	-	0.0	-	-	-	2.9	-	-	-
70.0	80.0	5.6	-	-	0.0	-	-	-	0.0	-	-	-
70.0	90.0	0.0	-	-	-	-	-	-	24.8	-	-	-
70.0	100.0	0.0	-	-	-	-	-	-	2.8	-	-	-
70.0	120.0	6.0	-	-	-	-	-	-	21.9	-	-	-
70.0	160.0	17.8	-	-	-	-	-	-	-	-	-	-
70.0	180.0	11.9	-	-	-	-	-	-	-	-	-	-
70.0	200.0	30.6	-	-	-	-	-	-	18.6	-	-	-
73.0	60.0	0.0	-	-	0.0	-	-	-	2.8	-	-	-
80.0	60.0	0.0	-	-	0.0	-	-	-	2.8	-	-	-
80.0	75.0	-	-	-	-	-	-	-	0.0	-	-	-
80.0	100.0	3.0	-	-	-	-	-	-	0.0	-	-	-
80.0	140.0	16.4	-	-	-	-	-	-	-	-	-	-
80.0	160.0	157.7	-	-	-	-	-	-	95.5	-	-	-
80.0	180.0	24.7	-	-	-	-	-	-	-	-	-	-
80.0	200.0	22.6	-	-	-	-	-	-	-	-	-	-
83.0	65.0	0.0	-	-	2.9	-	-	-	0.0	-	-	-
87.0	70.0	0.0	-	-	9.6	-	-	-	0.0	-	-	-
87.0	80.0	0.0	-	-	3.0	-	-	-	0.0	-	-	-
87.0	90.0	0.0	-	-	0.0	-	-	-	46.8	-	-	-
87.0	100.0	-	-	-	3.1	-	-	-	-	-	-	-
90.0	65.0	0.0	-	-	2.9	-	-	-	2.7	-	-	-
90.0	70.0	0.0	-	-	0.0	-	-	-	0.0	-	-	-
90.0	80.0	0.0	-	-	0.0	-	-	-	0.0	-	-	-
90.0	90.0	0.0	-	-	0.0	-	-	-	13.1	-	-	-
90.0	100.0	2.8	-	-	0.0	-	-	-	2.3	-	-	-
90.0	120.0	21.7	-	-	0.0	-	-	-	5.5	-	-	-
90.0	140.0	0.0	-	-	0.0	-	-	-	88.4	-	-	-
90.0	160.0	13.0	-	-	0.0	-	-	-	67.0	-	-	-
90.0	180.0	16.4	-	-	0.0	-	-	-	57.0	-	-	-
90.0	200.0	33.7	-	-	-	-	-	-	120.0	-	-	-
									24.8			

TABLE 4. (cont.)

*Ceratoscopelus townsendi* (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
93.0	85.0	-	0.0	-	-	6.7	0.0	-	-	26.9	-	-
93.0	90.0	0.0	-	-	8.5	-	9.7	-	-	2.8	-	-
93.0	100.0	-	0.0	-	-	7.0	-	-	-	-	58.0	-
97.0	80.0	0.0	-	-	0.0	-	-	-	2.8	0.0	-	-
97.0	100.0	-	0.0	-	0.0	-	-	-	3.1	3.0	-	-
100.0	40.0	0.0	-	-	0.0	-	-	-	0.0	15.4	-	-
100.0	65.0	0.0	-	-	0.0	-	-	-	-	-	-	-
100.0	70.0	3.1	-	-	0.0	-	-	-	-	-	-	-
100.0	85.0	-	-	-	17.3	-	-	-	-	-	-	-
100.0	90.0	0.0	-	-	19.9	-	-	-	2.9	-	-	-
100.0	100.0	-	-	3.0	-	-	-	-	9.0	-	-	-
100.0	120.0	-	-	3.0	-	-	-	-	15.0	-	-	-
103.0	60.0	0.0	-	-	0.0	-	-	-	-	15.4	-	-
103.0	65.0	0.0	-	-	0.0	-	-	-	-	16.0	-	-
103.0	70.0	0.0	-	-	0.0	-	-	-	-	0.0	-	-
103.0	75.0	-	-	-	-	3.2	-	-	-	-	-	-
103.0	80.0	-	-	-	0.0	-	-	-	3.0	-	-	-
103.0	85.0	-	-	-	17.4	-	-	-	-	-	-	-
103.0	90.0	-	-	-	3.2	-	-	-	-	-	-	-
105.0	140.0	-	-	-	27.8	-	-	-	16.7	-	-	-
107.0	60.0	-	-	-	0.0	-	-	-	-	3.0	-	-
107.0	65.0	-	-	-	0.0	-	-	-	-	26.2	-	-
107.0	70.0	-	-	-	0.0	-	-	-	-	15.1	-	-
107.0	90.0	-	-	-	0.0	-	-	-	-	-	-	-
110.0	40.0	-	-	-	27.8	-	-	-	-	-	-	-
110.0	45.0	-	-	-	0.0	-	-	-	-	-	-	-
110.0	60.0	-	-	-	0.0	-	-	-	-	-	-	-
110.0	65.0	-	-	-	0.0	-	-	-	-	-	-	-
110.0	75.0	-	-	-	3.2	-	-	-	-	-	-	-
110.0	80.0	-	-	-	7.7	-	-	-	-	-	-	-
110.0	90.0	-	-	-	2.6	-	-	-	-	-	-	-
110.0	100.0	-	-	-	10.4	-	-	-	-	-	-	-
110.0	120.0	-	-	-	2.6	-	-	-	-	-	-	-
110.0	140.0	-	-	-	13.5	-	-	-	-	-	-	-
110.0	160.0	-	-	-	5.6	-	-	-	-	-	-	-
113.0	35.0	-	-	-	0.0	-	-	-	3.3	-	-	-
113.0	45.0	-	-	-	0.0	-	-	-	0.0	4.9	-	-
113.0	50.0	-	-	-	6.2	-	-	-	0.0	30.0	-	-
113.0	65.0	-	-	-	0.0	-	-	-	6.5	-	2.8	-
113.0	90.0	-	-	-	0.0	-	-	-	0.0	-	8.7	-
117.0	50.0	-	-	-	0.0	-	-	-	0.0	-	15.1	-
117.0	60.0	-	-	-	0.0	-	-	-	0.0	-	-	-
117.0	80.0	-	-	-	0.0	-	-	-	0.0	-	-	-
117.0	85.0	-	-	-	3.1	-	-	-	3.1	-	-	-
117.0	90.0	-	-	-	5.8	-	-	-	5.8	-	-	-
117.0	90.0	-	-	-	0.0	-	-	-	0.0	-	-	-

TABLE 4. (cont.)

*Ceratoscopelus townsendi* (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
120.0	55.0	-	0.0	-	0.0	-	-	-	46.1	0.0	-	-
120.0	60.0	-	0.0	-	0.0	-	-	-	14.1	0.0	-	-
120.0	65.0	-	0.0	-	0.0	-	-	-	2.7	-	-	-
120.0	80.0	-	0.0	-	0.0	-	-	-	13.5	-	-	-
120.0	100.0	-	14.9	-	-	-	-	-	15.6	-	-	-
120.0	120.0	-	0.0	-	-	-	-	-	24.8	-	-	-
123.0	50.0	-	0.0	-	0.0	-	-	-	0.0	-	-	-
123.0	60.0	-	0.0	-	0.0	-	-	-	0.0	-	-	-
123.0	70.0	-	0.0	-	0.0	-	-	-	2.7	-	-	-
123.0	80.0	-	0.0	-	0.0	-	-	-	2.8	-	-	-
127.0	55.0	-	0.0	-	0.0	-	-	-	2.8	-	-	-
127.0	60.0	-	0.0	-	0.0	-	-	-	8.2	-	-	-
127.0	65.0	-	0.0	-	0.0	-	-	-	0.0	-	-	-
127.0	70.0	-	0.0	-	0.0	-	-	-	0.0	-	-	-
127.0	80.0	-	0.0	-	0.0	-	-	-	11.4	-	-	-
130.0	40.0	-	0.0	-	0.0	-	-	-	24.8	-	-	-
130.0	50.0	-	0.0	-	0.0	-	-	-	0.0	-	-	-
130.0	55.0	-	0.0	-	0.0	-	-	-	0.0	-	-	-
130.0	60.0	-	0.0	-	0.0	-	-	-	0.0	-	-	-
130.0	100.0	-	0.0	-	0.0	-	-	-	185.6	-	-	-
130.0	120.0	-	0.0	-	0.0	-	-	-	2.6	-	-	-
133.0	45.0	-	0.0	-	0.0	-	-	-	5.5	-	-	-
133.0	50.0	-	0.0	-	0.0	-	-	-	17.3	-	-	-
133.0	55.0	-	0.0	-	0.0	-	-	-	19.9	-	-	-
133.0	60.0	-	0.0	-	0.0	-	-	-	19.9	-	-	-
133.0	100.0	-	0.0	-	0.0	-	-	-	31.2	-	-	-
133.0	120.0	-	0.0	-	0.0	-	-	-	2.7	-	-	-
133.0	145.0	-	0.0	-	0.0	-	-	-	5.3	-	-	-
133.0	50.0	-	0.0	-	0.0	-	-	-	2.6	-	-	-
133.0	55.0	-	0.0	-	0.0	-	-	-	2.7	-	-	-
133.0	60.0	-	0.0	-	0.0	-	-	-	2.8	-	-	-
133.0	65.0	-	0.0	-	0.0	-	-	-	10.6	-	-	-
133.0	70.0	-	0.0	-	0.0	-	-	-	5.4	-	-	-
137.0	45.0	-	0.0	-	0.0	-	-	-	0.0	-	-	-
137.0	50.0	-	0.0	-	0.0	-	-	-	7.8	-	-	-
137.0	60.0	-	0.0	-	0.0	-	-	-	4.3	-	-	-
137.0	65.0	-	0.0	-	0.0	-	-	-	-	-	-	-
140.0	50.0	-	0.0	-	0.0	-	-	-	2.3	-	-	-

*Diaphus* spp.

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
60.0	90.0	0.0	-	-	-	-	-	-	5.6	-	-	-
60.0	100.0	0.0	-	-	-	-	-	-	28.9	-	-	-
60.0	120.0	0.0	-	-	-	-	-	-	2.7	-	-	-
60.0	140.0	0.0	-	-	-	-	-	-	15.7	-	-	-
60.0	200.0	0.0	-	-	-	-	-	-	2.7	-	-	-
63.0	55.0	0.0	-	-	-	-	-	-	2.8	-	-	-
63.0	60.0	0.0	-	-	-	-	-	-	5.7	-	-	-
67.0	55.0	0.0	-	-	-	-	-	-	14.4	-	-	-
70.0	51.0	0.0	-	-	-	-	-	-	2.9	-	-	-

TABLE 4. (cont.)

*Diaphus* spp. (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
70.0	80.0	0.0	-	-	0.0	-	-	-	-	3.2	-	-
70.0	90.0	0.0	-	-	-	-	-	-	-	16.5	-	-
70.0	100.0	0.0	-	-	-	-	-	-	-	2.8	-	-
70.0	120.0	0.0	-	-	-	-	-	-	-	13.7	-	-
70.0	200.0	0.0	-	-	-	-	-	-	-	8.0	-	-
73.0	51.0	0.0	-	-	-	-	-	-	-	12.7	-	-
73.0	53.0	0.0	-	-	-	-	-	-	-	13.9	-	-
73.0	60.0	0.0	-	-	-	-	-	-	-	33.8	-	-
77.0	57.0	0.0	-	-	-	-	-	-	-	5.5	-	-
77.0	70.0	-	-	-	4.6	-	-	-	-	-	-	-
77.0	80.0	-	-	-	2.8	-	-	-	-	21.5	0.0	-
80.0	70.0	0.0	-	-	0.0	-	-	-	-	-	-	-
80.0	80.0	0.0	-	-	0.0	-	-	-	-	-	-	-
80.0	90.0	0.0	-	-	-	-	-	-	-	-	-	-
80.0	100.0	0.0	-	-	-	-	-	-	-	-	-	-
80.0	120.0	0.0	-	-	-	-	-	-	-	-	-	-
80.0	140.0	2.7	-	-	-	-	-	-	-	-	-	-
80.0	200.0	0.0	-	-	-	-	-	-	-	-	-	-
82.0	47.0	0.0	-	-	-	3.1	-	-	-	-	-	-
83.0	51.0	0.0	-	-	-	0.0	-	-	-	-	-	-
83.0	60.0	0.0	-	-	-	3.0	-	-	-	-	-	-
83.0	65.0	0.0	-	-	-	0.0	-	-	-	-	-	-
83.0	70.0	0.0	-	-	-	0.0	-	-	-	-	-	-
83.0	80.0	0.0	-	-	-	0.0	-	-	-	-	-	-
83.0	90.0	0.0	-	-	-	0.0	-	-	-	-	-	-
87.0	34.0	0.0	-	-	-	0.0	-	-	-	-	-	-
87.0	35.0	0.0	-	-	-	0.0	-	-	-	-	-	-
87.0	40.0	0.0	-	-	-	0.0	-	-	-	-	-	-
87.0	60.0	0.0	-	-	-	0.0	-	-	-	-	-	-
87.0	65.0	0.0	-	-	-	0.0	-	-	-	-	-	-
87.0	70.0	0.0	-	-	-	0.0	-	-	-	-	-	-
87.0	80.0	0.0	-	-	-	0.0	-	-	-	-	-	-
90.0	85.0	0.0	-	-	-	0.0	-	-	-	-	-	-
90.0	90.0	0.0	-	-	-	0.0	-	-	-	-	-	-
90.0	100.0	0.0	-	-	-	0.0	-	-	-	-	-	-
90.0	120.0	0.0	-	-	-	0.0	-	-	-	-	-	-
90.0	140.0	0.0	-	-	-	0.0	-	-	-	-	-	-
90.0	160.0	0.0	-	-	-	0.0	-	-	-	-	-	-
90.0	180.0	0.0	-	-	-	0.0	-	-	-	-	-	-
93.0	27.0	0.0	-	-	-	0.0	-	-	-	-	-	-
93.0	50.0	0.0	-	-	-	0.0	-	-	-	-	-	-
93.0	55.0	0.0	-	-	-	0.0	-	-	-	-	-	-

TABLE 4. (cont.)

*Diaphus* spp. (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
93.0	60.0	0.0	-	0.0	-	3.3	-	-	0.0	-	-	-
93.0	70.0	0.0	-	0.0	-	3.3	-	-	0.0	-	-	-
93.0	75.0	-	-	-	0.0	12.6	-	-	-	2.9	-	-
93.0	80.0	0.0	-	-	-	19.6	-	-	0.0	-	-	-
93.0	85.0	-	-	0.0	-	9.5	-	-	9.0	-	-	-
93.0	90.0	0.0	-	0.0	-	6.6	-	-	3.0	-	-	-
97.0	40.0	0.0	-	0.0	-	0.0	-	-	0.0	-	-	-
97.0	45.0	0.0	-	0.0	-	-	-	-	0.0	-	-	-
97.0	50.0	0.0	-	0.0	-	-	-	-	0.0	-	-	-
97.0	75.0	-	-	-	0.0	-	-	-	-	3.2	-	-
97.0	80.0	0.0	-	-	-	-	-	-	-	5.8	-	-
100.0	35.0	0.0	-	0.0	-	0.0	-	-	0.0	-	-	-
100.0	65.0	0.0	-	0.0	-	-	-	-	3.0	-	-	-
103.0	30.0	0.0	-	0.0	-	5.3	-	-	0.0	-	-	-
103.0	40.0	0.0	-	0.0	-	3.3	-	-	0.0	-	-	-
103.0	45.0	0.0	-	0.0	-	10.1	-	-	0.0	-	-	-
103.0	50.0	0.0	-	0.0	-	11.0	-	-	0.0	-	-	-
103.0	55.0	0.0	-	0.0	-	0.0	-	-	6.2	-	-	-
103.0	60.0	0.0	-	0.0	-	6.6	-	-	0.0	-	-	-
110.0	140.0	-	0.0	-	-	-	-	-	2.8	-	-	-
110.0	160.0	-	0.0	-	-	-	-	-	2.9	-	-	-

*Lampadена urophaois*

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
60.0	180.0	0.0	-	-	-	-	-	-	2.7	-	-	-
60.0	200.0	0.0	-	-	-	-	-	-	11.0	-	-	-
70.0	200.0	0.0	-	-	-	-	-	-	2.7	-	-	-
80.0	180.0	5.5	-	-	-	-	-	-	8.4	-	-	-
80.0	200.0	2.8	-	-	-	-	-	-	5.4	-	-	-
90.0	120.0	0.0	-	-	-	-	-	-	11.2	-	-	-
90.0	180.0	0.0	-	-	-	-	-	-	5.5	-	-	-
90.0	200.0	0.0	-	-	-	-	-	-	-	16.4	-	-
97.0	85.0	-	-	3.6	-	-	-	-	-	5.6	-	-
107.0	65.0	-	0.0	-	0.0	0.0	-	-	-	0.0	-	-
110.0	40.0	-	0.0	-	0.0	-	-	-	-	0.1	-	-
110.0	65.0	-	0.0	-	0.0	-	-	-	-	5.4	-	-
110.0	120.0	-	0.0	-	-	-	-	-	-	5.5	-	-
110.0	140.0	-	0.0	-	-	-	-	-	-	8.8	-	-
110.0	160.0	-	0.0	-	-	-	-	-	-	-	3.0	-
113.0	50.0	-	0.0	-	0.0	-	-	-	-	-	5.9	-
113.0	55.0	-	0.0	-	0.0	-	-	-	-	-	2.9	-
113.0	60.0	-	0.0	-	0.0	-	-	-	-	-	-	3.0
113.0	85.0	-	3.0	-	-	-	-	-	-	-	-	0.0
113.0	90.0	-	0.0	-	0.0	-	-	-	-	-	-	0.0
117.0	60.0	-	-	-	-	-	-	-	-	-	-	3.0

TABLE 4. (cont.)

*Lampadena urophao*s (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
120.0	55.0	-	0.0	-	0.0	-	-	-	2.6	0.0	-	-
120.0	100.0	-	0.0	-	-	-	-	-	5.2	-	-	-
120.0	120.0	-	0.0	-	-	-	-	-	5.5	-	-	-
123.0	50.0	-	0.0	-	0.0	-	-	-	5.7	0.0	-	-
127.0	80.0	-	0.0	-	-	-	-	-	13.7	2.7	-	-
130.0	50.0	-	0.0	-	0.0	-	-	-	5.5	0.0	-	-
130.0	55.0	-	0.0	-	0.0	-	-	-	0.0	-	-	-
133.0	65.0	-	0.0	-	6.2	-	-	-	0.0	-	-	-
137.0	50.0	-	0.0	-	0.0	-	-	-	2.6	0.0	-	-
137.0	65.0	-	-	-	6.5	-	-	-	-	-	-	-

*Lampanyctus spp.*

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
60.0	140.0	0.0	-	-	-	-	-	-	5.2	-	-	-
60.0	160.0	0.0	-	-	-	-	-	-	2.9	-	-	-
60.0	180.0	5.9	-	-	-	-	-	-	0.0	-	-	-
60.0	200.0	6.2	-	-	-	-	-	-	30.1	-	-	-
70.0	160.0	35.6	-	-	-	-	-	-	0.0	-	-	-
70.0	200.0	15.3	-	-	-	-	-	-	0.0	-	-	-
80.0	80.0	0.0	-	-	0.0	-	-	-	0.0	-	-	-
80.0	140.0	2.7	-	-	-	-	-	-	-	-	-	-
80.0	160.0	14.6	-	-	-	-	-	-	-	-	-	-
80.0	180.0	2.7	-	-	-	-	-	-	-	-	-	-
80.0	200.0	0.0	-	-	-	-	-	-	36.5	-	-	-
83.0	39.0	0.0	-	-	-	0.0	-	-	-	1.9	-	-
83.0	65.0	0.0	-	-	0.0	-	-	-	9.9	0.0	-	-
83.0	80.0	0.0	-	-	0.0	-	-	-	0.0	5.7	-	-
87.0	60.0	0.0	-	-	0.0	-	-	-	0.0	3.0	-	-
87.0	65.0	0.0	-	-	0.0	-	-	-	32.0	0.0	-	-
87.0	70.0	0.0	-	-	0.0	-	-	-	6.1	0.0	-	-
87.0	80.0	0.0	-	-	0.0	-	-	-	6.5	-	-	-
90.0	65.0	3.2	-	-	-	-	-	-	0.0	8.3	-	-
90.0	70.0	0.0	-	-	0.0	-	-	-	-	16.1	-	-
90.0	85.0	-	-	-	0.0	-	-	-	0.0	5.2	-	-
90.0	100.0	5.5	-	-	-	-	-	-	-	16.7	-	-
90.0	120.0	2.7	-	-	-	-	-	-	5.5	-	-	-
90.0	140.0	0.0	-	-	-	-	-	-	-	3.2	-	-
90.0	160.0	5.2	-	-	-	-	-	-	-	3.1	-	-
90.0	180.0	2.7	-	-	-	-	-	-	-	6.6	-	-
93.0	200.0	0.0	-	-	-	-	-	-	-	0.0	-	-
93.0	28.0	0.0	-	-	-	-	-	-	-	2.7	-	-
93.0	30.0	0.0	-	-	-	-	-	-	-	0.0	-	-
93.0	50.0	0.0	-	-	-	-	-	-	-	0.0	-	-
93.0	60.0	0.0	-	-	-	-	-	-	-	13.3	-	-

TABLE 4. (cont.)

*Lampanyctus* spp. (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
93.0	65.0	0.0	-	-	0.0	-	0.0	-	-	-	3.0	-
93.0	80.0	0.0	-	-	0.0	-	3.2	-	-	-	11.8	-
93.0	85.0	-	-	-	0.0	-	6.3	-	-	-	6.0	-
93.0	90.0	0.0	-	-	0.0	-	0.0	-	-	-	5.7	-
97.0	35.0	3.1	-	0.0	-	-	13.1	-	-	-	0.0	-
97.0	40.0	0.0	-	0.0	-	-	6.2	-	-	-	0.0	-
97.0	70.0	0.0	-	-	0.0	-	-	-	-	-	2.9	-
97.0	75.0	-	-	-	0.0	-	0.0	-	-	-	3.2	-
97.0	80.0	0.0	-	-	-	-	3.1	-	-	-	8.7	-
97.0	90.0	0.0	-	-	-	-	-	-	-	-	-	-
100.0	35.0	0.0	-	-	0.0	-	-	-	-	-	3.0	-
100.0	40.0	0.0	-	-	0.0	-	-	-	-	-	6.2	-
100.0	45.0	0.0	-	-	0.0	-	-	-	-	-	-	-
100.0	55.0	0.0	-	-	0.0	-	-	-	-	-	3.0	-
100.0	65.0	0.0	-	-	0.0	-	-	-	-	-	0.0	-
100.0	70.0	0.0	-	-	0.0	-	-	-	-	-	3.1	-
100.0	100.0	-	0.0	-	-	-	-	-	-	-	-	-
103.0	30.0	0.0	-	-	0.0	-	0.0	-	-	-	2.6	-
103.0	45.0	0.0	-	-	0.0	-	3.4	-	-	-	0.0	-
103.0	65.0	0.0	-	-	0.0	-	6.6	-	-	-	0.0	-
105.0	140.0	-	-	2.8	-	-	-	2.8	-	-	3.3	-
107.0	65.0	-	0.0	-	0.0	-	0.0	-	-	-	3.0	-
107.0	70.0	0.0	-	-	0.0	-	0.0	-	-	-	0.0	-
107.0	80.0	-	2.9	-	0.0	-	0.0	-	-	-	5.1	-
110.0	45.0	-	0.0	-	-	-	-	-	-	-	2.7	-
110.0	120.0	0.0	-	-	-	-	-	-	-	-	11.0	-
110.0	140.0	-	-	-	-	-	-	-	-	-	14.6	-
110.0	160.0	-	-	-	-	-	-	-	-	-	-	-
113.0	29.0	-	0.0	-	-	-	1.8	-	-	-	0.0	-
113.0	45.0	-	0.0	-	-	-	3.1	-	-	-	9.0	-
113.0	50.0	-	0.0	-	-	-	0.0	-	-	-	3.0	-
113.0	55.0	-	0.0	-	-	-	8.9	-	-	-	-	-
113.0	70.0	-	0.0	-	-	-	0.0	-	-	-	-	-
113.0	80.0	-	0.0	-	-	-	18.5	-	-	-	-	-
113.0	85.0	-	-	-	-	-	9.0	-	-	-	-	-
117.0	40.0	-	0.0	-	-	-	-	-	-	-	7.8	-
117.0	45.0	-	0.0	-	-	-	2.9	-	-	-	6.6	-
117.0	50.0	-	0.0	-	-	-	3.2	-	-	-	0.0	-
117.0	65.0	-	0.0	-	-	-	5.7	-	-	-	-	-
117.0	70.0	-	0.0	-	-	-	2.8	-	-	-	0.0	-
117.0	80.0	-	0.0	-	-	-	0.0	-	-	-	0.0	-
117.0	85.0	-	-	-	-	-	12.2	-	-	-	-	-
117.0	90.0	-	-	-	-	-	20.4	-	-	-	0.0	-
120.0	30.0	-	-	-	-	-	11.3	-	-	-	0.0	-
120.0	50.0	-	-	-	-	-	2.8	-	-	-	0.0	-
120.0	55.0	-	-	-	-	-	5.9	-	-	-	0.0	-
							3.0	-	-	-	5.1	-

TABLE 4. (cont.)

*Lampanyctus* spp. (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
120.0	60.0	-	0.0	-	111.6	-	-	0.0	3.1	-	-	-
120.0	65.0	-	0.0	-	122.2	-	-	0.0	-	-	-	-
120.0	70.0	-	0.0	-	16.5	-	-	0.0	-	-	-	-
120.0	80.0	-	0.0	-	0.0	-	-	10.8	-	-	-	-
120.0	100.0	-	3.0	-	-	-	-	0.0	-	-	-	-
120.0	120.0	-	11.6	-	-	-	-	2.8	-	-	-	-
123.0	45.0	-	0.0	-	0.0	-	-	6.6	-	0.0	-	-
123.0	55.0	-	0.0	-	6.7	-	-	0.0	-	3.1	-	-
123.0	60.0	-	0.0	-	-	-	-	0.0	-	-	-	-
123.0	65.0	-	0.0	-	9.4	-	-	0.0	-	-	-	-
123.0	70.0	-	0.0	-	3.1	-	-	2.8	-	-	-	-
123.0	75.0	-	0.0	-	9.4	-	-	0.0	-	-	-	-
123.0	80.0	-	0.0	-	3.1	-	-	2.8	-	-	-	-
127.0	40.0	-	11.3	-	0.0	-	-	0.0	-	2.7	-	-
127.0	45.0	-	0.0	-	12.4	-	-	0.0	-	0.0	-	-
127.0	50.0	-	5.9	-	15.1	-	-	0.0	-	0.0	-	-
127.0	55.0	-	0.0	-	14.9	-	-	11.0	-	3.4	-	-
127.0	60.0	-	8.2	-	6.2	-	-	0.0	-	0.0	-	-
127.0	65.0	-	0.0	-	2.9	-	-	0.0	-	0.0	-	-
127.0	70.0	-	0.0	-	5.8	-	-	14.2	-	0.0	-	-
127.0	75.0	-	0.0	-	-	-	-	12.7	-	11.0	-	-
127.0	80.0	-	0.0	-	12.5	-	-	-	-	2.5	-	-
130.0	40.0	-	0.0	-	31.5	-	-	0.0	-	0.0	-	-
130.0	45.0	-	6.1	-	9.1	-	-	5.7	-	12.1	-	-
130.0	50.0	-	5.6	-	5.9	-	-	0.0	-	0.0	-	-
130.0	55.0	-	2.9	-	5.9	-	-	27.3	-	15.5	-	-
130.0	60.0	-	0.0	-	14.7	-	-	-	-	-	-	-
130.0	65.0	-	0.0	-	16.7	-	-	2.7	-	2.8	-	-
130.0	80.0	-	0.0	-	11.1	-	-	0.0	-	0.0	-	-
130.0	90.0	-	0.0	-	11.1	-	-	5.4	-	-	-	-
130.0	100.0	-	0.0	-	11.1	-	-	0.0	-	-	-	-
130.0	120.0	-	19.3	-	-	-	-	8.1	-	-	-	-
133.0	35.0	-	0.0	-	12.4	-	-	0.0	-	0.0	-	-
133.0	40.0	-	0.0	-	9.2	-	-	5.8	-	0.0	-	-
133.0	45.0	-	2.9	-	3.0	-	-	0.0	-	0.0	-	-
133.0	50.0	-	0.0	-	5.9	-	-	2.7	-	0.0	-	-
133.0	55.0	-	0.0	-	3.0	-	-	0.0	-	0.0	-	-
133.0	60.0	-	0.0	-	3.0	-	-	18.5	-	7.9	-	-
133.0	65.0	-	0.0	-	17.8	-	-	18.8	-	0.0	-	-
133.0	70.0	-	0.0	-	2.9	-	-	10.2	-	7.8	-	-
133.0	80.0	-	0.0	-	0.0	-	-	0.0	-	0.0	-	-
137.0	35.0	-	12.4	-	6.0	-	-	15.4	-	0.0	-	-
137.0	40.0	-	6.0	-	5.8	-	-	5.8	-	0.0	-	-
137.0	45.0	-	3.0	-	3.0	-	-	3.0	-	3.0	-	-
137.0	50.0	-	0.0	-	0.0	-	-	0.0	-	0.0	-	-
137.0	60.0	-	3.1	-	-	-	-	8.6	-	9.7	-	-

TABLE 4. (cont.)

*Lampanyctus spp.* (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
137.0	70.0	-	0.0	-	3.2	-	-	-	4.9	-	-	-
137.0	80.0	-	14.8	-	-	-	-	-	0.0	-	-	-
140.0	35.0	-	6.1	-	-	-	-	-	2.7	-	-	-
140.0	40.0	-	23.6	-	-	-	-	-	0.0	-	-	-
140.0	45.0	-	15.3	-	-	-	-	-	2.8	-	-	-
140.0	50.0	-	8.9	-	-	-	-	-	23.1	-	-	-

*Lampanyctus regalis*

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
60.0	60.0	0.0	-	-	-	-	-	2.8	-	-	-	-
70.0	80.0	0.0	-	-	0.0	-	-	3.2	-	-	-	-
73.0	51.0	0.0	-	-	0.0	-	-	3.2	-	-	-	-
80.0	65.0	0.0	-	-	0.0	-	-	0.0	3.0	-	-	-
87.0	65.0	0.0	-	-	0.0	-	-	6.4	-	-	-	-
87.0	80.0	0.0	-	-	0.0	-	-	6.5	-	-	0.0	-
90.0	65.0	0.0	-	-	0.0	-	-	0.0	-	-	3.0	-
93.0	55.0	0.0	-	-	0.0	-	-	6.6	-	-	0.0	-
93.0	85.0	-	-	-	-	-	-	-	-	-	3.0	-

*Lampanyctus ritteri*

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
60.0	60.0	2.7	-	-	-	-	-	0.0	-	-	-	-
60.0	70.0	0.0	-	-	-	-	-	2.8	-	-	-	-
60.0	90.0	0.0	-	-	-	-	-	2.8	-	-	-	-
60.0	100.0	6.2	-	-	-	-	-	0.0	-	-	-	-
60.0	140.0	0.0	-	-	-	-	-	15.7	-	-	-	-
63.0	51.0	0.0	-	-	0.0	-	-	4.8	-	-	-	-
63.0	55.0	0.0	-	-	0.0	-	-	2.8	-	-	-	-
63.0	60.0	6.5	-	-	0.0	-	-	14.3	-	-	-	-
67.0	47.3	0.7	-	-	0.0	-	-	-	-	-	-	-
67.0	55.0	0.0	-	-	2.7	-	-	2.9	-	-	-	-
67.0	60.0	0.0	-	-	5.5	-	-	0.0	-	-	-	-
67.0	65.0	-	-	-	2.9	-	-	0.0	-	-	-	-
67.0	70.0	-	-	-	5.7	-	-	-	-	-	-	-
67.0	80.0	-	-	-	2.7	-	-	-	-	-	-	-
70.0	53.0	0.0	-	-	2.7	-	-	-	-	-	-	-
70.0	60.0	0.0	-	-	5.7	-	-	-	-	-	-	-
70.0	65.0	-	-	-	2.7	-	-	-	-	-	-	-
70.0	70.0	-	-	-	5.8	-	-	-	-	-	-	-
70.0	80.0	8.3	-	-	0.0	-	-	0.0	-	-	-	-
70.0	90.0	2.9	-	-	-	-	-	-	-	-	-	-
70.0	100.0	2.8	-	-	-	-	-	13.8	-	-	-	-

TABLE 4. (cont.)

*Lampanyctus ritteri* (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
70.0	120.0	20.9	-	-	-	-	35.6	-	-	-	-	-
73.0	50.0	0.0	-	-	0.0	-	2.9	-	-	-	-	-
73.0	53.0	0.0	-	-	0.0	-	11.1	-	-	-	-	-
73.0	60.0	0.0	-	-	0.0	-	2.8	-	-	-	-	-
73.0	70.0	-	-	-	5.5	-	-	-	-	-	-	-
73.0	80.0	-	-	-	5.2	-	-	-	-	-	-	-
77.0	51.0	0.0	-	-	0.0	-	5.7	-	-	-	-	-
77.0	65.0	-	-	-	5.1	-	-	-	-	-	-	-
77.0	70.0	-	-	-	16.1	-	-	-	-	-	-	-
77.0	80.0	-	-	-	11.3	-	-	-	-	-	-	-
80.0	60.0	0.0	-	-	0.0	-	8.4	0.0	-	-	-	-
80.0	65.0	0.0	-	-	3.4	-	3.0	0.0	-	-	-	-
80.0	70.0	2.8	-	-	0.0	-	12.3	2.7	-	-	-	-
80.0	80.0	3.0	-	-	0.0	-	0.0	0.0	-	-	-	-
80.0	90.0	8.6	-	-	-	-	8.4	-	-	-	-	-
80.0	100.0	0.0	-	-	-	-	9.0	-	-	-	-	-
80.0	120.0	8.9	-	-	-	-	0.0	-	-	-	-	-
82.0	47.0	0.0	-	-	3.1	0.0	-	0.0	-	-	-	-
83.0	65.0	0.0	-	-	8.7	0.0	-	0.0	-	-	-	-
83.0	70.0	0.0	-	-	9.5	27.0	-	0.0	-	-	-	-
83.0	80.0	21.2	-	-	5.7	25.7	-	0.0	-	-	-	-
83.0	90.0	3.0	-	-	-	-	-	-	-	-	-	-
87.0	34.0	2.7	-	-	0.0	0.0	-	0.0	-	-	-	-
87.0	35.0	0.0	-	-	0.0	0.0	-	0.0	-	-	-	-
87.0	40.0	2.8	-	-	0.0	0.0	-	0.0	-	-	-	-
87.0	55.0	0.0	-	-	0.0	0.0	-	0.0	-	-	-	-
87.0	60.0	12.0	-	-	2.4	3.3	-	0.0	-	-	-	-
87.0	65.0	9.2	-	-	2.7	0.0	-	0.0	-	-	-	-
87.0	70.0	40.3	-	-	9.6	0.0	-	0.0	-	-	-	-
87.0	80.0	18.2	-	-	27.1	0.0	-	0.0	-	-	-	-
87.0	90.0	6.0	-	-	5.6	28.1	-	0.0	-	-	-	-
87.0	100.0	-	-	-	9.4	-	-	-	-	-	-	-
90.0	28.0	2.8	-	-	0.0	-	0.0	-	-	-	-	-
90.0	37.0	0.0	-	-	3.1	-	0.0	-	-	-	-	-
90.0	60.0	3.0	-	-	10.0	2.7	-	0.0	-	-	-	-
90.0	65.0	0.0	-	-	23.5	13.3	-	0.0	-	-	-	-
90.0	70.0	3.1	-	-	3.1	0.0	-	0.0	-	-	-	-
90.0	75.0	-	-	-	-	-	-	15.7	-	-	-	-
90.0	80.0	5.3	-	-	3.2	28.8	-	11.6	-	-	-	-
90.0	90.0	0.0	-	-	6.2	0.0	-	0.0	-	-	-	-
93.0	45.0	3.0	-	-	15.3	6.8	-	0.0	-	-	-	-
93.0	60.0	-	-	-	25.2	3.0	-	6.5	-	-	-	-
93.0	65.0	-	-	-	5.9	10.0	-	0.0	-	-	-	-
93.0	70.0	-	-	-	-	-	-	2.9	-	-	-	-
93.0	75.0	-	-	-	-	-	-	-	-	-	-	-

TABLE 4. (cont.)

*Lampanyctus ritteri* (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
93.0	80.0	3.0	-	-	3.2	-	0.0	-	-	0.0	-	-
93.0	85.0	-	-	-	16.8	-	0.0	-	-	6.0	-	-
93.0	90.0	3.2	-	-	5.7	-	-	-	-	0.0	-	-
93.0	100.0	-	-	-	6.3	-	3.3	-	-	0.0	-	-
97.0	35.0	0.0	-	-	3.2	-	0.0	-	-	6.1	-	-
97.0	40.0	0.0	-	-	0.0	-	0.0	-	-	0.0	-	-
97.0	50.0	3.2	-	-	0.0	-	0.0	-	-	0.0	-	-
97.0	55.0	0.0	-	-	0.0	-	0.0	-	-	0.0	-	-
97.0	65.0	12.3	-	-	0.0	-	0.0	-	-	0.0	-	-
97.0	80.0	6.7	-	-	-	-	9.7	-	-	-	-	-
97.0	85.0	-	-	-	17.9	-	-	-	-	-	-	-
97.0	90.0	3.0	-	-	-	-	0.0	-	-	-	-	-
97.0	95.0	-	-	-	10.9	-	-	-	-	0.0	-	-
100.0	30.0	0.0	-	-	-	-	3.2	-	-	0.0	-	-
100.0	35.0	5.8	-	-	-	-	3.2	-	-	5.5	-	-
100.0	40.0	0.0	-	-	-	-	6.4	-	-	0.0	-	-
100.0	45.0	0.0	-	-	-	-	22.7	-	-	2.9	-	-
100.0	50.0	0.0	-	-	-	-	9.0	-	-	0.0	-	-
100.0	60.0	14.9	-	-	-	-	10.1	-	-	3.1	-	-
100.0	65.0	6.0	-	-	-	-	0.0	-	-	0.0	-	-
100.0	70.0	3.1	-	-	-	-	6.7	-	-	2.3	-	-
100.0	80.0	14.6	-	-	-	-	16.7	-	-	0.0	-	-
100.0	85.0	-	-	-	-	-	17.3	-	-	0.0	-	-
100.0	90.0	12.1	-	-	-	-	46.3	-	-	0.0	-	-
100.0	100.0	-	-	-	-	-	-	-	-	0.0	-	-
103.0	35.0	17.6	-	-	-	-	3.1	-	-	0.0	-	-
103.0	40.0	29.2	-	-	-	-	15.3	-	-	0.0	-	-
103.0	45.0	14.8	-	-	-	-	18.2	-	-	0.0	-	-
103.0	50.0	5.8	-	-	-	-	6.6	-	-	0.0	-	-
103.0	55.0	25.8	-	-	-	-	6.7	-	-	3.4	-	-
103.0	60.0	24.3	-	-	-	-	3.3	-	-	0.0	-	-
103.0	65.0	14.1	-	-	-	-	9.5	-	-	0.0	-	-
103.0	70.0	18.1	-	-	-	-	9.3	-	-	0.0	-	-
103.0	75.0	-	-	-	-	-	6.4	-	-	0.0	-	-
103.0	80.0	-	-	-	-	-	6.0	-	-	0.0	-	-
103.0	85.0	-	-	-	-	-	-	-	-	-	-	-
103.0	90.0	-	-	-	-	-	-	-	-	-	-	-
107.0	31.0	-	-	-	-	-	3.0	-	-	0.0	-	-
107.0	32.0	-	-	-	-	-	4.3	-	-	0.0	-	-
107.0	40.0	-	-	-	-	-	0.0	-	-	0.0	-	-
107.0	45.0	-	-	-	-	-	5.9	-	-	5.9	-	-
107.0	50.0	-	-	-	-	-	6.0	-	-	6.0	-	-
107.0	55.0	-	-	-	-	-	0.0	-	-	0.0	-	-
107.0	60.0	-	-	-	-	-	36.4	-	-	9.7	-	-
107.0	65.0	-	-	-	-	-	5.9	-	-	3.1	-	-
107.0	70.0	-	-	-	-	-	9.5	-	-	21.4	-	-

TABLE 4. (cont.)

*Lampanyctus ritteri* (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
107.0	75.0	-	-	-	6.2	-	-	-	-	-	-	-
107.0	80.0	-	0.0	-	3.0	-	0.0	-	-	-	-	-
107.0	85.0	-	-	-	17.5	-	0.0	-	-	-	-	-
107.0	90.0	-	20.7	-	6.1	-	-	-	-	-	-	-
110.0	40.0	-	5.5	-	0.0	-	-	-	-	5.6	0.0	-
110.0	50.0	-	5.6	-	5.6	-	-	-	-	0.0	0.0	-
110.0	55.0	-	8.2	-	5.9	-	-	-	-	0.0	0.0	-
110.0	60.0	-	5.4	-	9.1	-	-	-	-	0.0	0.0	-
110.0	65.0	-	5.3	-	2.9	-	-	-	-	0.0	0.0	-
110.0	70.0	-	11.2	-	13.0	-	-	-	-	0.0	-	-
110.0	80.0	-	10.3	-	6.1	-	0.0	-	-	-	-	-
110.0	85.0	-	-	-	3.1	-	-	-	-	-	-	-
110.0	90.0	-	5.3	-	0.0	-	2.8	-	-	-	-	-
110.0	100.0	-	5.2	-	-	-	0.0	-	-	-	-	-
110.0	120.0	-	5.2	-	-	-	0.0	-	-	-	-	-
113.0	29.0	-	0.0	-	1.2	-	-	-	-	0.0	0.0	-
113.0	30.0	-	0.0	-	2.8	-	-	-	-	0.0	0.0	-
113.0	35.0	-	0.0	-	3.5	-	-	-	-	0.0	0.0	-
113.0	40.0	-	5.9	-	6.5	-	-	-	-	0.0	0.0	-
113.0	45.0	-	6.2	-	0.0	-	-	-	-	0.0	0.0	-
113.0	50.0	-	3.0	-	0.0	-	-	-	-	0.0	0.0	-
113.0	65.0	-	0.0	-	3.3	-	-	-	-	0.0	0.0	-
113.0	70.0	-	2.7	-	0.0	-	-	-	-	-	-	-
113.0	80.0	-	11.4	-	6.2	-	-	-	-	-	-	-
113.0	90.0	-	5.7	-	0.0	-	-	-	-	-	-	-
115.0	35.0	-	5.7	-	-	-	-	-	-	0.0	0.0	-
117.0	35.0	-	0.0	-	0.0	-	-	-	-	2.9	0.0	-
117.0	40.0	-	0.0	-	6.1	-	-	-	-	2.6	0.0	-
117.0	60.0	-	12.6	-	0.0	-	-	-	-	-	6.1	-
117.0	65.0	-	6.2	-	0.0	-	-	-	-	-	-	-
117.0	80.0	-	17.8	-	0.0	-	-	-	-	-	-	-
117.0	90.0	-	2.9	-	0.0	-	-	-	-	-	-	-
120.0	60.0	-	8.6	-	0.0	-	-	-	-	0.0	0.0	-
120.0	65.0	-	11.2	-	0.0	-	-	-	-	0.0	0.0	-
120.0	80.0	-	11.2	-	0.0	-	-	-	-	0.0	0.0	-
120.0	90.0	-	5.8	-	-	-	-	-	-	0.0	-	-
120.0	100.0	-	6.0	-	-	-	-	-	-	0.0	-	-
123.0	60.0	-	17.5	-	-	-	-	-	-	0.0	0.0	-
130.0	55.0	-	2.9	-	0.0	-	-	-	-	0.0	0.0	-
130.0	70.0	-	8.6	-	-	-	-	-	-	0.0	0.0	-

*Notolychnus valdiviae*

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
60.0	200.0	0.0	-	-	-	-	-	-	-	-	-	-

TABLE 4. (cont.)

*Notolychnus valdiviae* (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
70.0	120.0	3.0	-	-	-	-	0.0	-	-	-	-	-
70.0	160.0	5.9	-	-	-	-	-	-	-	-	-	-
		17.5	-	-	-	-	16.9	-	-	-	-	-
80.0	160.0	0.0	-	-	0.0	3.1	-	2.6	-	0.0	-	-
80.0	200.0	0.0	-	-	0.0	-	-	5.4	-	-	-	-
87.0	90.0	0.0	-	-	-	-	-	2.7	-	-	-	-
90.0	80.0	0.0	-	-	-	-	-	7.8	-	-	-	-
90.0	120.0	0.0	-	-	-	-	-	25.1	-	-	-	-
90.0	140.0	0.0	-	-	-	-	-	5.5	-	0.0	-	-
90.0	160.0	5.2	-	-	-	-	-	-	-	-	-	-
90.0	180.0	2.7	-	-	-	-	-	-	-	-	-	-
90.0	200.0	2.8	-	-	-	-	-	-	-	-	-	-
93.0	80.0	3.0	-	-	0.0	0.0	-	-	-	-	-	-
93.0	80.0	-	-	-	3.6	-	-	-	-	-	-	-
97.0	85.0	-	-	11.1	-	-	-	0.0	-	-	-	-
105.0	140.0	-	0.0	-	0.0	-	-	-	-	-	-	-
107.0	65.0	-	2.8	-	-	-	2.9	-	-	3.3	-	-
110.0	160.0	-	-	-	-	-	-	-	-	-	-	-

*Notoscopelus resplendens*

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
60.0	200.0	3.1	-	-	-	-	21.9	-	-	-	-	-
70.0	200.0	0.0	-	-	0.0	-	8.0	-	-	-	-	-
80.0	60.0	0.0	-	-	-	-	2.8	0.0	-	-	-	-
80.0	120.0	0.0	-	-	-	-	5.9	-	-	-	-	-
80.0	160.0	5.8	-	-	-	-	-	-	-	-	-	-
80.0	180.0	2.7	-	-	-	-	30.9	-	-	-	-	-
80.0	200.0	0.0	-	-	-	-	3.1	2.6	0.0	-	-	-
87.0	90.0	0.0	-	-	-	-	-	-	-	-	-	-
90.0	80.0	0.0	-	-	0.0	-	-	-	-	-	-	-
90.0	100.0	0.0	-	-	0.0	-	-	-	-	-	-	-
90.0	120.0	5.4	-	-	-	-	-	10.7	-	-	-	-
90.0	140.0	0.0	-	-	-	-	-	2.7	-	-	-	-
90.0	160.0	0.0	-	-	-	-	-	12.9	-	-	-	-
90.0	180.0	0.0	-	-	-	-	-	5.6	-	-	-	-
90.0	200.0	0.0	-	-	-	-	-	5.5	-	0.0	-	-
93.0	90.0	0.0	-	-	-	-	3.4	0.0	-	-	-	-
100.0	90.0	0.0	-	-	-	-	6.6	0.0	-	3.0	-	-
100.0	100.0	-	-	-	-	-	-	-	-	0.0	3.0	-
103.0	80.0	-	-	-	-	-	3.0	0.0	-	-	0.0	3.1
110.0	60.0	0.0	-	-	-	-	0.0	-	-	-	2.7	-
110.0	65.0	0.0	-	-	-	-	0.0	-	-	-	2.9	-
110.0	120.0	0.0	-	-	-	-	-	-	-	-	4.9	-
110.0	160.0	0.0	-	-	-	-	-	-	-	-	18.0	-
113.0	45.0	0.0	-	-	-	-	-	0.0	-	-	-	-
113.0	50.0	0.0	-	-	-	-	-	0.0	-	-	-	-

TABLE 4. (cont.)

*Notoscopelus resplendens* (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
117.0	60.0	-	0.0	-	0.0	-	0.0	-	-	6.1	-	-
117.0	80.0	-	0.0	-	3.1	-	0.0	-	-	2.6	0.0	-
120.0	55.0	-	0.0	-	0.0	-	-	-	-	2.8	0.0	-
120.0	60.0	-	0.0	-	0.0	-	-	-	-	2.7	-	-
120.0	65.0	-	0.0	-	0.0	-	-	-	-	0.7	-	-
120.0	120.0	-	5.8	-	-	-	-	-	-	-	-	-
123.0	150.0	-	0.0	-	0.0	-	17.0	-	-	0.0	-	-
127.0	45.0	-	0.0	-	0.0	-	13.2	-	-	0.0	-	-
127.0	70.0	-	0.0	-	0.0	-	5.7	-	-	-	-	-
127.0	80.0	-	0.0	-	-	-	2.7	-	-	-	-	-
130.0	50.0	-	0.0	-	0.0	-	-	-	-	2.7	0.0	-
130.0	55.0	-	0.0	-	0.0	-	-	-	-	2.7	0.0	-
130.0	70.0	-	0.0	-	0.0	-	-	-	-	7.5	-	-
130.0	100.0	-	0.0	-	-	-	-	-	-	2.8	-	-
130.0	120.0	-	30.4	-	-	-	-	-	-	0.0	-	-
133.0	35.0	-	0.0	-	3.1	-	-	-	-	0.0	-	-
137.0	70.0	-	0.0	-	3.2	-	-	-	-	0.0	-	-

*Stenobrachius leucopsarus*

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
60.0	49.5	0.9	-	-	-	-	-	-	-	-	-	-
60.0	51.0	8.1	-	-	-	-	-	-	-	2.0	-	-
60.0	52.0	2.3	-	-	-	-	-	-	-	0.0	-	-
60.0	55.0	18.5	-	-	-	-	-	-	-	2.8	-	-
60.0	60.0	92.8	-	-	-	-	-	-	-	11.3	-	-
60.0	70.0	48.6	-	-	-	-	-	-	-	9.2	-	-
60.0	80.0	48.3	-	-	-	-	-	-	-	0.0	-	-
60.0	90.0	33.5	-	-	-	-	-	-	-	-	-	-
63.0	49.7	1.2	-	-	-	-	-	-	-	-	-	-
63.0	50.0	4.1	-	-	-	-	-	-	-	0.0	0.0	-
63.0	51.0	44.7	-	-	-	-	-	-	-	0.0	0.0	-
63.0	52.0	226.2	-	-	-	-	-	-	-	20.5	0.0	-
63.0	55.0	361.4	-	-	-	-	-	-	-	91.1	0.0	-
63.0	60.0	52.2	-	-	-	-	-	-	-	55.4	0.0	-
63.0	70.0	-	-	-	-	-	-	-	-	11.4	-	-
63.0	80.0	-	-	-	-	-	-	-	-	44.0	-	-
67.0	47.3	3.4	-	-	-	-	-	-	-	0.0	-	-
67.0	48.0	1.2	-	-	-	-	-	-	-	29.0	-	-
67.0	49.0	19.4	-	-	-	-	-	-	-	17.0	0.0	-
67.0	50.0	28.2	-	-	-	-	-	-	-	13.5	2.8	-
67.0	55.0	29.9	-	-	-	-	-	-	-	13.5	0.0	-
67.0	60.0	111.4	-	-	-	-	-	-	-	412.5	3.0	-
67.0	65.0	-	-	-	-	-	-	-	-	206.6	-	-
67.0	70.0	-	-	-	-	-	-	-	-	215.8	-	-

TABLE 4. (cont.)

*Stenobrachius leucopsarus* (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
67.0	80.0	-	-	-	47.7	-	-	-	-	-	-	-
70.0	50.0	96.7	-	-	29.0	-	-	-	-	-	-	-
70.0	51.0	64.5	-	-	223.6	-	-	-	-	-	-	-
70.0	52.0	66.2	-	-	237.4	-	-	-	-	-	-	-
70.0	53.0	112.1	-	-	145.8	-	-	-	-	-	-	-
70.0	55.0	252.3	-	-	-	-	-	-	-	-	-	-
70.0	60.0	42.9	-	-	77.2	-	-	-	-	-	-	-
70.0	65.0	-	-	-	75.3	-	-	-	-	-	-	-
70.0	70.0	292.7	-	-	14.5	-	-	-	-	-	-	-
70.0	80.0	0.0	-	-	11.0	-	-	-	-	-	-	-
70.0	100.0	2.8	-	-	-	-	-	-	-	-	-	-
70.0	120.0	3.0	-	-	-	-	-	-	-	-	-	-
73.0	49.4	0.9	-	-	10.4	-	-	-	-	-	-	-
73.0	50.0	143.8	-	-	8.3	-	-	-	-	-	-	-
73.0	51.0	358.4	-	-	72.5	-	-	-	-	-	-	-
73.0	53.0	158.3	-	-	84.3	-	-	-	-	-	-	-
73.0	60.0	195.3	-	-	59.2	-	-	-	-	-	-	-
73.0	65.0	-	-	-	5.8	-	-	-	-	-	-	-
73.0	70.0	-	-	-	2.8	-	-	-	-	-	-	-
73.0	80.0	-	-	-	20.7	-	-	-	-	-	-	-
77.0	47.6	1.7	-	-	0.0	-	-	-	-	-	-	-
77.0	48.0	0.6	-	-	0.0	-	-	-	-	-	-	-
77.0	51.0	83.2	-	-	15.1	-	-	-	-	-	-	-
77.0	55.0	25.9	-	-	21.7	-	-	-	-	-	-	-
77.0	57.0	86.7	-	-	98.0	-	-	-	-	-	-	-
77.0	60.0	-	-	-	15.7	-	-	-	-	-	-	-
77.0	65.0	-	-	-	155.6	-	-	-	-	-	-	-
77.0	70.0	-	-	-	9.2	-	-	-	-	-	-	-
77.0	80.0	-	-	-	5.7	-	-	-	-	-	-	-
80.0	50.2	1.1	-	-	1.5	-	-	-	-	-	-	-
80.0	51.0	37.9	-	-	3.5	-	-	-	-	-	-	-
80.0	52.0	194.1	-	-	59.9	-	-	-	-	-	-	-
80.0	55.0	193.3	-	-	179.2	-	-	-	-	-	-	-
80.0	60.0	74.5	-	-	132.2	-	-	-	-	-	-	-
80.0	65.0	68.6	-	-	111.2	-	-	-	-	-	-	-
80.0	70.0	25.6	-	-	44.2	-	-	-	-	-	-	-
80.0	80.0	17.9	-	-	29.0	-	-	-	-	-	-	-
80.0	90.0	5.8	-	-	-	-	-	-	-	-	-	-
80.0	120.0	185.9	-	-	-	-	-	-	-	-	-	-
82.0	47.0	11.4	-	-	-	-	-	-	-	-	-	-
83.0	40.0	9.6	-	-	-	-	-	-	-	-	-	-
83.0	43.0	3.0	-	-	-	-	-	-	-	-	-	-
83.0	51.0	9.4	-	-	-	-	-	-	-	-	-	-
83.0	55.0	290.0	-	-	-	-	-	-	-	-	-	-
83.0	60.0	69.9	-	-	-	-	-	-	-	-	-	-
83.0	65.0	71.8	-	-	-	-	-	-	-	-	-	-

TABLE 4. (cont.)

*Stenobrachius leucopsarus* (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
83.0	70.0	38.3	-	-	31.5	-	13.5	-	-	0.0	-	-
83.0	80.0	141.2	-	-	48.8	-	4.1	-	-	0.0	-	-
83.0	90.0	39.4	-	-	0.0	0.0	0.0	-	-	0.0	-	-
87.0	32.5	30.0	-	-	5.3	0.0	0.0	-	-	0.0	-	-
87.0	33.0	131.3	-	-	22.1	2.6	6.1	-	-	0.0	-	-
87.0	34.0	419.2	-	-	27.3	-	-	-	-	0.0	-	-
87.0	35.0	449.8	-	-	107.6	0.0	0.0	-	-	0.0	-	-
87.0	40.0	119.7	-	-	13.7	0.0	0.0	-	-	0.0	-	-
87.0	45.0	58.1	-	-	4.6	0.0	0.0	-	-	0.0	-	-
87.0	50.0	59.1	-	-	5.6	52.3	-	-	-	0.0	-	-
87.0	55.0	226.5	-	-	45.4	0.0	0.0	-	-	0.0	-	-
87.0	60.0	60.0	-	-	24.1	3.2	-	-	-	0.0	-	-
87.0	65.0	9.2	-	-	9.6	0.0	0.0	-	-	0.0	-	-
87.0	70.0	52.7	-	-	9.0	0.0	0.0	-	-	0.0	-	-
87.0	80.0	90.9	-	-	0.0	-	-	-	-	0.0	-	-
87.0	90.0	35.9	-	-	9.4	-	-	-	-	0.0	-	-
87.0	100.0	-	-	-	5.1	4.5	0.0	-	-	0.0	-	-
90.0	27.5	-	-	-	44.6	3.1	0.0	-	-	0.0	-	-
90.0	28.0	-	-	-	72.9	18.3	0.0	-	-	0.0	-	-
90.0	32.0	-	-	-	116.6	19.3	34.4	26.6	2.1	0.0	-	-
90.0	37.0	-	-	-	27.9	34.4	0.0	-	-	0.0	-	-
90.0	45.0	-	-	-	190.6	19.3	3.3	3.3	0.0	0.0	-	-
90.0	53.0	-	-	-	60.0	26.6	0.0	-	-	0.0	-	-
90.0	60.0	-	-	-	65.0	19.4	2.9	0.0	-	0.0	-	-
90.0	65.0	-	-	-	70.0	6.2	0.0	-	-	0.0	-	-
90.0	80.0	-	-	-	80.0	47.9	0.0	-	-	0.0	-	-
90.0	90.0	-	-	-	100.0	9.0	9.6	18.6	-	0.0	-	-
93.0	27.0	-	-	-	100.0	2.8	0.0	-	-	0.0	-	-
93.0	28.0	-	-	-	93.0	0.0	0.0	-	-	0.0	-	-
93.0	30.0	-	-	-	69.9	26.6	2.9	2.9	-	0.0	-	-
93.0	35.0	-	-	-	78.0	17.2	-	-	-	0.0	-	-
93.0	40.0	-	-	-	76.0	14.6	3.3	24.3	-	0.0	-	-
93.0	45.0	-	-	-	15.5	-	0.0	0.0	-	6.6	-	-
93.0	50.0	-	-	-	12.3	-	4.2	0.0	-	0.0	-	-
93.0	55.0	-	-	-	15.3	-	13.5	13.5	-	0.0	-	-
93.0	60.0	-	-	-	12.6	-	8.9	8.9	-	0.0	-	-
93.0	65.0	-	-	-	70.0	2.9	20.0	20.0	-	3.3	-	-
93.0	80.0	-	-	-	57.2	-	3.2	3.2	-	0.0	-	-
93.0	90.0	-	-	-	28.7	-	6.7	6.7	-	0.0	-	-
96.8	28.7	-	-	-	7.1	-	0.0	0.0	-	0.0	-	-
97.0	29.0	-	-	-	37.4	-	0.0	0.0	-	0.0	-	-
97.0	30.0	-	-	-	54.0	-	12.6	12.6	-	0.0	-	-
97.0	32.0	-	-	-	25.0	-	34.1	34.1	-	0.0	-	-

TABLE 4. (cont.)

*Stenobrachius leucopsarus* (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
97.0	35.0	3.1	-	15.8	-	-	3.3	-	-	0.0	-	-
97.0	40.0	3.2	-	0.0	-	-	0.0	-	-	0.0	-	-
97.0	45.0	6.1	-	0.0	-	-	0.0	-	-	0.0	-	-
97.0	50.0	37.8	-	0.0	-	-	0.0	-	-	0.0	-	-
97.0	55.0	15.0	-	-	-	-	0.0	-	-	0.0	-	-
97.0	60.0	14.8	-	-	-	-	0.0	-	-	0.0	-	-
97.0	65.0	58.3	-	17.3	-	-	0.0	-	-	0.0	-	-
97.0	70.0	17.7	-	-	21.8	-	-	0.0	-	0.0	-	-
97.0	75.0	-	-	-	-	-	-	-	-	0.0	-	-
97.0	80.0	0.0	-	-	-	-	-	-	-	0.0	-	-
97.0	90.0	15.3	-	-	-	-	0.0	-	-	0.0	-	-
100.0	29.0	29.5	-	0.0	-	-	0.0	-	-	0.0	-	-
100.0	30.0	56.2	-	0.0	-	-	0.0	-	-	0.0	-	-
100.0	35.0	0.0	-	-	-	3.2	-	-	0.0	-	-	-
100.0	45.0	3.1	-	0.0	-	-	0.0	-	-	0.0	-	-
100.0	50.0	19.5	-	-	-	6.0	-	-	0.0	-	-	-
100.0	55.0	3.0	-	-	-	0.0	-	-	0.0	-	-	-
100.0	60.0	20.9	-	-	-	0.0	-	-	0.0	-	-	-
100.0	65.0	20.9	-	-	-	12.8	-	-	0.0	-	-	-
100.0	70.0	24.6	-	-	-	0.0	-	-	0.0	-	-	-
100.0	85.0	-	-	-	-	3.5	-	-	-	-	-	-
100.0	100.0	-	-	-	17.7	-	-	0.0	-	-	-	-
103.0	28.5	0.0	-	-	-	-	1.8	-	-	0.0	-	-
103.0	29.0	0.0	-	-	-	-	0.0	-	-	5.0	-	-
103.0	30.0	2.5	-	-	-	-	15.3	-	-	2.6	-	-
103.0	35.0	14.7	-	-	-	-	15.5	-	-	0.0	-	-
103.0	40.0	5.8	-	-	-	-	0.0	-	-	0.0	-	-
103.0	50.0	2.9	-	-	-	-	0.0	-	-	0.0	-	-
103.0	55.0	2.9	-	-	-	-	0.0	-	-	0.0	-	-
103.0	60.0	3.0	-	-	-	-	0.0	-	-	0.0	-	-
103.0	65.0	42.5	-	-	-	-	3.2	-	-	0.0	-	-
103.0	70.0	15.1	-	-	-	-	0.0	-	-	0.0	-	-
103.0	80.0	-	-	6.0	-	-	0.0	-	-	0.0	-	-
107.0	31.0	-	-	2.2	-	-	0.0	-	-	0.0	-	-
107.0	32.0	-	-	0.0	-	-	6.1	-	-	0.0	-	-
107.0	45.0	-	3.0	-	-	-	0.0	-	-	0.0	-	-
107.0	50.0	-	2.9	-	-	-	0.0	-	-	0.0	-	-
107.0	60.0	-	18.2	-	-	-	0.0	-	-	0.0	-	-
110.0	70.0	-	2.8	-	-	-	0.0	-	-	0.0	-	-
110.0	140.0	-	2.7	-	-	-	0.0	-	-	0.0	-	-

*Triphoturus mexicanus*

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
70.0	120.0	0.0	-	-	-	-	-	-	-	-	-	-

TABLE 4. (cont.)

*Triphoturus mexicanus* (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
80.0	60.0	0.0	-	-	0.0	-	-	2.8	2.7	-	-	-
80.0	65.0	0.0	-	-	0.0	-	-	3.0	0.7	-	-	-
80.0	70.0	0.0	-	-	0.0	-	-	0.0	2.7	-	-	-
80.0	80.0	0.0	-	-	0.0	-	-	0.0	2.5	-	-	-
82.0	47.0	0.0	-	-	0.0	-	-	-	-	-	-	-
83.0	39.0	0.0	-	-	-	-	-	-	-	-	-	-
83.0	40.0	0.0	-	-	0.0	-	-	-	-	-	-	-
83.0	43.0	0.0	-	-	0.0	-	-	-	-	-	-	-
83.0	60.0	0.0	-	-	0.0	-	-	-	-	-	-	-
83.0	75.0	-	-	-	-	-	-	-	-	-	-	-
83.0	90.0	0.0	-	-	-	-	-	-	-	-	-	-
87.0	32.5	0.0	-	-	0.0	-	-	6.4	-	-	-	-
87.0	33.0	0.0	-	-	0.0	-	-	0.0	-	-	-	-
87.0	34.0	0.0	-	-	0.0	-	-	0.0	-	-	-	-
87.0	35.0	0.0	-	-	0.0	-	-	0.0	-	-	-	-
87.0	40.0	0.0	-	-	0.0	-	-	0.0	-	-	-	-
87.0	45.0	0.0	-	-	0.0	-	-	0.0	-	-	-	-
87.0	50.0	0.0	-	-	0.0	-	-	0.0	-	-	-	-
87.0	55.0	0.0	-	-	0.0	-	-	0.0	-	-	-	-
87.0	60.0	0.0	-	-	0.0	-	-	0.0	-	-	-	-
87.0	65.0	0.0	-	-	0.0	-	-	0.0	-	-	-	-
87.0	70.0	0.0	-	-	0.0	-	-	3.2	3.0	0.0	0.0	-
87.0	80.0	0.0	-	-	0.0	-	-	21.1	9.8	0.0	0.0	-
87.0	90.0	0.0	-	-	0.0	-	-	18.7	-	-	-	-
90.0	32.0	0.0	-	-	0.0	-	-	-	7.7	-	-	-
90.0	37.0	0.0	-	-	0.0	-	-	-	35.5	-	-	-
90.0	45.0	0.0	-	-	0.0	-	-	-	7.7	-	-	-
90.0	53.0	0.0	-	-	0.0	-	-	-	0.0	-	-	-
90.0	60.0	0.0	-	-	0.0	-	-	-	5.4	-	-	-
90.0	65.0	0.0	-	-	0.0	-	-	-	2.7	-	-	-
90.0	70.0	0.0	-	-	0.0	-	-	-	8.3	-	-	-
90.0	80.0	0.0	-	-	0.0	-	-	-	18.3	-	-	-
90.0	85.0	-	-	-	-	-	-	-	-	-	-	-
90.0	90.0	0.0	-	-	0.0	-	-	-	-	-	-	-
90.0	100.0	0.0	-	-	0.0	-	-	-	-	-	-	-
90.0	140.0	0.0	-	-	0.0	-	-	-	-	-	-	-
93.0	26.5	0.0	-	-	-	-	-	-	2.5	-	-	-
93.0	27.0	0.0	-	-	-	-	-	-	17.2	-	-	-
93.0	28.0	0.0	-	-	-	-	-	-	38.5	-	-	-
93.0	30.0	0.0	-	-	-	-	-	-	49.1	-	-	-
93.0	35.0	0.0	-	-	-	-	-	-	12.2	-	-	-
93.0	40.0	0.0	-	-	-	-	-	-	3.3	-	-	-
93.0	45.0	0.0	-	-	-	-	-	-	18.9	-	-	-
93.0	50.0	0.0	-	-	-	-	-	-	0.0	-	-	-
93.0	55.0	0.0	-	-	-	-	-	-	19.8	-	-	-
93.0	60.0	0.0	-	-	-	-	-	-	6.6	-	-	-
					-	-	-	-	3.4	-	-	-

TABLE 4. (cont.)

*Triphoturus mexicanus* (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
93.0	65.0	0.0	-	-	3.0	-	9.8	-	-	8.9	-	-
93.0	70.0	0.0	-	-	6.7	-	3.3	-	-	0.0	-	-
93.0	75.0	-	-	-	-	9.5	6.3	-	-	8.6	-	-
93.0	80.0	0.0	-	-	-	-	-	-	-	52.9	-	-
93.0	85.0	-	-	-	0.0	-	18.8	-	-	47.8	-	-
93.0	90.0	0.0	-	-	0.0	-	0.0	-	-	22.8	-	-
96.8	28.7	0.0	-	-	0.0	-	0.0	-	-	1.5	-	-
97.0	30.0	0.0	-	-	0.0	-	2.8	-	-	4.5	-	-
97.0	32.0	0.0	-	-	0.0	-	10.1	-	-	12.3	-	-
97.0	35.0	0.0	-	-	6.3	-	26.0	-	-	26.2	-	-
97.0	40.0	0.0	-	-	9.6	-	26.2	-	-	42.4	-	-
97.0	45.0	0.0	-	-	6.9	-	9.5	-	-	69.9	-	-
97.0	50.0	0.0	-	-	6.4	-	33.2	-	-	50.3	-	-
97.0	55.0	0.0	-	-	-	-	15.1	-	-	0.0	-	-
97.0	60.0	0.0	-	-	-	-	31.4	-	-	2.8	-	-
97.0	65.0	0.0	-	-	0.0	-	73.8	-	-	2.8	-	-
97.0	70.0	0.0	-	-	-	-	215.6	-	-	20.2	-	-
97.0	75.0	-	-	-	0.0	-	-	-	-	6.5	-	-
97.0	80.0	0.0	-	-	-	-	54.7	-	-	63.8	-	-
97.0	85.0	-	-	-	3.6	-	9.3	-	-	-	-	-
97.0	90.0	0.0	-	-	-	-	0.0	-	-	24.8	-	-
100.0	30.0	0.0	-	-	0.0	-	-	-	-	149.0	-	-
100.0	35.0	0.0	-	-	0.0	-	-	-	-	135.2	-	-
100.0	40.0	0.0	-	-	22.5	-	-	-	-	129.7	-	-
100.0	45.0	0.0	-	-	22.7	-	-	-	-	111.8	-	-
100.0	50.0	0.0	-	-	18.1	-	-	-	-	14.7	-	-
100.0	55.0	0.0	-	-	6.0	-	-	-	-	5.8	-	-
100.0	60.0	0.0	-	-	10.1	-	-	-	-	9.2	-	-
100.0	65.0	0.0	-	-	3.2	-	-	-	-	15.3	-	-
100.0	70.0	0.0	-	-	0.0	-	-	-	-	18.7	-	-
100.0	80.0	0.0	-	-	10.0	-	-	-	-	0.0	-	-
100.0	85.0	-	-	-	10.4	-	-	-	-	12.0	-	-
100.0	100.0	-	-	-	0.0	-	-	-	-	3.0	-	-
100.0	120.0	-	-	-	0.0	-	-	-	-	-	-	-
103.0	28.5	0.0	-	-	-	-	0.0	-	-	7.0	-	-
103.0	29.0	0.0	-	-	0.0	-	-	-	-	29.1	-	-
103.0	30.0	0.0	-	-	3.1	-	-	-	-	10.4	-	-
103.0	35.0	0.0	-	-	3.1	-	-	-	-	25.6	-	-
103.0	40.0	0.0	-	-	24.5	-	108.1	-	-	278.6	-	-
103.0	45.0	0.0	-	-	39.4	-	69.9	-	-	145.3	-	-
103.0	50.0	0.0	-	-	16.5	-	80.7	-	-	12.6	-	-
103.0	55.0	0.0	-	-	30.0	-	23.5	-	-	9.3	-	-
103.0	60.0	0.0	-	-	10.1	-	0.0	-	-	111.2	-	-
103.0	65.0	0.0	-	-	9.5	-	52.8	-	-	159.5	-	-
103.0	70.0	-	-	-	6.2	-	43.7	-	-	81.2	-	-
							6.4	-	-	-	-	-

TABLE 4. (cont.)

*Triphoturus mexicanus* (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
103.0	80.0	-	0.0	-	8.9	-	18.1	-	-	-	-	-
103.0	85.0	-	-	-	8.7	-	-	-	-	-	-	-
103.0	90.0	-	0.0	-	6.3	-	16.6	-	-	-	-	-
105.0	140.0	-	0.0	-	0.0	-	0.0	-	5.6	-	-	-
107.0	30.0	-	0.0	-	5.7	-	0.0	-	-	-	-	-
107.0	31.0	-	0.0	-	12.2	-	13.8	-	-	-	-	-
107.0	32.0	-	0.0	-	0.0	-	110.7	-	-	-	-	-
107.0	35.0	-	0.0	-	0.0	-	12.0	-	113.3	-	-	-
107.0	40.0	-	0.0	-	0.0	-	11.9	-	20.7	-	-	-
107.0	45.0	-	0.0	-	0.0	-	21.1	-	50.0	-	-	-
107.0	50.0	-	0.0	-	0.0	-	25.4	-	26.5	-	-	-
107.0	55.0	-	0.0	-	0.0	-	48.6	-	33.7	-	-	-
107.0	60.0	-	0.0	-	0.0	-	9.4	-	6.3	-	-	-
107.0	65.0	-	0.0	-	0.0	-	15.3	-	61.4	-	-	-
107.0	70.0	-	0.0	-	0.0	-	24.6	-	-	-	-	-
107.0	75.0	-	0.0	-	0.0	-	32.7	-	138.5	-	-	-
107.0	80.0	-	0.0	-	0.0	-	26.2	-	23.4	-	-	-
107.0	85.0	-	0.0	-	0.0	-	9.2	-	-	-	-	-
107.0	90.0	-	0.0	-	0.0	-	0.0	-	-	-	-	-
110.0	32.0	-	0.0	-	0.0	-	3.1	-	-	-	-	-
110.0	35.0	-	0.0	-	0.0	-	11.4	-	-	-	-	-
110.0	40.0	-	0.0	-	0.0	-	0.0	-	-	-	-	-
110.0	45.0	-	0.0	-	0.0	-	2.8	-	-	-	-	-
110.0	50.0	-	0.0	-	0.0	-	17.7	-	-	-	-	-
110.0	55.0	-	0.0	-	0.0	-	58.0	-	-	-	-	-
110.0	60.0	-	0.0	-	0.0	-	17.0	-	-	-	-	-
110.0	65.0	-	0.0	-	0.0	-	2.7	-	-	-	-	-
110.0	70.0	-	0.0	-	0.0	-	35.2	-	-	-	-	-
110.0	75.0	-	0.0	-	0.0	-	35.6	-	-	-	-	-
110.0	80.0	-	0.0	-	0.0	-	25.6	-	-	-	-	-
110.0	85.0	-	0.0	-	0.0	-	3.0	-	-	-	-	-
110.0	90.0	-	0.0	-	0.0	-	18.4	-	-	-	-	-
110.0	95.0	-	0.0	-	0.0	-	49.4	-	-	-	-	-
110.0	100.0	-	0.0	-	0.0	-	-	-	-	-	-	-
110.0	105.0	-	0.0	-	0.0	-	-	-	-	-	-	-
110.0	110.0	-	0.0	-	0.0	-	-	-	-	-	-	-
110.0	115.0	-	0.0	-	0.0	-	-	-	-	-	-	-
110.0	120.0	-	0.0	-	0.0	-	-	-	-	-	-	-
110.0	125.0	-	0.0	-	0.0	-	-	-	-	-	-	-
113.0	29.0	-	0.0	-	0.0	-	-	-	-	-	-	-
113.0	30.0	-	0.0	-	0.0	-	-	-	-	-	-	-
113.0	35.0	-	0.0	-	0.0	-	-	-	-	-	-	-
113.0	40.0	-	0.0	-	0.0	-	-	-	-	-	-	-
113.0	45.0	-	0.0	-	0.0	-	-	-	-	-	-	-
113.0	50.0	-	0.0	-	0.0	-	-	-	-	-	-	-
113.0	55.0	-	0.0	-	0.0	-	-	-	-	-	-	-
113.0	60.0	-	0.0	-	0.0	-	-	-	-	-	-	-
113.0	65.0	-	0.0	-	0.0	-	-	-	-	-	-	-
113.0	70.0	-	0.0	-	0.0	-	-	-	-	-	-	-
113.0	75.0	-	0.0	-	0.0	-	-	-	-	-	-	-
113.0	80.0	-	0.0	-	0.0	-	-	-	-	-	-	-

TABLE 4. (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.	<i>Triphoturus mexicanus</i> (cont.)												
													1	2	3	4	5	6	7	8	9	10	11	12	13
1113.0	85.0	-	-	-	65.8	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1113.0	90.0	-	0.0	0.0	17.0	-	3.3	-	-	20.9	28.2	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1115.0	35.0	-	0.0	0.0	0.0	0.0	-	0.0	-	-	2.3	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1117.0	24.8	-	0.0	0.0	0.0	0.0	-	0.0	-	-	13.2	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1117.0	26.0	-	0.0	0.0	0.0	0.0	-	0.0	-	-	2.6	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1117.0	30.0	-	0.0	0.0	3.1	0.0	-	28.6	-	-	29.2	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1117.0	35.0	-	0.0	0.0	85.7	-	-	98.4	-	-	40.4	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1117.0	40.0	-	0.0	0.0	101.9	-	-	3.4	-	-	26.0	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1117.0	45.0	-	0.0	0.0	32.0	-	-	49.8	-	-	106.4	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1117.0	50.0	-	0.0	0.0	26.2	-	-	207.9	-	-	107.7	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1117.0	55.0	-	0.0	0.0	61.5	-	23.1	-	-	78.7	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1117.0	60.0	-	0.0	0.0	31.1	56.6	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1117.0	65.0	-	0.0	0.0	2.8	40.3	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1117.0	70.0	-	0.0	0.0	18.2	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1117.0	75.0	-	0.0	0.0	134.6	29.7	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1117.0	80.0	-	0.0	0.0	101.9	3.3	-	221.2	-	4.3	0.0	-	-	-	-	-	-	-	-	-	-	-	-	-	
1117.0	85.0	-	0.0	0.0	183.3	9.3	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1117.0	90.0	-	2.9	2.9	6.2	9.2	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1118.0	39.0	-	0.0	0.0	0.0	0.0	0.0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1119.0	33.0	-	0.0	0.0	0.0	0.0	0.0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
120.0	25.0	-	0.0	0.0	0.0	0.0	0.0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
120.0	30.0	-	0.0	0.0	0.0	0.0	5.7	-	-	2.7	4.7	-	-	-	-	-	-	-	-	-	-	-	-	-	-
120.0	40.0	-	0.0	0.0	0.0	0.0	0.0	-	-	0.0	0.0	-	-	-	-	-	-	-	-	-	-	-	-	-	-
120.0	45.0	-	0.0	0.0	0.0	0.0	18.5	-	-	68.6	29.8	-	-	-	-	-	-	-	-	-	-	-	-	-	
120.0	50.0	-	0.0	0.0	0.0	0.0	29.5	-	-	75.3	38.5	-	-	-	-	-	-	-	-	-	-	-	-	-	
120.0	55.0	-	0.0	0.0	0.0	14.9	84.4	-	-	158.7	127.7	-	-	-	-	-	-	-	-	-	-	-	-	-	
120.0	60.0	-	0.0	0.0	28.8	116.3	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
120.0	65.0	-	0.0	0.0	14.6	125.4	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
120.0	70.0	-	0.0	0.0	21.6	30.9	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
120.0	75.0	-	0.0	0.0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
120.0	80.0	-	0.0	0.0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
120.0	100.0	-	0.0	0.0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
120.0	120.0	-	0.0	0.0	2.9	0.0	0.0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
123.0	36.0	-	0.0	0.0	0.0	0.0	0.0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
123.0	37.0	-	0.0	0.0	2.9	8.6	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
123.0	42.0	-	0.0	0.0	17.8	8.7	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
123.0	45.0	-	0.0	0.0	5.8	9.3	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
123.0	50.0	-	0.0	0.0	2.9	0.0	0.0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
123.0	55.0	-	0.0	0.0	3.0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
123.0	60.0	-	0.0	0.0	0.0	0.0	0.0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
123.0	65.0	-	0.0	0.0	0.0	0.0	0.0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
123.0	70.0	-	0.0	0.0	0.0	0.0	0.0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
123.0	75.0	-	0.0	0.0	0.0	0.0	0.0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
123.0	80.0	-	0.0	0.0	0.0	0.0	0.0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
123.0	85.0	-	0.0	0.0	0.0	0.0	0.0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
123.0	90.0	-	0.0	0.0	0.0	0.0	0.0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
127.0	33.0	-	0.0	0.0	0.0	0.0	0.0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
127.0	34.0	-	0.0	0.0	0.0	0.0	0.0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

TABLE 4. (cont.)

*Triphoturus mexicanus* (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
127.0	40.0	-	36.8	-	5.9	-	-	0.0	46.4	12.6	-	-
127.0	45.0	-	6.0	-	71.5	-	-	74.3	-	18.6	-	-
127.0	50.0	-	17.6	-	54.5	-	-	81.5	-	15.1	-	-
127.0	55.0	-	0.0	-	20.8	-	-	21.9	-	3.2	-	-
127.0	60.0	-	16.4	-	3.1	-	-	73.3	-	6.7	-	-
127.0	65.0	-	8.7	-	8.8	-	-	25.9	-	-	-	-
127.0	70.0	-	5.9	-	8.6	-	-	25.6	-	-	-	-
127.0	80.0	-	5.8	-	-	-	-	13.7	-	-	-	-
130.0	28.0	-	0.0	-	2.7	-	-	-	0.0	0.0	-	-
130.0	35.0	-	3.3	-	408.7	-	-	38.3	0.0	-	-	-
130.0	40.0	-	27.5	-	239.4	-	-	28.8	75.0	-	-	-
130.0	45.0	-	8.4	-	6.1	-	-	36.8	21.2	-	-	-
130.0	50.0	-	20.2	-	8.9	-	-	54.6	16.6	-	-	-
130.0	55.0	-	11.5	-	14.7	-	-	368.5	65.1	-	-	-
130.0	60.0	-	41.8	-	0.0	-	-	441.2	0.0	-	-	-
130.0	65.0	-	-	-	6.2	-	-	-	-	-	-	-
130.0	70.0	-	11.5	-	0.0	-	-	34.9	-	-	-	-
130.0	80.0	-	11.1	-	-	-	-	5.5	-	-	-	-
130.0	90.0	-	8.0	-	-	-	-	24.4	-	-	-	-
130.0	100.0	-	0.0	-	-	-	-	5.7	-	-	-	-
130.0	120.0	-	2.8	-	-	-	-	10.8	-	0.0	-	-
133.0	25.0	-	0.0	-	21.3	-	-	-	-	-	-	-
133.0	30.0	-	0.0	-	15.5	-	-	0.0	-	0.0	-	-
133.0	35.0	-	8.6	-	18.5	-	-	3.0	-	0.0	-	-
133.0	40.0	-	35.8	-	15.4	-	-	0.0	-	9.1	-	-
133.0	45.0	-	32.2	-	12.1	-	-	180.4	-	0.0	-	-
133.0	50.0	-	17.0	-	20.7	-	-	31.9	-	0.0	-	-
133.0	55.0	-	13.7	-	8.9	-	-	15.7	-	3.0	-	-
133.0	60.0	-	17.3	-	15.1	-	-	-	-	-	-	-
133.0	65.0	-	8.6	-	37.1	-	-	-	-	-	-	-
133.0	70.0	-	2.8	-	50.5	-	-	-	-	-	-	-
133.0	80.0	-	5.8	-	-	-	-	56.1	-	-	-	-
137.0	30.0	-	27.7	-	71.5	-	-	-	0.0	0.0	-	-
137.0	35.0	-	24.9	-	30.1	-	-	-	2.7	0.0	-	-
137.0	40.0	-	55.4	-	6.0	-	-	-	0.0	3.0	-	-
137.0	45.0	-	14.5	-	33.0	-	-	-	6.5	3.0	-	-
137.0	50.0	-	0.0	-	16.3	-	-	-	52.0	0.0	-	-
137.0	55.0	-	0.0	-	12.8	-	-	-	27.6	-	-	-
137.0	60.0	-	-	-	3.1	-	-	-	19.4	-	-	-
137.0	65.0	-	-	-	16.1	-	-	-	-	-	-	-
137.0	70.0	-	0.0	-	0.0	-	-	-	4.9	-	-	-
137.0	80.0	-	-	-	-	-	-	-	-	19.9	-	-
140.0	35.0	-	-	-	-	-	-	-	2.7	0.0	-	-
140.0	40.0	-	-	-	-	-	-	-	0.0	0.0	-	-
140.0	45.0	-	-	-	-	-	-	-	16.6	-	-	-
140.0	50.0	-	-	-	-	-	-	-	-	23.1	-	-

TABLE 4. (cont.)

*Triphoturus nigrescens*

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
80.0	200.0	0.0	-	-	-	-	-	2.8	-	-	-	-
80.0	200.0	0.0	-	-	-	-	-	5.6	-	-	-	-
90.0	200.0	0.0	-	-	-	-	-	2.8	-	-	-	-
<i>Centrobranchus</i> spp.												

*Diogenichthys* spp.

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
93.0	90.0	0.0	-	-	0.0	-	-	3.1	-	-	0.0	-
100.0	55.0	0.0	-	-	0.0	-	-	0.0	-	-	3.0	-
103.0	30.0	0.0	-	-	0.0	-	-	0.0	-	-	2.6	-
107.0	31.0	-	0.0	-	0.0	-	-	0.0	-	-	4.8	-
107.0	50.0	-	0.0	-	0.0	-	-	0.0	-	-	3.2	-
107.0	70.0	-	3.2	-	0.0	-	-	0.0	-	-	6.1	-
110.0	45.0	-	0.0	-	0.0	-	-	-	-	2.6	0.0	-
110.0	120.0	-	0.0	-	-	-	-	-	-	-	-	-
110.0	140.0	-	0.0	-	-	-	-	-	-	-	-	-
130.0	45.0	-	0.0	-	0.0	-	-	-	-	-	-	-

*Diogenichthys atlanticus*

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
60.0	100.0	6.2	-	-	-	-	-	-	-	-	-	-
60.0	140.0	0.0	-	-	-	-	-	-	-	-	-	-
60.0	160.0	6.5	-	-	-	-	-	-	-	-	-	-
60.0	200.0	0.0	-	-	-	-	-	-	-	-	-	-
70.0	70.0	0.0	-	-	-	-	-	2.9	-	-	-	-
70.0	80.0	8.3	-	-	-	-	-	0.0	-	-	-	-
70.0	90.0	2.9	-	-	-	-	-	-	-	-	-	-
70.0	100.0	2.8	-	-	-	-	-	-	-	-	-	-
70.0	120.0	14.9	-	-	-	-	-	-	-	-	-	-
70.0	160.0	8.9	-	-	-	-	-	-	-	-	-	-
70.0	180.0	3.0	-	-	-	-	-	-	-	-	-	-
70.0	200.0	6.1	-	-	-	-	-	-	-	-	-	-
80.0	65.0	3.1	-	-	-	-	-	-	-	-	-	-
80.0	75.0	-	-	-	-	-	-	-	-	-	-	-
80.0	90.0	2.9	-	-	-	-	-	-	-	-	-	-
80.0	100.0	0.0	-	-	-	-	-	-	-	-	-	-
80.0	120.0	0.0	-	-	-	-	-	-	-	-	-	-
80.0	160.0	26.3	-	-	-	-	-	-	-	-	-	-

TABLE 4. (cont.)

*Diogenichthys atlanticus* (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
80.0	180.0	2.7	-	-	-	-	-	-	-	-	-	-
80.0	200.0	2.8	-	-	0.0	-	-	-	-	-	-	-
83.0	60.0	0.0	-	-	0.0	-	-	-	-	-	-	-
87.0	45.0	0.0	-	-	0.0	-	-	-	-	-	-	-
87.0	55.0	0.0	-	-	2.8	-	-	-	-	-	-	-
87.0	60.0	0.0	-	-	2.4	-	-	-	-	-	-	-
87.0	65.0	3.1	-	-	2.7	-	-	-	-	-	-	-
87.0	70.0	0.0	-	-	22.5	-	-	-	-	-	-	-
87.0	80.0	0.0	-	-	6.0	-	-	-	-	-	-	-
87.0	90.0	0.0	-	-	2.8	-	-	-	-	-	-	-
90.0	60.0	0.0	-	-	13.4	-	-	-	-	-	-	-
90.0	65.0	3.2	-	-	2.9	-	-	-	-	-	-	-
90.0	80.0	0.0	-	-	16.1	-	-	-	-	-	-	-
90.0	90.0	0.0	-	-	0.0	-	-	-	-	-	-	-
90.0	100.0	0.0	-	-	0.0	-	-	-	-	-	-	-
90.0	120.0	5.4	-	-	-	-	-	-	-	-	-	-
90.0	140.0	0.0	-	-	-	-	-	-	-	-	-	-
90.0	160.0	2.6	-	-	-	-	-	-	-	-	-	-
90.0	180.0	8.2	-	-	-	-	-	-	-	-	-	-
90.0	200.0	2.8	-	-	-	-	-	-	-	-	-	-
93.0	28.0	3.1	-	-	0.0	-	-	-	-	-	-	-
93.0	35.0	0.0	-	-	0.0	-	-	-	-	-	-	-
93.0	65.0	0.0	-	-	0.0	-	-	-	-	-	-	-
93.0	70.0	5.9	-	-	0.0	-	-	-	-	-	-	-
93.0	80.0	3.0	-	-	6.3	-	-	-	-	-	-	-
93.0	90.0	0.0	-	-	37.0	-	-	-	-	-	-	-
93.0	100.0	-	-	-	25.5	-	-	-	-	-	-	-
97.0	40.0	0.0	-	-	3.2	-	-	-	-	-	-	-
97.0	65.0	0.0	-	-	0.0	-	-	-	-	-	-	-
97.0	70.0	3.0	-	-	-	-	-	-	-	-	-	-
97.0	80.0	3.3	-	-	-	-	-	-	-	-	-	-
97.0	85.0	-	-	-	7.2	-	-	-	-	-	-	-
100.0	35.0	0.0	-	-	3.2	-	-	-	-	-	-	-
100.0	45.0	0.0	-	-	2.3	-	-	-	-	-	-	-
100.0	50.0	0.0	-	-	12.0	-	-	-	-	-	-	-
100.0	55.0	0.0	-	-	6.0	-	-	-	-	-	-	-
100.0	80.0	2.9	-	-	6.7	-	-	-	-	-	-	-
100.0	85.0	-	-	-	31.1	-	-	-	-	-	-	-
100.0	90.0	6.1	-	-	13.2	-	-	-	-	-	-	-
100.0	100.0	-	-	-	-	-	-	-	-	-	-	-
100.0	120.0	-	-	-	17.7	-	-	-	-	-	-	-
103.0	45.0	0.0	-	-	3.0	-	-	-	-	-	-	-
103.0	50.0	0.0	-	-	3.3	-	-	-	-	-	-	-
103.0	65.0	0.0	-	-	3.2	-	-	-	-	-	-	-
103.0	70.0	6.0	-	-	9.3	-	-	-	-	-	-	-
103.0	75.0	-	-	-	25.5	-	-	-	-	-	-	-

TABLE 4. (cont.)

*Diogenichthys atlanticus* (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
103.0	80.0	-	0.0	-	8.9	-	0.0	-	-	-	-	-
103.0	85.0	-	-	5.8	-	-	-	-	-	-	-	-
103.0	90.0	-	6.1	-	18.9	-	0.0	-	-	-	-	-
105.0	140.0	-	16.7	-	-	-	-	-	-	-	-	-
107.0	31.0	-	0.0	-	0.0	-	0.0	-	-	-	-	-
107.0	35.0	-	3.2	-	0.0	-	0.0	-	-	-	-	-
107.0	50.0	-	0.0	-	0.0	-	3.3	-	-	-	-	-
107.0	60.0	-	9.1	-	0.0	-	0.0	-	-	-	-	-
107.0	65.0	-	2.9	-	3.1	-	3.2	-	-	-	-	-
107.0	70.0	-	0.0	-	6.1	-	0.0	-	-	-	-	-
107.0	80.0	-	0.0	-	3.0	-	9.7	-	-	-	-	-
107.0	90.0	-	0.0	-	6.1	-	0.0	-	-	-	-	-
110.0	65.0	-	0.0	-	5.9	-	-	-	-	-	-	-
110.0	70.0	-	2.8	-	0.0	-	-	-	-	-	-	-
110.0	90.0	-	2.6	-	0.0	-	-	-	-	-	-	-
110.0	120.0	-	7.7	-	-	-	-	-	-	-	-	-
110.0	140.0	-	32.4	-	-	-	-	-	-	-	-	-
110.0	160.0	-	19.6	-	-	-	-	-	-	-	-	-
120.0	100.0	-	6.0	-	0.0	-	-	-	-	-	-	-
130.0	70.0	-	2.9	-	-	-	-	-	-	-	-	-

*Diogenichthys laternatus*

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
100.0	40.0	0.0	-	0.0	-	-	11.0	-	-	3.1	-	-
103.0	60.0	0.0	-	0.0	-	0.0	-	-	-	15.4	-	-
103.0	65.0	0.0	-	0.0	-	0.0	-	-	-	60.6	-	-
103.0	70.0	0.0	-	0.0	-	13.4	-	-	-	5.8	-	-
103.0	80.0	-	0.0	-	0.0	3.0	-	-	-	-	-	-
103.0	90.0	-	0.0	-	0.0	3.3	-	-	-	-	-	-
107.0	32.0	-	0.0	-	0.0	0.0	-	-	-	3.0	-	-
107.0	60.0	-	0.0	-	0.0	0.0	-	-	-	-	-	-
107.0	65.0	-	0.0	-	0.0	0.0	-	-	-	-	-	-
110.0	32.0	-	0.0	-	0.0	-	-	-	-	3.3	-	-
110.0	35.0	-	0.0	-	0.0	-	-	-	-	4.3	-	-
110.0	40.0	-	0.0	-	0.0	-	-	-	-	5.7	11.9	-
110.0	45.0	-	0.0	-	0.0	-	-	-	-	19.7	40.8	-
110.0	60.0	-	0.0	-	0.0	-	-	-	-	7.7	10.1	-
110.0	70.0	-	0.0	-	0.0	-	-	-	-	0.0	0.0	-
110.0	85.0	-	9.2	-	-	-	-	-	-	9.1	-	-
110.0	100.0	-	7.8	-	-	-	-	-	-	2.1	-	-
113.0	35.0	-	0.0	-	0.0	-	-	-	-	6.5	-	0.0
113.0	40.0	-	0.0	-	3.2	-	-	-	-	6.5	-	21.6
113.0	45.0	-	0.0	-	9.3	-	-	-	-	24.6	-	59.3
113.0	50.0	-	3.1	-	-	-	-	-	-	0.0	-	21.0

TABLE 4. (cont.)

*Diogenichthys laternatus* (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
113.0	55.0	-	0.0	-	3.6	-	0.0	-	-	3.0	-	-
113.0	60.0	-	0.0	-	3.2	-	3.4	-	-	2.9	-	-
113.0	65.0	-	0.0	-	9.9	-	0.0	-	-	-	-	-
113.0	70.0	-	0.0	-	37.0	-	3.4	-	-	-	-	-
113.0	75.0	-	-	-	13.2	-	-	-	-	-	-	-
113.0	80.0	-	0.0	-	37.1	0.0	-	-	-	-	-	-
113.0	85.0	-	-	-	20.9	-	6.5	-	-	-	-	-
113.0	90.0	0.0	0.0	-	0.0	-	-	-	-	-	-	-
117.0	35.0	0.0	0.0	-	0.0	-	-	-	-	-	-	-
117.0	40.0	0.0	0.0	-	0.0	-	-	-	-	-	-	-
117.0	45.0	0.0	0.0	-	2.9	-	-	-	-	-	-	-
117.0	50.0	0.0	0.0	-	0.0	-	-	-	-	-	-	-
117.0	55.0	0.0	0.0	-	0.0	-	-	-	-	-	-	-
117.0	60.0	0.0	0.0	-	2.9	-	-	-	-	-	-	-
117.0	65.0	0.0	0.0	-	39.6	0.0	-	-	-	-	-	-
117.0	75.0	-	-	-	3.0	-	-	-	-	-	-	-
117.0	80.0	0.0	0.0	-	61.2	0.0	-	-	-	-	-	-
117.0	85.0	-	-	-	52.4	-	-	-	-	-	-	-
117.0	90.0	-	-	-	84.6	6.5	-	-	-	-	-	-
118.0	39.0	0.0	0.0	-	0.0	-	-	-	-	-	-	-
120.0	45.0	0.0	0.0	-	3.1	-	-	-	-	-	-	-
120.0	50.0	0.0	0.0	-	0.0	-	-	-	-	-	-	-
120.0	55.0	0.0	0.0	-	8.9	-	-	-	-	-	-	-
120.0	60.0	0.0	0.0	-	116.4	-	-	-	-	-	-	-
120.0	65.0	0.0	0.0	-	5.8	-	-	-	-	-	-	-
120.0	70.0	0.0	0.0	-	2.8	-	-	-	-	-	-	-
120.0	80.0	0.0	0.0	-	17.5	9.9	-	-	-	-	-	-
120.0	100.0	0.0	0.0	-	2.8	-	-	-	-	-	-	-
123.0	37.0	0.0	0.0	-	17.4	-	-	-	-	-	-	-
123.0	42.0	0.0	0.0	-	5.7	0.0	-	-	-	-	-	-
123.0	45.0	0.0	0.0	-	24.5	2.9	-	-	-	-	-	-
123.0	50.0	0.0	0.0	-	8.7	-	-	-	-	-	-	-
123.0	55.0	0.0	0.0	-	58.6	40.2	-	-	-	-	-	-
123.0	60.0	0.0	0.0	-	6.0	10.0	-	-	-	-	-	-
123.0	65.0	0.0	0.0	-	0.0	-	-	-	-	-	-	-
123.0	70.0	0.0	0.0	-	25.0	-	-	-	-	-	-	-
123.0	75.0	-	-	-	14.1	7.7	-	-	-	-	-	-
123.0	80.0	-	-	-	-	40.6	-	-	-	-	-	-
123.0	85.0	-	-	-	-	-	-	-	-	-	-	-
127.0	33.0	-	-	-	-	-	-	-	-	-	-	-
127.0	34.0	-	-	-	-	-	-	-	-	-	-	-
127.0	40.0	-	-	-	-	-	-	-	-	-	-	-
127.0	45.0	-	-	-	-	-	-	-	-	-	-	-
127.0	50.0	-	-	-	-	-	-	-	-	-	-	-
127.0	55.0	-	-	-	-	-	-	-	-	-	-	-
127.0	60.0	-	-	-	-	-	-	-	-	-	-	-

TABLE 4. (cont.)

*Diogenichthys laternatus* (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
127.0	65.0	-	8.7	-	8.8	-	-	2.9	-	-	-	-
127.0	70.0	-	38.2	-	11.5	-	-	82.4	-	-	-	-
127.0	80.0	-	2.9	-	90.5	-	-	68.3	-	-	-	-
130.0	35.0	-	10.0	-	100.8	-	-	35.7	0.0	-	-	-
130.0	40.0	-	52.0	-	30.4	-	-	0.0	93.0	-	-	-
130.0	45.0	-	19.5	-	20.6	-	-	0.0	33.3	-	-	-
130.0	50.0	-	66.2	-	23.4	-	-	109.2	10.0	-	-	-
130.0	55.0	-	14.3	-	21.6	-	-	41.0	86.8	-	-	-
130.0	60.0	-	50.2	-	9.2	-	-	59.0	6.3	-	-	-
130.0	65.0	-	-	-	6.2	-	-	-	-	-	-	-
130.0	70.0	-	17.2	-	-	-	-	27.4	-	-	-	-
130.0	80.0	-	5.6	-	-	-	-	62.8	-	-	-	-
130.0	90.0	-	16.0	-	-	-	-	43.4	-	-	-	-
130.0	100.0	-	8.3	-	-	-	-	8.5	-	-	-	-
130.0	120.0	-	44.2	-	-	-	-	8.1	-	-	-	-
133.0	23.0	-	0.0	-	2.6	-	-	0.0	-	-	-	-
133.0	25.0	-	0.0	-	12.2	-	-	0.0	-	-	-	-
133.0	30.0	-	0.0	-	15.5	-	-	0.0	-	-	-	-
133.0	35.0	-	37.4	-	27.8	-	-	0.0	-	-	-	-
133.0	40.0	-	38.7	-	18.5	-	-	72.8	-	-	-	-
133.0	45.0	-	111.3	-	36.2	-	-	18.6	-	-	-	-
133.0	50.0	-	28.3	-	17.8	-	-	0.0	-	-	-	-
133.0	55.0	-	35.6	-	20.7	-	-	10.6	-	-	-	-
133.0	60.0	-	11.5	-	39.3	-	-	36.4	-	-	-	-
133.0	65.0	-	2.9	-	77.3	-	-	15.8	-	-	-	-
133.0	70.0	-	5.7	-	0.0	-	-	211.7	-	-	-	-
133.0	80.0	-	8.7	-	-	-	-	53.5	-	-	-	-
137.0	30.0	-	49.3	-	108.8	-	-	0.0	9.1	-	-	-
137.0	35.0	-	62.2	-	102.3	-	-	0.0	6.2	-	-	-
137.0	40.0	-	21.6	-	78.0	-	-	11.4	56.8	-	-	-
137.0	45.0	-	17.4	-	18.2	-	-	9.8	54.7	-	-	-
137.0	50.0	-	3.3	-	13.6	-	-	23.4	3.0	-	-	-
137.0	55.0	-	3.1	-	22.3	-	-	38.6	-	-	-	-
137.0	60.0	-	-	-	27.6	-	-	38.9	-	-	-	-
137.0	65.0	-	-	-	19.4	-	-	-	-	-	-	-
137.0	70.0	-	9.0	-	25.3	-	-	-	-	-	-	-
137.0	80.0	-	11.8	-	-	-	-	-	4.9	-	-	-
140.0	30.0	-	0.0	-	-	-	-	-	7.5	-	-	-
140.0	35.0	-	9.1	-	-	-	-	-	2.8	-	-	-
140.0	40.0	-	14.8	-	-	-	-	-	21.3	-	-	-
140.0	45.0	-	15.3	-	-	-	-	-	5.6	-	-	-
140.0	50.0	-	-	-	-	-	-	-	60.7	-	-	-
										138.6		

TABLE 4. (cont.)

*Electrona rissoii*

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
70.0	120.0	3.0	-	-	-	-	-	0.0	-	-	-	-
90.0	100.0	0.0	-	-	-	3.1	-	0.0	-	-	-	-

*Gonichthys tenuiculus*

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
107.0	32.0	-	0.0	-	0.0	-	-	3.4	-	0.0	-	-
107.0	90.0	-	0.0	-	0.0	-	-	3.3	-	-	-	-
113.0	45.0	-	0.0	-	6.2	-	-	0.0	-	2.5	-	-
113.0	50.0	-	0.0	-	9.2	-	-	0.0	-	3.0	-	-
113.0	55.0	-	0.0	-	3.6	-	-	0.0	-	0.0	-	-
113.0	65.0	-	0.0	-	3.3	-	-	0.0	-	-	-	-
113.0	70.0	-	0.0	-	12.3	-	-	0.0	-	-	-	-
113.0	75.0	-	0.0	-	3.3	-	-	0.0	-	-	-	-
113.0	80.0	-	0.0	-	18.5	-	-	0.0	-	-	-	-
113.0	85.0	-	0.0	-	12.0	-	-	0.0	-	-	-	-
117.0	60.0	-	0.0	-	2.9	-	-	0.0	-	-	-	-
117.0	65.0	-	0.0	-	2.8	-	-	0.0	-	-	-	-
117.0	70.0	-	0.0	-	2.8	-	-	0.0	-	-	-	-
117.0	75.0	-	0.0	-	3.0	-	-	0.0	-	-	-	-
117.0	80.0	-	0.0	-	12.2	-	-	0.0	-	-	-	-
117.0	85.0	-	0.0	-	11.6	-	-	0.0	-	-	-	-
117.0	90.0	-	0.0	-	2.8	-	-	3.3	-	-	-	-
120.0	60.0	-	2.9	-	0.0	-	-	0.0	-	-	-	-
120.0	65.0	-	2.8	-	0.0	-	-	0.0	-	-	-	-
120.0	70.0	-	0.0	-	6.6	-	-	0.0	-	-	-	-
120.0	80.0	-	0.0	-	3.4	-	-	0.0	-	-	-	-
120.0	120.0	-	2.9	-	-	-	-	0.0	-	-	-	-
123.0	42.0	-	3.1	-	0.0	-	-	0.0	-	-	-	-
123.0	55.0	-	0.0	-	3.3	-	-	0.0	-	-	-	-
123.0	65.0	-	0.0	-	3.1	-	-	0.0	-	-	-	-
123.0	70.0	-	0.0	-	9.2	-	-	5.5	-	-	-	-
123.0	75.0	-	0.0	-	3.1	-	-	0.0	-	-	-	-
123.0	80.0	-	0.0	-	0.0	-	-	5.6	-	-	-	-
127.0	40.0	-	4.3	-	3.0	-	-	2.8	-	0.0	-	-
127.0	45.0	-	0.0	-	6.2	-	-	0.0	-	-	-	-
127.0	50.0	-	2.9	-	-	-	-	0.0	-	-	-	-
127.0	55.0	-	0.0	-	8.9	-	-	0.0	-	-	-	-
127.0	60.0	-	5.5	-	0.0	-	-	0.0	-	-	-	-
127.0	65.0	-	2.9	-	-	-	-	5.5	-	-	-	-
127.0	80.0	-	0.0	-	-	-	-	0.0	-	-	-	-
130.0	26.0	-	1.0	-	-	-	-	0.0	-	-	-	-
130.0	35.0	-	1.0	-	-	-	-	0.0	-	-	-	-
130.0	40.0	-	6.1	-	-	-	-	0.0	-	-	-	-
130.0	45.0	-	0.0	-	-	-	-	0.0	-	-	-	-

TABLE 4. (cont.)

*Gonichthys tenuiculus* (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
130.0	55.0	-	5.7	-	0.0	-	-	-	5.5	0.0	-	-
130.0	60.0	-	5.6	-	0.0	-	-	-	2.8	3.1	-	-
130.0	70.0	-	0.0	-	0.0	-	-	-	2.5	-	-	-
130.0	80.0	-	8.3	-	-	-	-	-	0.0	-	-	-
130.0	120.0	-	8.3	-	-	-	-	-	0.0	-	-	-
133.0	35.0	-	5.8	-	3.1	-	-	0.0	-	0.0	-	-
133.0	45.0	-	0.0	-	6.0	-	-	0.0	-	0.0	-	-
133.0	50.0	-	5.7	-	0.0	-	-	0.0	-	0.0	-	-
133.0	55.0	-	5.5	-	0.0	-	-	0.0	-	0.0	-	-
133.0	60.0	-	2.9	-	0.0	-	-	0.0	-	0.0	-	-
133.0	65.0	-	0.0	-	6.2	-	-	0.0	-	0.0	-	-
133.0	70.0	-	0.0	-	3.0	-	-	2.7	-	-	-	-
137.0	35.0	-	9.3	-	0.0	-	-	0.0	-	0.0	-	-
137.0	40.0	-	12.3	-	0.0	-	-	0.0	-	0.0	-	-
137.0	45.0	-	14.5	-	0.0	-	-	0.0	-	9.1	-	-
137.0	50.0	-	0.0	-	0.0	-	-	0.0	-	2.6	0.0	-
137.0	55.0	-	0.0	-	6.4	-	-	0.0	-	-	-	-
137.0	65.0	-	-	-	12.9	-	-	-	-	-	-	-
137.0	80.0	-	8.9	-	-	-	-	-	-	2.5	-	-
140.0	35.0	-	3.0	-	-	-	-	-	-	0.0	-	-
140.0	40.0	-	3.0	-	-	-	-	-	-	0.0	-	-
140.0	45.0	-	3.1	-	-	-	-	-	-	2.8	-	-
140.0	50.0	-	3.0	-	-	-	-	-	-	13.9	-	-

*Hygophum* spp.

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
60.0	200.0	12.4	-	-	-	-	-	0.0	-	-	-	-
90.0	80.0	0.0	-	-	0.0	-	-	2.6	-	0.0	-	-
107.0	70.0	-	0.0	-	0.0	-	-	0.0	-	3.0	-	-
117.0	70.0	-	0.0	-	2.8	-	-	-	-	-	-	-
123.0	80.0	-	0.0	-	3.1	-	-	0.0	-	-	-	-
127.0	40.0	-	0.0	-	3.0	-	-	0.0	-	0.0	-	-
127.0	45.0	-	0.0	-	12.4	-	-	0.0	-	0.0	-	-
127.0	55.0	-	0.0	-	11.9	-	-	0.0	-	3.2	-	-
127.0	60.0	-	0.0	-	0.0	-	-	0.0	-	6.7	-	-
130.0	55.0	-	0.0	-	2.9	-	-	0.0	-	0.0	-	-

*Hygophum atratum*

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
103.0	65.0	0.0	-	-	0.0	-	-	0.0	-	16.0	-	-
103.0	70.0	0.0	-	-	0.0	-	-	3.4	-	2.9	-	-
107.0	65.0	-	0.0	-	0.0	-	-	0.0	-	3.3	-	-

TABLE 4. (cont.)

*Hygophum atratum* (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
110.0	65.0	-	5.3	-	2.9	-	-	0.0	0.0	-	-	-
113.0	35.0	-	0.0	-	7.0	-	0.0	0.0	0.0	-	-	-
113.0	40.0	-	0.0	-	0.0	-	0.0	-	-	3.1	-	-
113.0	45.0	-	0.0	-	3.1	-	0.0	-	-	2.5	-	-
113.0	50.0	-	0.0	-	9.2	-	0.0	-	-	6.0	-	-
113.0	55.0	-	0.0	-	3.6	-	0.0	-	-	3.0	-	-
113.0	75.0	-	0.0	-	3.3	-	-	-	-	-	-	-
113.0	80.0	-	0.0	-	9.3	-	-	-	-	-	-	-
117.0	80.0	-	0.0	-	12.2	0.0	-	-	-	-	-	-
117.0	85.0	-	0.0	-	2.9	-	-	-	-	-	-	-
120.0	40.0	-	0.0	-	1.9	-	0.0	-	-	0.0	-	-
120.0	50.0	-	0.0	-	0.0	-	-	-	-	-	-	-
120.0	60.0	-	0.0	-	0.0	-	-	-	-	-	-	-
120.0	65.0	-	0.0	-	0.0	-	-	-	-	-	-	-
120.0	70.0	-	0.0	-	2.9	-	9.2	-	-	0.0	-	-
123.0	45.0	-	0.0	-	2.8	-	0.0	-	-	0.0	-	-
123.0	50.0	-	0.0	-	5.9	-	0.0	-	-	0.0	-	-
123.0	60.0	-	0.0	-	0.0	-	-	-	-	-	-	-
123.0	65.0	-	0.0	-	0.0	-	6.2	-	-	5.6	-	-
123.0	70.0	-	0.0	-	2.8	-	9.2	-	-	2.8	-	-
127.0	33.0	-	0.0	-	2.8	-	0.0	-	-	0.0	-	-
127.0	40.0	-	0.0	-	2.8	-	0.0	-	-	0.0	-	-
127.0	45.0	-	0.0	-	3.0	-	0.0	-	-	0.0	-	-
127.0	50.0	-	0.0	-	0.0	-	18.2	-	-	0.0	-	-
127.0	55.0	-	0.0	-	2.7	-	0.0	-	-	5.5	-	-
127.0	60.0	-	0.0	-	2.7	-	3.0	-	-	0.0	-	-
127.0	65.0	-	0.0	-	2.9	-	3.1	-	-	0.0	-	-
127.0	70.0	-	0.0	-	5.9	-	2.9	-	-	2.8	-	-
127.0	80.0	-	0.0	-	0.0	-	0.0	-	-	10.9	-	-
130.0	35.0	-	0.0	-	3.1	-	3.1	-	-	0.0	-	-
130.0	40.0	-	0.0	-	12.6	-	-	-	-	2.6	-	-
130.0	45.0	-	0.0	-	0.0	-	-	-	-	5.7	-	-
130.0	50.0	-	0.0	-	2.9	-	3.0	-	-	0.0	-	-
130.0	55.0	-	0.0	-	2.9	-	0.0	-	-	10.9	-	-
130.0	60.0	-	0.0	-	5.6	-	3.1	-	-	5.6	-	-
130.0	65.0	-	0.0	-	3.1	-	3.1	-	-	0.0	-	-
130.0	70.0	-	0.0	-	6.2	-	-	-	-	-	-	-
133.0	25.0	-	0.0	-	0.0	-	0.0	-	-	0.0	-	-
133.0	30.0	-	0.0	-	0.0	-	0.0	-	-	0.0	-	-
133.0	35.0	-	0.0	-	8.6	-	3.1	-	-	0.0	-	-
133.0	40.0	-	0.0	-	3.0	-	3.1	-	-	11.6	-	-
133.0	45.0	-	0.0	-	3.0	-	0.0	-	-	2.7	-	-
133.0	50.0	-	0.0	-	8.8	-	0.0	-	-	10.0	-	-
133.0	60.0	-	0.0	-	2.8	-	0.0	-	-	40.3	-	-
133.0	65.0	-	0.0	-	2.9	-	0.0	-	-	2.8	-	-
133.0	70.0	-	0.0	-	3.1	-	3.1	-	-	2.6	-	-

TABLE 4. (cont.)

*Hygophum atratum* (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
133.0	80.0	-	2.9	-	-	-	-	10.2	-	6.2	-	-
137.0	35.0	-	15.5	-	0.0	-	-	-	0.0	6.5	18.2	-
137.0	45.0	-	8.7	-	6.1	-	-	-	0.0	13.0	0.0	-
137.0	50.0	-	6.6	-	0.0	-	-	-	-	2.2	-	-
137.0	60.0	-	-	-	0.0	-	-	-	-	-	-	-
137.0	65.0	-	-	-	3.2	-	-	-	-	2.4	-	-
137.0	70.0	-	0.0	-	3.2	-	-	-	-	0.0	-	-
137.0	80.0	-	8.9	-	-	-	-	-	-	0.0	-	-
140.0	40.0	-	8.9	-	-	-	-	-	-	33.1	-	-
140.0	45.0	-	6.1	-	-	-	-	-	-	30.0	-	-
140.0	50.0	-	3.0	-	-	-	-	-	-	-	-	-

*Hygophum reinhardtii*

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
60.0	160.0	6.5	-	-	-	-	-	0.0	-	-	-	-
60.0	200.0	0.0	-	-	-	-	-	19.2	-	-	-	-
70.0	200.0	6.1	-	-	-	-	-	2.7	-	-	-	-
80.0	160.0	11.7	-	-	-	-	-	33.7	-	-	-	-
80.0	200.0	0.0	-	-	-	-	-	5.4	-	-	-	-
90.0	120.0	2.7	-	-	-	-	-	8.0	-	-	-	-
90.0	140.0	0.0	-	-	-	-	-	27.9	-	-	-	-
90.0	180.0	5.5	-	-	-	-	-	2.8	-	-	-	-
90.0	200.0	8.4	-	-	-	-	-	0.0	-	2.8	-	-
97.0	60.0	0.0	-	-	-	-	-	0.0	-	5.8	-	-
97.0	80.0	0.0	-	-	-	-	-	0.0	-	-	-	-
100.0	90.0	3.0	-	-	0.0	-	-	0.0	-	3.0	-	-
100.0	120.0	-	0.0	-	-	-	-	-	-	0.0	-	-
105.0	140.0	-	2.8	-	-	-	-	-	-	2.7	-	-
110.0	120.0	-	0.0	-	-	-	-	-	-	29.2	-	-
110.0	160.0	-	0.0	-	-	-	-	3.3	-	-	-	-
113.0	75.0	-	-	-	0.0	-	-	3.3	-	-	-	-
113.0	90.0	-	0.0	-	-	-	-	0.0	-	0.0	-	-
127.0	80.0	-	2.9	-	-	-	-	-	-	19.0	-	-
130.0	80.0	-	2.8	-	-	-	-	-	-	0.0	-	-
130.0	90.0	-	0.0	-	-	-	-	-	-	0.0	-	-
130.0	120.0	-	35.9	-	-	-	-	-	-	2.5	-	-
137.0	80.0	-	0.0	-	-	-	-	-	-	-	-	-

*Loweina rara*

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
100.0	90.0	0.0	-	2.8	-	0.0	-	-	2.9	-	-	-
105.0	140.0	-	-	-	-	-	-	-	0.0	-	-	-

TABLE 4. (cont.)

*Loweina rara* (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
110.0	100.0	-	2.6	-	-	-	-	0.0	-	-	-	-
110.0	140.0	-	2.7	-	-	-	-	0.0	-	-	-	-
110.0	160.0	-	0.0	-	-	-	-	2.9	-	-	-	-
117.0	90.0	-	2.9	-	0.0	-	-	0.0	-	-	-	-
130.0	80.0	-	2.8	-	-	-	-	-	0.0	-	-	-
130.0	90.0	-	0.0	-	-	-	-	-	2.7	-	-	-

*Myctophum nitidulum*

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
60.0	200.0	3.1	-	-	-	-	-	0.0	-	-	-	-
80.0	140.0	2.7	-	-	-	-	-	-	-	-	-	-
80.0	160.0	2.9	-	-	-	-	-	0.0	-	-	-	-
90.0	160.0	7.8	-	-	-	-	-	2.8	-	-	-	-
90.0	180.0	2.7	-	-	-	-	-	0.0	-	-	-	-
105.0	140.0	-	8.3	-	-	-	-	0.0	-	-	-	-
107.0	80.0	-	5.8	-	0.0	-	-	0.0	-	-	-	-
110.0	60.0	-	0.0	-	3.0	-	-	2.7	-	0.0	0.0	-
110.0	120.0	-	5.2	-	-	-	-	5.5	-	0.0	-	-
110.0	140.0	-	2.7	-	-	-	-	0.0	-	0.0	-	-
113.0	30.0	-	1.1	-	-	-	-	2.8	-	-	-	-
117.0	65.0	-	0.0	-	-	-	-	3.1	-	-	-	-
117.0	80.0	-	0.0	-	-	-	-	0.0	-	-	-	-
123.0	50.0	-	0.0	-	-	-	-	-	2.8	-	0.0	-
130.0	100.0	-	2.8	-	-	-	-	-	0.0	-	-	-
130.0	120.0	-	5.5	-	-	-	-	-	0.0	-	-	-

*Protomyctophum crockeri*

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
60.0	60.0	2.7	-	-	-	-	-	0.0	-	-	-	-
60.0	70.0	13.0	-	-	-	-	-	0.0	-	-	-	-
60.0	100.0	3.1	-	-	-	-	-	0.0	-	-	-	-
60.0	120.0	7.4	-	-	-	-	-	0.0	-	-	-	-
60.0	140.0	0.0	-	-	-	-	-	2.6	-	-	-	-
60.0	160.0	16.1	-	-	-	-	-	0.0	-	-	-	-
60.0	200.0	12.4	-	-	-	-	-	0.0	-	-	-	-
63.0	60.0	0.0	-	-	-	-	-	2.5	-	-	-	-
63.0	70.0	-	-	-	-	-	-	2.3	-	-	-	-
67.0	55.0	0.0	-	-	-	-	-	0.0	-	-	5.8	-
67.0	60.0	3.0	-	-	-	-	-	0.5	-	3.0	-	-
67.0	65.0	-	-	-	-	-	-	5.8	-	-	-	-
67.0	70.0	-	-	-	-	-	-	2.8	-	-	-	-
67.0	80.0	-	-	-	-	-	-	2.7	-	-	-	-

TABLE 4. (cont.)

## Protomyctophum crockeri (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
70.0	52.0	0.0	-	-	5.2	-	-	-	-	-	-	-
70.0	53.0	0.0	-	-	5.4	-	-	-	-	-	-	-
70.0	55.0	2.9	-	-	-	-	-	-	-	-	-	-
70.0	70.0	5.4	-	-	0.0	-	-	-	-	-	-	-
70.0	80.0	0.0	-	-	0.0	-	-	-	-	-	-	-
70.0	90.0	0.0	-	-	-	-	-	-	-	-	-	-
70.0	100.0	0.0	-	-	-	-	-	-	-	-	-	-
70.0	120.0	8.9	-	-	-	-	-	-	-	-	-	-
70.0	160.0	5.9	-	-	-	-	-	-	-	-	-	-
73.0	53.0	2.7	-	-	3.0	-	-	-	-	-	-	-
73.0	60.0	5.6	-	-	0.0	-	-	-	-	-	-	-
73.0	70.0	-	-	-	2.8	-	-	-	-	-	-	-
73.0	80.0	-	-	-	2.6	-	-	-	-	-	-	-
77.0	51.0	0.0	-	-	0.0	-	-	-	-	-	-	-
77.0	55.0	2.9	-	-	0.0	-	-	-	-	-	-	-
77.0	57.0	3.0	-	-	2.5	-	-	-	-	-	-	-
77.0	65.0	-	-	-	5.1	-	-	-	-	-	-	-
80.0	52.0	0.0	-	-	0.0	-	-	-	-	-	-	-
80.0	55.0	0.0	-	-	3.2	-	-	-	-	-	-	-
80.0	60.0	0.0	-	-	6.8	-	-	-	-	-	-	-
80.0	65.0	6.2	-	-	13.5	-	-	-	-	-	-	-
80.0	70.0	2.8	-	-	0.0	-	-	-	-	-	-	-
80.0	80.0	9.0	-	-	0.0	-	-	-	-	-	-	-
80.0	90.0	2.9	-	-	-	-	-	-	-	-	-	-
80.0	100.0	0.0	-	-	-	-	-	-	-	-	-	-
80.0	120.0	5.9	-	-	-	-	-	-	-	-	-	-
83.0	60.0	0.0	-	-	0.0	-	-	-	-	-	-	-
83.0	65.0	0.0	-	-	2.9	-	-	-	-	-	-	-
83.0	70.0	3.2	-	-	6.3	-	-	-	-	-	-	-
83.0	80.0	3.5	-	-	0.0	-	-	-	-	-	-	-
87.0	45.0	0.0	-	-	0.0	-	-	-	-	-	-	-
87.0	55.0	0.0	-	-	0.0	-	-	-	-	-	-	-
87.0	60.0	6.0	-	-	4.8	-	-	-	-	-	-	-
87.0	65.0	0.0	-	-	0.0	-	-	-	-	-	-	-
87.0	70.0	3.1	-	-	6.4	-	-	-	-	-	-	-
87.0	80.0	6.1	-	-	3.0	-	-	-	-	-	-	-
87.0	90.0	6.0	-	-	5.6	-	-	-	-	-	-	-
87.0	100.0	-	-	-	6.2	-	-	-	-	-	-	-
90.0	32.0	2.7	-	-	0.0	-	-	-	-	-	-	-
90.0	45.0	0.0	-	-	0.0	-	-	-	-	-	-	-
90.0	60.0	3.0	-	-	3.3	-	-	-	-	-	-	-
90.0	65.0	3.2	-	-	8.8	-	-	-	-	-	-	-
90.0	70.0	6.2	-	-	0.0	-	-	-	-	-	-	-
90.0	80.0	0.0	-	-	0.0	-	-	-	-	-	-	-
90.0	85.0	-	-	-	0.0	-	-	-	-	-	-	-
90.0	90.0	0.0	-	-	-	-	-	-	-	-	-	-

TABLE 4. (cont.)

*Protomyctophum crockeri* (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
90.0	100.0	5.5	-	0.0	-	-	0.0	-	-	-	-	-
90.0	160.0	2.6	-	0.0	-	-	0.0	-	-	-	-	-
93.0	28.0	0.0	-	0.0	-	-	0.0	-	-	-	2.7	0.0
93.0	30.0	6.1	-	0.0	-	-	0.0	-	-	-	3.7	0.0
93.0	35.0	0.0	2.9	-	-	-	0.0	-	-	-	3.2	0.0
93.0	40.0	3.0	-	3.3	-	-	3.2	-	-	-	8.7	0.0
93.0	50.0	0.0	-	0.0	-	-	0.0	-	-	-	2.9	0.0
93.0	55.0	3.1	-	0.0	-	-	0.0	-	-	-	5.6	0.0
93.0	60.0	6.1	-	0.0	-	-	0.0	-	-	-	0.0	0.0
93.0	65.0	0.0	-	6.0	-	-	6.5	-	-	-	0.0	0.0
93.0	70.0	2.9	-	0.0	-	-	0.0	-	-	-	2.9	0.0
93.0	75.0	-	-	6.3	-	-	3.2	-	-	-	3.0	0.0
93.0	80.0	3.0	-	-	-	-	6.3	-	-	-	0.0	0.0
93.0	85.0	-	-	0.0	-	-	6.3	-	-	-	6.2	0.0
93.0	90.0	3.2	-	2.8	-	-	0.0	-	-	-	3.0	0.0
93.0	100.0	-	-	0.0	-	-	0.0	-	-	-	3.0	0.0
97.0	32.0	3.1	0.0	-	-	-	0.0	-	-	-	0.0	0.0
97.0	35.0	3.1	3.2	-	-	-	3.3	-	-	-	0.0	0.0
97.0	40.0	0.0	3.2	-	-	-	0.0	-	-	-	3.0	0.0
97.0	45.0	0.0	13.8	-	-	-	0.0	-	-	-	3.0	0.0
97.0	50.0	0.0	0.0	-	-	-	3.3	-	-	-	3.0	0.0
97.0	55.0	3.0	0.0	-	-	-	0.0	-	-	-	0.0	0.0
97.0	65.0	0.0	3.2	-	-	-	3.0	-	-	-	3.0	0.0
97.0	70.0	5.9	13.8	-	-	-	0.0	-	-	-	6.2	0.0
97.0	80.0	3.3	0.0	-	-	-	9.7	-	-	-	0.0	0.0
97.0	90.0	6.1	0.0	-	-	-	0.0	-	-	-	0.0	0.0
100.0	30.0	2.8	0.0	-	-	-	0.0	-	-	-	0.0	0.0
100.0	35.0	0.0	0.0	-	-	-	0.0	-	-	-	0.0	0.0
100.0	40.0	0.0	3.2	-	-	-	0.0	-	-	-	0.0	0.0
100.0	45.0	0.0	2.3	-	-	-	0.0	-	-	-	0.0	0.0
100.0	50.0	0.0	12.0	-	-	-	0.0	-	-	-	0.0	0.0
100.0	55.0	0.0	9.1	-	-	-	0.0	-	-	-	0.0	0.0
100.0	60.0	8.9	10.1	-	-	-	0.0	-	-	-	0.0	0.0
100.0	70.0	15.4	0.0	-	-	-	0.0	-	-	-	0.0	0.0
100.0	75.0	-	6.7	-	-	-	0.0	-	-	-	0.0	0.0
100.0	80.0	0.0	3.3	-	-	-	0.0	-	-	-	0.0	0.0
100.0	85.0	-	10.4	-	-	-	0.0	-	-	-	0.0	0.0
100.0	90.0	3.0	6.6	-	-	-	0.0	-	-	-	0.0	0.0
103.0	35.0	2.9	0.0	-	-	-	0.0	-	-	-	3.4	0.0
103.0	40.0	5.8	0.0	-	-	-	0.0	-	-	-	6.6	0.0
103.0	45.0	0.0	6.1	-	-	-	0.0	-	-	-	0.0	0.0
103.0	50.0	0.0	6.6	-	-	-	0.0	-	-	-	0.0	0.0
103.0	55.0	2.9	0.0	-	-	-	0.0	-	-	-	6.6	0.0
103.0	60.0	6.1	0.0	-	-	-	0.0	-	-	-	6.6	0.0
103.0	65.0	2.8	0.0	-	-	-	0.0	-	-	-	3.1	0.0
103.0	70.0	0.0	6.2	-	-	-	0.0	-	-	-	6.2	0.0

TABLE 4. (cont.)

*Protomyctophum crockeri* (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
103.0	75.0	-	-	-	9.6	-	-	-	-	-	-	-
103.0	80.0	-	3.0	-	8.9	-	-	-	-	-	-	-
103.0	90.0	-	3.0	-	0.0	-	-	-	-	-	-	-
107.0	32.0	-	3.1	-	6.1	-	-	-	-	-	-	-
107.0	35.0	-	6.4	-	0.0	-	-	-	-	-	-	-
107.0	40.0	-	0.0	-	0.0	-	-	-	-	-	-	-
107.0	45.0	-	0.0	-	8.9	-	-	-	-	-	-	-
107.0	60.0	-	0.0	-	0.0	-	-	-	-	-	-	-
107.0	65.0	-	11.8	-	3.1	-	-	-	-	-	-	-
107.0	70.0	-	15.9	-	0.0	-	-	-	-	-	-	-
107.0	75.0	-	-	-	3.1	-	-	-	-	-	-	-
107.0	80.0	-	0.0	-	3.0	-	-	-	-	-	-	-
107.0	90.0	-	0.0	-	0.0	-	-	-	-	-	-	-
110.0	35.0	-	0.0	-	0.0	-	-	-	-	-	-	-
110.0	40.0	-	2.8	-	2.8	-	-	-	-	-	-	-
110.0	45.0	-	5.6	-	0.0	-	-	-	-	-	-	-
110.0	55.0	-	2.7	-	0.0	-	-	-	-	-	-	-
110.0	60.0	-	5.4	-	3.0	-	-	-	-	-	-	-
110.0	65.0	-	2.7	-	11.7	-	-	-	-	-	-	-
110.0	70.0	-	0.0	-	6.5	-	-	-	-	-	-	-
110.0	75.0	-	-	-	6.4	-	-	-	-	-	-	-
110.0	80.0	-	10.3	-	0.0	-	-	-	-	-	-	-
110.0	85.0	-	-	-	3.1	-	-	-	-	-	-	-
110.0	90.0	-	0.0	-	3.1	-	-	-	-	-	-	-
110.0	100.0	-	5.2	-	-	-	-	-	-	-	-	-
110.0	120.0	-	5.2	-	-	-	-	-	-	-	-	-
113.0	40.0	-	0.0	-	9.7	-	-	-	-	-	-	-
113.0	45.0	-	0.0	-	6.2	-	-	-	-	-	-	-
113.0	50.0	-	0.0	-	0.0	-	-	-	-	-	-	-
113.0	65.0	-	0.0	-	6.6	-	-	-	-	-	-	-
113.0	70.0	-	0.0	-	3.1	-	-	-	-	-	-	-
113.0	75.0	-	-	-	6.6	-	-	-	-	-	-	-
113.0	80.0	-	0.0	-	3.1	-	-	-	-	-	-	-
113.0	85.0	-	-	-	3.0	-	-	-	-	-	-	-
113.0	90.0	-	0.0	-	3.4	-	-	-	-	-	-	-
117.0	40.0	-	0.0	-	0.0	-	-	-	-	-	-	-
117.0	45.0	-	-	-	2.9	-	-	-	-	-	-	-
117.0	50.0	-	-	-	0.0	-	-	-	-	-	-	-
117.0	55.0	-	-	-	3.1	-	-	-	-	-	-	-
117.0	65.0	-	-	-	0.0	-	-	-	-	-	-	-
117.0	70.0	-	-	-	0.0	-	-	-	-	-	-	-
117.0	75.0	-	-	-	6.7	-	-	-	-	-	-	-
117.0	80.0	-	-	-	0.0	-	-	-	-	-	-	-
117.0	85.0	-	-	-	3.4	-	-	-	-	-	-	-
117.0	90.0	-	-	-	3.3	-	-	-	-	-	-	-
120.0	50.0	-	-	-	0.0	-	-	-	-	-	-	-
120.0	55.0	-	-	-	0.0	-	-	-	-	-	-	-
120.0	60.0	-	-	-	5.8	-	-	-	-	-	-	-

TABLE 4. (cont.)

*Protomyctophum crockeri* (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
120.0	80.0	-	2.8	-	0.0	-	-	-	0.0	-	-	-
120.0	90.0	-	5.8	-	-	-	-	-	0.0	-	-	-
120.0	100.0	-	6.0	-	0.0	-	-	-	2.8	-	0.0	-
123.0	50.0	-	0.0	-	0.0	-	-	-	0.0	-	0.0	-
123.0	55.0	-	6.0	-	0.0	-	-	-	0.0	-	0.0	-
123.0	60.0	-	2.9	-	0.0	-	-	-	2.7	-	3.1	-
123.0	65.0	-	2.9	-	0.0	-	-	-	0.0	-	-	-
123.0	70.0	-	0.0	-	9.2	-	-	-	0.0	-	-	-
123.0	80.0	-	0.0	-	0.0	-	-	-	5.6	-	-	-
127.0	40.0	-	0.0	-	3.0	-	-	-	0.0	-	0.0	-
127.0	45.0	-	0.0	-	0.0	-	-	-	0.0	-	3.1	-
127.0	50.0	-	2.9	-	0.0	-	-	-	0.0	-	0.0	-
127.0	60.0	-	2.7	-	0.0	-	-	-	0.0	-	0.0	-
127.0	80.0	-	2.9	-	0.0	-	-	-	0.0	-	0.0	-
130.0	40.0	-	0.0	-	3.2	-	-	-	0.0	-	0.0	-
130.0	50.0	-	0.0	-	0.0	-	-	-	2.7	-	0.0	-
130.0	60.0	-	0.0	-	0.0	-	-	-	2.8	-	0.0	-
130.0	70.0	-	8.6	-	0.0	-	-	-	0.0	-	-	-
130.0	100.0	-	2.8	-	-	-	-	-	0.0	-	-	-
130.0	120.0	-	2.8	-	-	-	-	-	0.0	-	-	-
133.0	30.0	-	0.0	-	3.1	-	-	-	0.0	-	0.0	-
133.0	45.0	-	0.0	-	3.0	-	-	-	0.0	-	0.0	-
133.0	70.0	-	0.0	-	3.0	-	-	-	0.0	-	2.6	-
137.0	50.0	-	0.0	-	0.0	-	-	-	-	-	-	-

*Symbolophorus californiensis*

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
60.0	100.0	3.1	-	-	-	-	-	-	0.0	-	-	-
60.0	140.0	3.5	-	-	-	-	-	-	18.3	-	-	-
60.0	160.0	0.0	-	-	-	-	-	-	2.9	-	-	-
60.0	180.0	3.0	-	-	-	-	-	-	2.7	-	-	-
70.0	50.0	0.0	-	-	-	-	-	-	2.2	-	0.0	-
70.0	60.0	0.0	-	-	-	-	-	-	0.0	-	2.7	-
70.0	90.0	2.9	-	-	-	-	-	-	5.5	-	2.8	-
70.0	100.0	0.0	-	-	-	-	-	-	16.4	-	-	-
70.0	120.0	20.9	-	-	-	-	-	-	-	-	-	-
70.0	160.0	11.9	-	-	-	-	-	-	-	-	-	-
73.0	65.0	-	-	-	-	-	-	-	2.9	-	-	-
77.0	70.0	-	-	-	-	-	-	-	2.3	-	-	-
77.0	80.0	-	-	-	-	-	-	-	2.8	-	-	-
80.0	60.0	0.0	-	-	-	-	-	-	5.6	-	0.0	-
80.0	80.0	3.0	-	-	-	-	-	-	3.0	-	-	-
80.0	100.0	3.0	-	-	-	-	-	-	6.0	-	-	-
80.0	120.0	0.0	-	-	-	-	-	-	5.9	-	-	-

TABLE 4. (cont.)

*Symbolophorus californiensis* (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
80.0	160.0	5.8	-	-	-	-	-	-	-	-	-	-
80.0	180.0	2.7	-	-	-	-	-	-	-	-	-	-
83.0	70.0	0.0	-	-	0.0	-	3.4	-	-	-	-	-
87.0	60.0	3.0	-	-	7.2	-	0.0	-	-	-	-	-
87.0	70.0	3.1	-	-	0.0	-	0.0	-	-	-	-	-
87.0	80.0	3.0	-	-	0.0	-	3.3	-	-	-	-	-
87.0	90.0	0.0	-	-	0.0	-	12.5	-	-	-	-	-
87.0	100.0	-	-	-	3.1	-	-	-	-	-	-	-
90.0	45.0	0.0	-	-	3.1	-	-	-	-	-	-	-
90.0	60.0	3.0	-	-	13.4	-	-	-	-	-	-	-
90.0	65.0	0.0	-	-	2.9	-	-	-	-	-	-	-
90.0	70.0	3.1	-	-	6.2	-	-	-	-	-	-	-
90.0	80.0	0.0	-	-	22.5	-	-	-	-	-	-	-
90.0	90.0	0.0	-	-	3.2	-	-	-	-	-	-	-
90.0	100.0	11.0	-	-	0.0	-	-	-	-	-	-	-
90.0	120.0	2.7	-	-	-	-	-	-	-	-	-	-
93.0	35.0	0.0	-	-	0.0	-	-	-	-	-	-	-
93.0	65.0	0.0	-	-	0.0	-	-	-	-	-	-	-
93.0	70.0	14.7	-	-	0.0	-	-	-	-	-	-	-
93.0	80.0	0.0	-	-	3.2	-	-	-	-	-	-	-
93.0	85.0	-	-	-	-	-	-	-	-	-	-	-
93.0	90.0	0.0	-	-	10.1	-	-	-	-	-	-	-
93.0	100.0	-	-	-	11.3	-	-	-	-	-	-	-
97.0	35.0	0.0	-	-	3.5	-	-	-	-	-	-	-
97.0	45.0	0.0	-	-	0.0	-	-	-	-	-	-	-
97.0	50.0	0.0	-	-	3.5	-	-	-	-	-	-	-
97.0	60.0	0.0	-	-	0.0	-	-	-	-	-	-	-
97.0	65.0	0.0	-	-	0.0	-	-	-	-	-	-	-
97.0	70.0	3.0	-	-	-	-	-	-	-	-	-	-
97.0	80.0	10.1	-	-	-	-	-	-	-	-	-	-
97.0	85.0	-	-	-	3.6	-	-	-	-	-	-	-
97.0	90.0	15.3	-	-	-	-	-	-	-	-	-	-
100.0	35.0	0.0	-	-	3.2	-	-	-	-	-	-	-
100.0	40.0	0.0	-	-	3.2	-	-	-	-	-	-	-
100.0	45.0	0.0	-	-	4.5	-	-	-	-	-	-	-
100.0	50.0	0.0	-	-	15.1	-	-	-	-	-	-	-
100.0	55.0	6.1	-	-	0.0	-	-	-	-	-	-	-
100.0	65.0	0.0	-	-	0.0	-	-	-	-	-	-	-
100.0	70.0	0.0	-	-	0.0	-	-	-	-	-	-	-
100.0	80.0	2.9	-	-	16.7	-	-	-	-	-	-	-
100.0	85.0	-	-	-	10.4	-	-	-	-	-	-	-
100.0	90.0	9.1	-	-	59.6	-	-	-	-	-	-	-
100.0	100.0	-	-	-	17.7	-	-	-	-	-	-	-
100.0	120.0	-	-	-	3.0	-	-	-	-	-	-	-
103.0	29.0	0.0	-	-	0.0	-	-	-	-	-	-	-
103.0	45.0	0.0	-	-	3.0	-	-	-	-	-	-	-

TABLE 4. (cont.)

*Symbolophorus californiensis* (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
103.0	50.0	0.0	-	-	9.9	-	0.0	-	0.0	-	-	-
103.0	55.0	0.0	-	-	3.3	-	0.0	-	0.0	-	-	-
103.0	65.0	2.8	-	-	0.0	-	0.0	-	0.0	-	-	-
103.0	70.0	3.0	-	-	0.0	-	3.4	-	0.0	-	-	-
103.0	75.0	-	-	-	9.6	-	-	-	-	-	-	-
103.0	80.0	-	0.0	-	8.9	-	0.0	-	-	-	-	-
103.0	90.0	-	6.1	-	6.3	-	13.8	-	0.0	-	-	-
107.0	45.0	-	3.0	-	0.0	-	0.0	-	0.0	-	-	-
107.0	60.0	-	3.0	-	3.2	-	0.0	-	0.0	-	-	-
107.0	65.0	-	0.0	-	3.1	-	0.0	-	0.0	-	-	-
107.0	70.0	-	3.2	-	0.0	-	3.1	-	0.0	-	-	-
107.0	75.0	-	-	-	3.1	-	-	-	-	-	-	-
107.0	80.0	-	11.6	-	3.0	-	3.2	-	-	-	-	-
107.0	90.0	-	0.0	-	6.1	-	0.0	-	-	-	-	-
110.0	75.0	-	-	-	3.2	-	-	-	-	-	-	-
110.0	80.0	-	5.1	-	0.0	-	0.0	-	-	-	-	-
113.0	60.0	-	0.0	-	0.0	-	3.4	-	0.0	-	-	-
117.0	90.0	-	2.9	-	0.0	-	0.0	-	-	-	-	-

*Tarletonbeania crenularis*

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
60.0	49.5	0.9	-	-	-	-	-	-	-	-	-	-
60.0	51.0	3.2	-	-	-	-	-	-	0.0	-	-	-
60.0	52.0	6.8	-	-	-	-	-	-	0.0	-	-	-
60.0	55.0	9.3	-	-	-	-	-	-	2.6	-	-	-
60.0	60.0	15.9	-	-	-	-	-	-	14.2	-	-	-
60.0	70.0	13.0	-	-	-	-	-	-	14.1	-	-	-
60.0	80.0	27.9	-	-	-	-	-	-	6.1	-	-	-
60.0	90.0	6.7	-	-	-	-	-	-	8.3	-	-	-
60.0	100.0	0.0	-	-	-	-	-	-	2.6	-	-	-
60.0	120.0	4.9	-	-	-	-	-	-	2.7	-	-	-
63.0	51.0	10.5	-	-	-	-	-	-	0.0	-	-	-
63.0	52.0	34.8	-	-	-	-	-	-	0.0	-	-	-
63.0	55.0	34.9	-	-	-	-	-	-	0.0	-	-	-
63.0	60.0	22.8	-	-	-	-	-	-	0.0	-	-	-
63.0	65.0	-	-	-	-	-	-	-	-	-	-	-
63.0	80.0	-	-	-	-	-	-	-	-	-	-	-
67.0	47.3	0.7	-	-	-	-	-	-	-	-	-	-
67.0	48.0	0.0	-	-	-	-	-	-	-	-	-	-
67.0	49.0	3.2	-	-	-	-	-	-	-	-	-	-
67.0	50.0	4.7	-	-	-	-	-	-	-	-	-	-
67.0	55.0	6.6	-	-	-	-	-	-	-	-	-	-
67.0	60.0	18.1	-	-	-	-	-	-	-	-	-	-
67.0	65.0	-	-	-	-	-	-	-	-	-	-	-

TABLE 4. (cont.)

*Tarletonbeania crenularis* (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
67.0	70.0	-	-	-	17.0	-	-	-	-	-	-	-
67.0	80.0	-	-	-	21.2	-	-	-	-	-	-	-
70.0	50.0	29.3	-	-	0.0	-	-	-	-	-	-	-
70.0	51.0	9.2	-	-	11.0	-	-	-	-	-	-	-
70.0	52.0	9.5	-	-	23.2	-	-	-	-	-	-	-
70.0	53.0	3.0	-	-	18.9	-	-	-	-	-	-	-
70.0	55.0	11.6	-	-	-	-	-	-	-	-	-	-
70.0	60.0	28.6	-	-	0.0	-	-	-	-	-	-	-
70.0	65.0	-	-	-	8.1	-	-	-	-	-	-	-
70.0	70.0	59.6	-	-	8.7	-	-	-	-	-	-	-
70.0	80.0	0.0	-	-	2.7	-	-	-	-	-	-	-
70.0	50.0	7.0	-	-	0.0	-	-	-	-	-	-	-
73.0	51.0	6.2	-	-	0.0	-	-	-	-	-	-	-
73.0	53.0	5.5	-	-	12.0	-	-	-	-	-	-	-
73.0	60.0	75.3	-	-	11.3	-	-	-	-	-	-	-
73.0	65.0	-	-	-	8.6	-	-	-	-	-	-	-
73.0	70.0	-	-	-	2.8	-	-	-	-	-	-	-
73.0	80.0	-	-	-	7.8	-	-	-	-	-	-	-
77.0	49.0	-	-	-	0.0	-	-	-	-	-	-	-
77.0	55.0	8.6	-	-	7.2	-	-	-	-	-	-	-
77.0	57.0	-	-	-	7.4	-	-	-	-	-	-	-
77.0	60.0	-	-	-	6.3	-	-	-	-	-	-	-
77.0	65.0	-	-	-	2.5	-	-	-	-	-	-	-
77.0	70.0	-	-	-	4.6	-	-	-	-	-	-	-
77.0	80.0	-	-	-	17.0	-	-	-	-	-	-	-
80.0	50.2	0.0	-	-	2.3	-	-	-	-	-	-	-
80.0	51.0	0.0	-	-	0.0	-	-	-	-	-	-	-
80.0	55.0	0.0	-	-	3.2	-	-	-	-	-	-	-
80.0	60.0	7.7	-	-	3.4	-	-	-	-	-	-	-
80.0	65.0	25.0	-	-	3.4	-	-	-	-	-	-	-
80.0	70.0	0.0	-	-	13.6	-	-	-	-	-	-	-
80.0	75.0	-	-	-	-	-	-	-	-	-	-	-
80.0	80.0	-	-	-	6.4	-	-	-	-	-	-	-
80.0	120.0	0.0	-	-	20.6	-	-	-	-	-	-	-
83.0	55.0	14.5	-	-	2.8	-	-	-	-	-	-	-
83.0	43.0	0.0	-	-	0.0	-	-	-	-	-	-	-
83.0	60.0	6.1	-	-	3.0	-	-	-	-	-	-	-
83.0	65.0	5.7	-	-	0.0	-	-	-	-	-	-	-
83.0	70.0	6.4	-	-	6.3	-	-	-	-	-	-	-
83.0	80.0	0.0	-	-	2.9	-	-	-	-	-	-	-
83.0	90.0	9.1	-	-	-	-	-	-	-	-	-	-
83.0	55.0	-	-	-	0.0	-	-	-	-	-	-	-
87.0	55.0	-	-	-	11.9	-	-	-	-	-	-	-
87.0	60.0	0.0	-	-	0.0	-	-	-	-	-	-	-
87.0	65.0	0.0	-	-	8.0	-	-	-	-	-	-	-
87.0	70.0	0.0	-	-	3.2	-	-	-	-	-	-	-
87.0	75.0	-	-	-	0.0	-	-	-	-	-	-	-

TABLE 4. (cont.)

*Tarletonbeania crenularis* (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
87.0	80.0	3.0	-	-	0.0	-	3.3	-	-	0.0	-	-
90.0	53.0	3.2	-	-	0.0	-	-	0.0	-	0.0	-	-
90.0	60.0	3.0	-	-	0.0	-	-	0.0	-	0.0	-	-
90.0	65.0	3.2	-	-	0.0	-	-	2.7	-	0.0	-	-
90.0	80.0	8.0	-	-	0.0	-	-	0.0	-	0.0	-	-
90.0	90.0	3.0	-	-	0.0	-	-	2.3	-	0.0	-	-
90.0	100.0	0.0	-	-	3.1	-	-	0.0	-	0.0	-	-
93.0	30.0	0.0	-	-	2.9	-	-	0.0	-	0.0	-	-
93.0	50.0	0.0	-	-	-	0.0	-	6.6	-	0.0	-	-
93.0	60.0	0.0	-	-	-	0.0	-	3.3	-	0.0	-	-
93.0	90.0	3.2	-	-	-	0.0	-	0.0	-	0.0	-	-
97.0	45.0	3.0	-	-	0.0	-	-	0.0	-	0.0	-	-
97.0	50.0	6.3	-	-	0.0	-	-	0.0	-	0.0	-	-
100.0	45.0	0.0	-	-	-	2.3	-	-	0.0	-	0.0	-

*Synodus spp.*

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
1113.0	28.5	-	0.0	-	0.0	-	0.0	-	-	7.8	-	-
1113.0	29.0	-	0.0	-	0.0	-	0.0	-	-	17.8	-	-
1113.0	30.0	-	0.0	-	0.0	-	0.0	-	0.0	5.4	-	-
1115.0	35.0	-	0.0	-	-	0.0	-	-	0.0	3.1	-	-
1117.0	24.8	-	0.0	-	0.0	-	-	-	-	5.8	-	-
1117.0	25.0	-	0.0	-	0.0	-	0.0	-	-	7.7	-	-
1117.0	26.0	-	0.0	-	0.0	-	0.0	-	-	86.8	-	-
1117.0	35.0	-	0.0	-	0.0	-	0.0	-	-	23.4	-	-
1118.0	39.0	-	0.0	-	0.0	-	0.0	-	-	54.4	-	-
1119.0	33.0	-	0.0	-	0.0	-	0.0	-	-	8.6	-	-
120.0	23.0	-	0.0	-	0.0	-	0.0	-	-	35.7	-	-
120.0	24.0	-	0.0	-	0.0	-	0.0	-	-	69.0	-	-
120.0	25.0	-	0.0	-	0.0	-	0.0	-	-	9.3	-	-
120.0	30.0	-	0.0	-	0.0	-	0.0	-	-	15.1	-	-
120.0	35.0	-	0.0	-	0.0	-	0.0	-	-	21.9	-	-
120.0	40.0	-	0.0	-	0.0	-	0.0	-	-	13.7	-	-
127.0	32.6	-	0.0	-	0.0	-	0.0	-	-	0.9	-	-
127.0	34.0	-	0.0	-	0.0	-	0.0	-	-	3.0	-	-
130.0	30.0	-	0.0	-	0.0	-	0.0	-	-	2.6	-	-
130.0	35.0	-	3.3	-	0.0	-	0.0	-	-	3.0	-	-
133.0	21.0	-	0.0	-	0.0	-	0.0	-	-	11.4	-	-
133.0	23.0	-	0.0	-	0.0	-	0.0	-	-	2.5	-	-
133.0	25.0	-	0.0	-	0.0	-	0.0	-	-	5.9	-	-
133.0	30.0	-	0.0	-	0.0	-	0.0	-	-	14.3	-	-
133.0	35.0	-	2.9	-	0.0	-	0.0	-	-	5.9	-	-
137.0	20.0	-	0.0	-	0.0	-	0.0	-	-	0.4	-	-
137.0	20.0	-	0.6	-	0.0	-	0.0	-	-	1.4	-	-

TABLE 4. (cont.)

*Synodus spp.* (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
137.0	22.0	-	3.9	-	0.0	-	-	-	0.0	12.2	-	-
137.0	23.0	-	5.6	-	0.0	-	-	-	7.6	13.8	-	-
137.0	30.0	-	0.0	-	0.0	-	-	-	0.0	36.6	-	-
137.0	35.0	-	0.0	-	0.0	-	-	-	0.0	3.1	-	-
137.0	40.0	-	3.1	-	0.0	-	-	-	0.0	0.0	-	-
140.0	28.8	-	0.0	-	-	-	-	-	1.0	-	-	-
140.0	29.0	-	0.0	-	-	-	-	-	23.5	-	-	-

*Merluccius productus*

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
60.0	52.0	0.0	-	-	-	-	-	-	-	2.5	-	-
63.0	55.0	3.2	-	-	-	-	-	-	-	0.0	-	-
67.0	55.0	3.3	-	-	-	-	-	-	-	0.0	-	-
67.0	60.0	15.1	-	-	-	-	-	-	-	0.0	-	-
67.0	65.0	-	-	-	-	-	-	-	-	-	-	-
67.0	70.0	-	-	-	-	-	-	-	-	-	-	-
70.0	50.0	11.7	-	-	-	-	-	-	-	-	-	-
70.0	51.0	12.3	-	-	-	-	-	-	-	-	-	-
70.0	52.0	66.2	-	-	-	-	-	-	-	-	-	-
70.0	53.0	38.4	-	-	-	-	-	-	-	-	-	-
70.0	60.0	0.0	-	-	-	-	-	-	-	-	-	-
70.0	65.0	-	-	-	-	-	-	-	-	-	-	-
70.0	70.0	2.7	-	-	-	-	-	-	-	-	-	-
73.0	49.4	0.0	-	-	-	-	-	-	-	-	-	-
73.0	50.0	0.0	-	-	-	-	-	-	-	-	-	-
73.0	51.0	30.9	-	-	-	-	-	-	-	-	-	-
73.0	53.0	32.8	-	-	-	-	-	-	-	-	-	-
73.0	60.0	2.8	-	-	-	-	-	-	-	-	-	-
73.0	80.0	-	-	-	-	-	-	-	-	-	-	-
77.0	47.6	0.8	-	-	-	-	-	-	-	-	-	-
77.0	51.0	80.2	-	-	-	-	-	-	-	-	-	-
77.0	55.0	2.9	-	-	-	-	-	-	-	-	-	-
77.0	57.0	0.0	-	-	-	-	-	-	-	-	-	-
77.0	60.0	-	-	-	-	-	-	-	-	-	-	-
77.0	65.0	-	-	-	-	-	-	-	-	-	-	-
77.0	70.0	-	-	-	-	-	-	-	-	-	-	-
80.0	50.2	25.9	-	-	-	-	-	-	-	-	-	-
80.0	51.0	543.5	-	-	-	-	-	-	-	-	-	-
80.0	52.0	243.5	-	-	-	-	-	-	-	-	-	-
80.0	55.0	307.9	-	-	-	-	-	-	-	-	-	-
80.0	60.0	25.7	-	-	-	-	-	-	-	-	-	-
80.0	65.0	0.0	-	-	-	-	-	-	-	-	-	-
80.0	70.0	0.0	-	-	-	-	-	-	-	-	-	-

TABLE 4. (cont.)

*Merluccius productus* (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
80.0	80.0	0.0	-	-	77.3	-	-	0.0	-	0.0	-	-
82.0	47.0	105.8	-	-	31.0	-	-	0.0	-	0.0	-	-
83.0	40.0	119.1	-	-	0.0	-	-	0.0	-	0.0	-	-
83.0	43.0	159.8	-	-	30.8	-	-	6.3	-	2.5	-	-
83.0	51.0	111.8	-	-	2.6	-	-	0.0	-	0.0	-	-
83.0	55.0	72.5	-	-	15.8	-	-	0.0	-	0.0	-	-
83.0	60.0	30.4	-	-	0.0	-	-	3.5	-	0.0	-	-
83.0	65.0	37.3	-	-	2.9	-	-	6.6	-	0.0	-	-
83.0	70.0	6.4	-	-	22.1	-	-	3.4	-	0.0	-	-
83.0	80.0	3.5	-	-	109.1	-	-	0.0	-	0.0	-	-
87.0	32.5	9.5	-	-	0.0	-	-	2.7	-	0.0	-	-
87.0	33.0	29.9	-	-	2.6	-	-	0.0	-	0.0	-	-
87.0	34.0	251.0	-	-	2.5	-	-	0.0	-	0.0	-	-
87.0	35.0	461.6	-	-	37.5	-	-	0.0	-	0.0	-	-
87.0	40.0	946.2	-	-	33.1	-	-	0.0	-	0.0	-	-
87.0	45.0	364.1	-	-	33.4	-	-	0.0	-	0.0	-	-
87.0	50.0	82.7	-	-	6.9	-	-	0.0	-	0.0	-	-
87.0	55.0	426.1	-	-	8.4	-	-	0.0	-	0.0	-	-
87.0	60.0	0.0	-	-	33.5	-	-	0.0	-	0.0	-	-
87.0	65.0	0.0	-	-	32.7	-	-	0.0	-	0.0	-	-
87.0	70.0	0.0	-	-	44.9	-	-	0.0	-	0.0	-	-
87.0	80.0	0.0	-	-	24.1	-	-	0.0	-	0.0	-	-
87.0	90.0	0.0	-	-	11.3	-	-	0.0	-	0.0	-	-
87.0	100.0	-	-	-	6.2	-	-	0.0	-	0.0	-	-
90.0	127.5	4.6	-	-	3.9	-	-	0.0	-	0.0	-	-
90.0	28.0	106.0	-	-	12.2	-	-	0.0	-	0.0	-	-
90.0	32.0	229.5	-	-	6.1	-	-	0.0	-	0.0	-	-
90.0	37.0	418.6	-	-	24.7	-	-	0.0	-	0.0	-	-
90.0	45.0	91.4	-	-	62.6	-	-	0.0	-	0.0	-	-
90.0	53.0	16.1	-	-	36.5	-	-	0.0	-	0.0	-	-
90.0	60.0	9.0	-	-	30.1	-	-	0.0	-	0.0	-	-
90.0	65.0	0.0	-	-	29.4	-	-	0.0	-	0.0	-	-
90.0	70.0	0.0	-	-	68.4	-	-	0.0	-	0.0	-	-
90.0	80.0	0.0	-	-	9.7	-	-	0.0	-	0.0	-	-
90.0	90.0	0.0	-	-	9.6	-	-	0.0	-	0.0	-	-
90.0	100.0	0.0	-	-	31.2	-	-	0.0	-	0.0	-	-
93.0	26.5	0.0	-	-	-	-	-	1.3	-	0.0	-	-
93.0	28.0	52.5	-	-	-	-	-	0.0	-	0.0	-	-
93.0	30.0	513.8	-	-	11.5	-	-	0.0	-	0.0	-	-
93.0	35.0	54.0	-	-	5.8	-	-	0.0	-	0.0	-	-
93.0	45.0	294.9	-	-	-	-	-	3.3	-	0.0	-	-
93.0	50.0	46.6	-	-	0.0	-	-	0.0	-	0.0	-	-
93.0	55.0	30.7	-	-	12.5	-	-	0.0	-	0.0	-	-
93.0	60.0	6.1	-	-	13.5	-	-	0.0	-	0.0	-	-
93.0	65.0	0.0	-	-	17.9	-	-	0.0	-	0.0	-	-
93.0	70.0	0.0	-	-	66.6	-	-	0.0	-	0.0	-	-

TABLE 4. (cont.)

*Merluccius productus* (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
93.0	80.0	0.0	-	-	3.2	-	0.0	-	-	0.0	-	-
93.0	100.0	-	-	-	8.5	-	-	-	-	-	-	-
96.8	28.7	4.7	0.0	-	0.0	-	0.0	-	-	0.0	-	-
97.0	29.0	3.7	14.5	-	-	-	0.0	-	-	0.0	-	-
97.0	30.0	38.9	0.0	-	-	-	0.0	-	-	0.0	-	-
97.0	32.0	81.1	27.9	-	-	-	3.4	-	-	0.0	-	-
97.0	35.0	18.4	34.7	-	-	-	0.0	-	-	0.0	-	-
97.0	40.0	18.9	9.6	-	-	-	0.0	-	-	0.0	-	-
97.0	45.0	106.4	10.4	-	-	-	0.0	-	-	0.0	-	-
97.0	50.0	63.0	35.4	-	-	-	0.0	-	-	0.0	-	-
97.0	55.0	3.0	-	-	-	-	0.0	-	-	0.0	-	-
97.0	60.0	17.8	-	-	-	-	0.0	-	-	0.0	-	-
97.0	65.0	110.5	20.7	-	-	-	0.0	-	-	0.0	-	-
97.0	70.0	14.8	-	-	-	-	0.0	-	-	0.0	-	-
97.0	75.0	-	-	-	-	-	34.2	-	-	-	-	-
97.0	85.0	-	-	-	-	-	3.6	-	-	-	-	-
97.0	95.0	-	-	-	-	-	54.5	-	-	-	-	-
97.0	100.0	-	-	-	-	-	7.0	-	-	-	-	-
100.0	29.0	78.8	-	-	-	-	11.6	-	-	0.0	-	-
100.0	30.0	22.5	-	-	-	-	19.3	-	-	0.0	-	-
100.0	35.0	8.8	6.3	-	-	-	0.0	-	-	0.0	-	-
100.0	40.0	11.8	6.4	-	-	-	0.0	-	-	0.0	-	-
100.0	45.0	90.2	13.6	-	-	-	0.0	-	-	0.0	-	-
100.0	50.0	89.3	21.1	-	-	-	0.0	-	-	0.0	-	-
100.0	55.0	69.7	24.2	-	-	-	0.0	-	-	0.0	-	-
100.0	60.0	44.7	6.7	-	-	-	0.0	-	-	0.0	-	-
100.0	65.0	38.7	41.5	-	-	-	0.0	-	-	0.0	-	-
100.0	70.0	46.2	83.5	-	-	-	0.0	-	-	0.0	-	-
100.0	75.0	-	43.8	-	-	-	0.0	-	-	0.0	-	-
100.0	80.0	0.0	-	-	-	-	0.0	-	-	0.0	-	-
100.0	90.0	0.0	-	-	-	-	0.0	-	-	0.0	-	-
103.0	28.5	1.3	-	-	-	-	3.3	-	-	0.0	-	-
103.0	30.0	0.0	-	-	-	-	3.6	-	-	0.0	-	-
103.0	35.0	47.0	-	-	-	-	18.4	-	-	0.0	-	-
103.0	40.0	20.4	-	-	-	-	9.3	-	-	0.0	-	-
103.0	45.0	0.0	-	-	-	-	47.0	-	-	0.0	-	-
103.0	55.0	11.5	-	-	-	-	42.8	-	-	0.0	-	-
103.0	60.0	3.0	-	-	-	-	9.1	-	-	0.0	-	-
103.0	65.0	48.1	-	-	-	-	0.0	-	-	0.0	-	-
103.0	70.0	18.1	-	-	-	-	0.0	-	-	0.0	-	-
107.0	30.0	-	-	-	-	-	33.0	-	-	0.0	-	-
107.0	31.0	-	-	-	-	-	150.5	-	-	8.5	-	-
107.0	32.0	-	-	-	-	-	908.5	-	-	49.0	-	-
107.0	35.0	-	-	-	-	-	35.2	-	-	22.1	-	-
107.0	40.0	-	-	-	-	-	52.9	-	-	12.0	-	-
107.0	60.0	-	-	-	-	-	30.3	-	-	0.0	-	-

TABLE 4. (cont.)

*Merluccius productus* (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
110.0	32.0	-	13.4	-	0.0	-	-	-	0.0	0.0	-	-
110.0	35.0	-	73.6	-	0.0	-	-	-	0.0	0.0	-	-
110.0	40.0	-	617.7	-	2.8	-	-	-	0.0	0.0	-	-
110.0	55.0	-	0.0	-	3.0	-	-	-	0.0	0.0	-	-
113.0	30.0	-	0.0	-	2.8	-	-	-	0.0	0.0	-	-
113.0	45.0	-	18.5	-	0.0	-	-	-	0.0	0.0	-	-
113.0	50.0	-	3.0	-	0.0	-	-	-	0.0	0.0	-	-
113.0	70.0	-	0.0	-	6.2	-	-	-	0.0	0.0	-	-
113.0	75.0	-	-	-	6.6	-	-	-	0.0	0.0	-	-
115.0	35.0	-	8.6	-	-	-	-	-	2.4	-	-	-
117.0	25.0	-	44.4	-	0.0	-	-	-	0.0	0.0	-	-
117.0	26.0	-	61.6	-	0.0	-	-	-	0.0	0.0	-	-
117.0	30.0	-	471.1	-	0.0	-	-	-	0.0	0.0	-	-
117.0	40.0	-	6.1	-	0.0	-	-	-	0.0	0.0	-	-
117.0	45.0	-	32.6	-	0.0	-	-	-	0.0	0.0	-	-
117.0	50.0	-	16.0	-	0.0	-	-	-	0.0	0.0	-	-
117.0	65.0	-	0.0	-	2.8	-	-	-	0.0	0.0	-	-
118.0	39.0	-	3.1	-	6.2	-	-	-	0.0	0.0	-	-
119.0	33.0	-	23.9	-	0.0	-	-	-	0.0	0.0	-	-
120.0	22.5	-	0.0	-	2.0	-	-	-	0.0	0.0	-	-
120.0	25.0	-	0.0	-	0.0	-	-	-	0.0	0.0	-	-
120.0	30.0	-	11.6	-	0.0	-	-	-	2.7	-	-	-
120.0	35.0	-	3.0	-	0.0	-	-	-	0.0	0.0	-	-
120.0	45.0	-	2.9	-	3.1	-	-	-	0.0	0.0	-	-
120.0	50.0	-	12.3	-	0.0	-	-	-	0.0	0.0	-	-
120.0	70.0	-	0.0	-	3.3	-	-	-	0.0	0.0	-	-
123.0	36.0	-	0.0	-	94.0	-	-	-	0.0	0.0	-	-
123.0	37.0	-	14.3	-	0.0	-	-	-	0.0	0.0	-	-
123.0	42.0	-	333.8	-	5.7	-	-	-	0.0	0.0	-	-
123.0	45.0	-	17.5	-	14.6	-	-	-	0.0	0.0	-	-
123.0	65.0	-	2.9	-	0.0	-	-	-	0.0	0.0	-	-
123.0	70.0	-	2.8	-	0.0	-	-	-	0.0	0.0	-	-
123.5	35.6	-	2.2	-	91.5	-	-	-	0.0	0.0	-	-
127.0	32.6	-	22.4	-	0.0	-	-	-	0.0	0.0	-	-
127.0	33.0	-	168.1	-	21.5	-	-	-	0.0	0.0	-	-
127.0	34.0	-	26.3	-	12.2	-	-	-	0.0	0.0	-	-
127.0	40.0	-	12.7	-	44.5	-	-	-	0.0	0.0	-	-
127.0	45.0	-	0.0	-	12.4	-	-	-	0.0	0.0	-	-
130.0	25.5	-	0.6	-	0.0	-	-	-	0.0	0.0	-	-
130.0	28.0	-	2.1	-	0.0	-	-	-	0.0	0.0	-	-
130.0	30.0	-	2.4	-	0.0	-	-	-	0.0	0.0	-	-
130.0	35.0	-	116.5	-	28.1	-	-	-	0.0	0.0	-	-
130.0	40.0	-	52.0	-	44.1	-	-	-	0.0	0.0	-	-
130.0	45.0	-	16.7	-	3.0	-	-	-	0.0	0.0	-	-
130.0	50.0	-	708.5	-	0.0	-	-	-	0.0	0.0	-	-
133.0	21.0	-	4.0	-	0.0	-	-	-	0.0	0.0	-	-

TABLE 4. (cont.)

*Merluccius productus* (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
133.0	23.0	-	10.6	-	0.0	-	-	0.0	-	0.0	-	-
133.0	25.0	-	13.5	-	3.0	-	-	0.0	-	0.0	-	-
133.0	30.0	-	16.6	-	9.3	-	-	0.0	-	0.0	-	-
133.0	35.0	-	141.1	-	12.4	-	-	0.0	-	0.0	-	-
133.0	40.0	-	128.1	-	0.0	-	-	0.0	-	0.0	-	-
133.0	45.0	-	224.4	-	3.0	-	-	0.0	-	0.0	-	-
133.0	50.0	-	203.8	-	3.0	-	-	0.0	-	0.0	-	-
133.0	55.0	-	8.2	-	0.0	-	-	0.0	-	0.0	-	-
137.0	23.0	-	1.9	-	0.0	-	-	0.0	-	0.0	-	-
137.0	30.0	-	86.2	-	6.2	-	-	0.0	-	0.0	-	-
137.0	35.0	-	2428.9	-	3.0	-	-	0.0	-	0.0	-	-
137.0	40.0	-	40.0	-	3.0	-	-	0.0	-	0.0	-	-
137.0	45.0	-	0.0	-	3.0	-	-	0.0	-	0.0	-	-
137.0	50.0	-	6.6	-	0.0	-	-	0.0	-	0.0	-	-
140.0	28.8	-	8.2	-	-	-	-	-	-	-	-	-
140.0	29.0	-	7.0	-	-	-	-	-	-	-	-	-
140.0	30.0	-	124.7	-	-	-	-	-	-	-	-	-
140.0	35.0	-	751.4	-	-	-	-	-	-	-	-	-
140.0	40.0	-	663.8	-	-	-	-	-	-	-	-	-
140.0	45.0	-	6.1	-	-	-	-	-	-	-	-	-
140.0	50.0	-	3.0	-	-	-	-	-	-	-	-	-

*Physiculus* spp.

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
80.0	200.0	0.0	-	-	-	-	-	2.8	-	-	-	-
133.0	45.0	-	2.9	-	0.0	-	-	0.0	-	0.0	-	-
133.0	50.0	-	2.8	-	0.0	-	-	0.0	-	0.0	-	-

*Macrouridae*

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
100.0	45.0	3.1	-	0.0	-	-	-	0.0	-	0.0	-	-
133.0	35.0	-	0.0	-	3.1	-	-	0.0	-	0.0	-	-
137.0	35.0	-	3.1	-	0.0	-	-	0.0	-	0.0	-	-

*Ophidiiformes*

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
67.0	60.0	0.0	-	-	0.0	-	-	3.0	-	-	-	-
73.0	50.0	0.0	-	-	0.0	-	-	2.9	-	-	-	-
73.0	51.0	0.0	-	-	0.0	-	-	6.3	-	-	-	-
93.0	27.0	0.0	-	0.0	-	-	-	2.9	-	0.0	-	-

TABLE 4. (cont.)

## Ophidiiformes (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
97.0	32.0	0.0	-	0.0	-	0.0	-	3.4	-	0.0	-	-
100.0	30.0	0.0	-	0.0	-	0.0	-	2.5	-	0.0	-	-
103.0	35.0	0.0	-	0.0	-	3.1	0.0	-	0.0	-	-	-
110.0	32.0	-	0.0	-	0.0	-	0.0	-	0.0	0.9	-	-
113.0	28.5	-	0.0	-	0.0	-	0.0	-	-	6.7	-	-
113.0	29.0	-	0.0	-	0.0	-	0.0	-	-	7.1	-	-
115.0	35.0	-	0.0	-	0.0	-	0.0	-	0.0	3.1	-	-
117.0	24.8	-	0.0	-	0.0	-	0.0	-	0.0	1.1	-	-
117.0	26.0	-	0.0	-	0.0	-	0.0	-	0.0	52.6	-	-
117.0	30.0	-	0.0	-	0.0	-	0.0	-	0.0	7.8	-	-
117.0	40.0	-	0.0	-	0.0	-	6.1	0.0	0.0	0.0	-	-
118.0	39.0	-	0.0	-	0.0	-	0.0	-	0.0	10.9	-	-
119.0	33.0	-	0.0	-	0.0	-	0.0	-	0.0	28.6	-	-
120.0	23.0	-	0.0	-	0.0	-	0.0	-	0.0	16.9	-	-
120.0	24.0	-	0.0	-	0.0	-	0.0	-	0.0	40.5	-	-
120.0	25.0	-	0.0	-	0.0	-	0.0	-	0.0	21.0	-	-
120.0	30.0	-	0.0	-	0.0	-	0.0	-	0.0	42.1	-	-
120.0	35.0	-	0.0	-	0.0	-	0.0	-	0.0	19.2	-	-
130.0	26.0	-	0.0	-	0.0	-	0.0	-	0.0	2.3	-	-
130.0	28.0	-	0.0	-	0.0	-	0.0	-	0.0	7.1	-	-
130.0	30.0	-	0.0	-	0.0	-	0.0	-	0.0	2.6	-	-
130.0	30.0	-	0.0	-	0.0	-	0.0	-	0.0	4.6	-	-
133.0	21.0	-	0.0	-	0.0	-	0.0	-	0.0	2.5	-	-
133.0	23.0	-	0.0	-	0.0	-	0.0	-	0.0	20.0	-	-
133.0	30.0	-	0.0	-	0.0	-	0.0	-	0.0	3.0	-	-
133.0	35.0	-	0.0	-	0.0	-	0.0	-	0.0	4.3	-	-
137.0	21.0	-	0.0	-	0.0	-	0.0	-	0.0	15.0	-	-
137.0	22.0	-	0.0	-	0.0	-	0.0	-	1.9	16.6	-	-
137.0	23.0	-	0.0	-	0.0	-	0.0	-	0.0	18.3	-	-
137.0	30.0	-	0.0	-	0.0	-	0.0	-	0.0	17.6	-	-
140.0	29.0	-	0.0	-	0.0	-	-	-	11.4	-	-	-
140.0	30.0	-	0.0	-	0.0	-	-	-	-	-	-	-

*Brosomophycis marginata*

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
60.0	55.0	0.0	-	-	-	-	-	-	-	2.6	-	-
67.0	50.0	0.0	-	-	-	-	-	-	0.0	-	-	-
70.0	52.0	0.0	-	-	-	-	-	-	-	-	-	-
83.0	43.0	0.0	-	-	-	-	-	-	0.0	-	0.0	-
87.0	50.0	0.0	-	-	-	-	-	-	0.0	-	0.0	-
103.0	29.0	0.0	-	0.0	-	-	-	-	1.7	-	0.0	-
107.0	30.0	-	0.0	-	0.0	-	-	-	1.1	-	0.0	-

TABLE 4. (cont.)

*Chilara taylori*

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
63.0	49.7	0.6	-	-	0.0	-	-	-	-	-	-	-
63.0	50.0	0.8	-	-	0.0	-	-	0.0	-	-	-	-
63.0	52.0	2.9	-	-	0.0	-	-	0.0	-	-	-	-
63.0	55.0	3.2	-	-	0.0	-	-	0.0	-	-	-	-
70.0	52.0	3.2	-	-	0.0	-	-	0.0	-	-	-	-
80.0	70.0	0.0	-	-	0.0	-	-	0.0	-	-	-	-
83.0	40.0	0.0	-	-	0.0	-	-	0.0	-	-	-	-
83.0	43.0	0.0	-	-	0.0	-	-	0.0	-	-	-	-
83.0	60.0	0.0	-	-	0.0	-	-	0.0	-	-	-	-
87.0	32.5	0.0	-	-	0.0	-	-	0.0	-	-	-	-
87.0	50.0	0.0	-	-	0.0	-	-	0.0	-	-	-	-
87.0	50.0	0.0	-	-	0.0	-	-	0.0	-	-	-	-
93.0	30.0	0.0	-	0.0	-	-	-	-	-	-	-	-
110.0	35.0	-	0.0	-	0.0	-	-	-	-	-	-	-
110.0	65.0	-	2.7	-	0.0	-	-	-	-	-	-	-
113.0	29.0	-	0.0	-	0.0	-	-	-	-	-	-	-
113.0	30.0	-	0.0	-	0.0	-	-	-	-	-	-	-
117.0	25.0	-	0.0	-	0.0	-	-	-	-	-	-	-
117.0	26.0	-	0.0	-	0.0	-	-	-	-	-	-	-
117.0	30.0	-	0.0	-	0.0	-	-	-	-	-	-	-
117.0	35.0	-	0.0	-	0.0	-	-	-	-	-	-	-
117.0	50.0	-	0.0	-	0.0	-	-	-	-	-	-	-
119.0	33.0	-	0.0	-	0.0	-	-	-	-	-	-	-
120.0	23.0	-	0.0	-	0.0	-	-	-	-	-	-	-
120.0	35.0	-	0.0	-	0.0	-	-	-	-	-	-	-
123.5	35.6	-	0.0	-	0.0	-	-	-	-	-	-	-
130.0	28.0	-	0.0	-	0.0	-	-	-	-	-	-	-
130.0	35.0	-	0.0	-	0.0	-	-	-	-	-	-	-
133.0	50.0	-	0.0	-	0.0	-	-	-	-	-	-	-
137.0	80.0	-	0.0	-	-	-	-	-	-	-	-	-

*Ophidion scripttiae*

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
83.0	39.0	0.0	-	-	-	0.0	-	-	-	-	3.9	-
87.0	32.5	0.0	-	-	0.0	-	-	0.0	-	-	31.5	-
87.0	34.0	0.0	-	-	0.0	-	-	0.0	-	-	2.6	-
87.0	35.0	0.0	-	-	0.0	-	-	0.0	-	-	3.0	-
96.8	28.7	0.0	-	-	0.0	-	-	0.0	-	-	1.5	-
97.0	29.0	0.0	-	-	0.0	-	-	0.0	-	-	12.2	-
97.0	30.0	0.0	-	-	0.0	-	-	0.0	-	-	2.3	-
100.0	30.0	0.0	-	-	0.0	-	-	2.5	-	-	0.0	-
103.0	30.0	0.0	-	-	0.0	-	-	7.9	-	-	0.0	-
110.0	32.0	-	0.0	-	0.0	-	-	-	-	-	0.9	-
115.0	35.0	-	0.0	-	0.0	-	-	-	-	-	3.1	-
117.0	26.0	-	0.0	-	0.0	-	-	-	-	-	28.9	-

TABLE 4. (cont.)

*Ophidion scriptpsae* (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
117.0	30.0	-	0.0	-	0.0	-	0.0	-	2.6	-	-	-
117.0	35.0	-	0.0	-	0.0	-	0.0	-	5.8	-	-	-
117.0	50.0	-	0.0	-	2.9	-	0.0	-	0.0	-	-	-
118.0	39.0	-	0.0	-	0.0	-	0.0	-	19.0	-	-	-
119.0	33.0	-	0.0	-	0.0	-	-	106.3	22.9	-	-	-
120.0	24.0	-	0.0	-	0.0	-	0.0	-	21.4	-	-	-
120.0	25.0	-	0.0	-	0.0	-	0.0	-	11.6	-	-	-
120.0	30.0	-	0.0	-	0.0	-	0.0	-	42.1	-	-	-
120.0	35.0	-	0.0	-	0.0	-	0.0	-	32.9	-	-	-
120.0	40.0	-	0.0	-	0.0	-	2.6	-	2.7	-	-	-
120.0	45.0	-	0.0	-	0.0	-	-	5.7	0.0	-	-	-
123.5	35.6	-	0.0	-	0.0	-	0.0	-	1.4	-	-	-
127.0	33.0	-	0.0	-	0.0	-	0.0	-	2.6	-	-	-
130.0	28.0	-	0.0	-	0.0	-	-	0.0	7.1	-	-	-
130.0	30.0	-	0.0	-	0.0	-	-	2.0	5.2	-	-	-
130.0	35.0	-	0.0	-	0.0	-	-	0.0	6.1	-	-	-
133.0	21.0	-	0.0	-	0.0	-	0.0	-	18.3	-	-	-
133.0	23.0	-	0.0	-	0.0	-	0.0	-	40.0	-	-	-
133.0	25.0	-	0.0	-	0.0	-	0.0	-	26.5	-	-	-
133.0	35.0	-	0.0	-	0.0	-	0.0	-	3.0	-	-	-
137.0	21.0	-	0.0	-	0.0	-	-	0.0	6.5	-	-	-
137.0	22.0	-	0.0	-	0.0	-	-	0.0	2.0	-	-	-
137.0	23.0	-	0.0	-	0.0	-	-	1.9	0.0	-	-	-
137.0	30.0	-	0.0	-	0.0	-	-	0.0	12.2	-	-	-
140.0	29.0	-	0.0	-	-	-	-	2.0	-	-	-	-

## Ceratioidei

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
70.0	200.0	0.0	-	-	-	-	-	2.7	-	-	-	-
90.0	120.0	0.0	-	-	-	-	-	2.7	-	-	-	-
93.0	90.0	0.0	-	0.0	0.0	-	0.0	-	2.8	-	-	-
100.0	45.0	0.0	-	0.0	-	-	0.0	-	3.1	-	-	-
100.0	100.0	-	0.0	-	0.0	-	3.0	-	-	3.3	-	-
107.0	65.0	-	0.0	-	0.0	-	-	0.0	0.0	3.1	-	-
110.0	50.0	-	0.0	-	0.0	-	-	0.0	0.0	3.1	-	-
110.0	55.0	-	0.0	-	0.0	-	-	0.0	0.0	11.9	-	-
110.0	60.0	-	0.0	-	0.0	-	-	0.0	0.0	3.1	-	-
110.0	65.0	-	0.0	-	0.0	-	-	0.0	0.0	3.1	-	-
110.0	120.0	-	0.0	-	-	-	2.7	-	-	3.0	-	-
113.0	50.0	-	0.0	-	0.0	-	0.0	-	-	3.0	-	-
117.0	60.0	-	0.0	-	0.0	-	0.0	-	-	3.0	-	-
120.0	80.0	-	0.0	-	0.0	-	-	2.7	-	-	3.1	-
127.0	45.0	-	0.0	-	0.0	-	0.0	-	-	3.4	-	-
127.0	60.0	-	0.0	-	0.0	-	-	0.0	-	-	-	-

TABLE 4. (cont.)

## Ceratioidei (cont.)

Ceratioidei (cont.)												
STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
130.0	45.0	-	0.0	-	0.0	-	-	-	0.0	3.0	-	-
130.0	55.0	-	0.0	-	0.0	-	-	-	5.5	0.0	-	-
<b>Gobiesocidae</b>												
STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
97.0	30.0	0.0	-	0.0	-	-	-	0.0	-	-	4.5	-
97.0	32.0	0.0	-	0.0	-	-	-	3.4	-	-	0.0	-
103.0	29.0	0.0	-	0.0	-	-	-	0.0	-	-	5.1	-
110.0	32.0	-	0.0	-	0.0	-	-	-	0.8	0.0	-	-
120.0	40.0	-	0.0	-	0.0	-	-	0.0	-	2.7	-	-
127.0	32.6	-	0.0	-	0.0	-	-	0.0	-	0.9	-	-
133.0	21.0	-	0.0	-	0.0	-	-	0.0	-	13.7	-	-
137.0	21.0	-	0.0	-	0.0	-	-	-	0.0	2.2	-	-
140.0	28.8	-	0.0	-	-	-	-	-	2.4	-	-	-
<b>Exocoetidae</b>												
STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
107.0	65.0	-	0.0	-	0.0	-	-	0.0	-	3.3	-	-
133.0	21.0	-	0.0	-	0.0	-	-	-	0.0	2.3	-	-
<b>Hemiramphidae</b>												
STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
140.0	28.8	-	0.0	-	-	-	-	-	1.0	-	-	-
<b>Cololabis saira</b>												
STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
80.0	200.0	2.8	-	-	-	-	-	0.0	-	-	-	-
87.0	35.0	0.0	-	-	-	0.0	-	3.1	-	-	0.0	-
93.0	26.5	0.0	-	-	-	-	-	1.3	-	-	0.0	-
93.0	60.0	0.0	-	-	-	3.4	-	0.0	-	-	0.0	-
93.0	65.0	0.0	-	-	-	0.0	-	3.3	-	-	0.0	-
100.0	40.0	0.0	-	-	-	0.0	-	-	5.5	-	-	-
100.0	80.0	0.0	-	-	-	3.3	-	0.0	-	-	0.0	-
100.0	100.0	-	3.0	-	-	-	-	0.0	-	-	-	-
130.0	70.0	-	0.0	-	-	6.2	-	0.0	-	-	-	-

TABLE 4. (cont.)

## Atherinidae

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
67.0	47.3	0.0	-	-	5.3	-	-	-	-	-	-	-
67.0	48.0	0.0	-	-	1.3	-	-	0.0	-	-	-	-
96.8	28.7	0.0	-	1.1	-	-	0.0	-	-	-	-	-
97.0	29.0	0.0	-	2.4	-	-	0.0	-	-	-	-	-
107.0	30.0	-	0.0	-	67.3	-	-	0.0	-	-	-	-
107.0	31.0	-	0.0	-	8.5	-	-	0.0	-	-	-	-
113.0	28.5	-	7.0	-	0.0	-	-	0.0	-	-	-	-
117.0	25.0	-	0.0	-	0.0	-	-	2.4	-	-	-	-

## Trachipteridae

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
63.0	55.0	3.2	-	-	0.0	-	-	0.0	-	-	-	-
70.0	70.0	0.0	-	-	2.9	-	-	0.0	-	-	-	-
70.0	90.0	0.0	-	-	-	-	-	2.8	-	-	-	-
70.0	160.0	3.0	-	-	-	-	-	-	-	-	-	-
80.0	60.0	0.0	-	-	0.0	-	-	2.8	-	-	-	-
80.0	65.0	0.0	-	-	3.4	-	-	0.0	-	-	-	-
80.0	70.0	0.0	-	-	0.0	-	-	0.0	-	-	-	-
80.0	80.0	0.0	-	-	0.0	-	-	0.0	-	-	-	-
83.0	80.0	0.0	-	-	2.9	-	-	0.0	-	-	-	-
87.0	60.0	3.0	-	-	0.0	-	-	0.0	-	-	-	-
90.0	65.0	0.0	-	-	2.9	-	-	0.0	-	-	-	-
93.0	65.0	0.0	-	-	0.0	-	-	0.0	-	-	-	-
93.0	100.0	-	-	-	2.8	-	-	-	-	-	-	-
100.0	90.0	0.0	-	-	3.3	-	-	0.0	-	-	-	-
103.0	40.0	0.0	-	-	0.0	-	-	3.3	-	-	-	-
107.0	50.0	-	0.0	-	0.0	-	-	0.0	-	-	-	-
110.0	35.0	-	0.0	-	0.0	-	-	-	-	-	-	-
120.0	45.0	-	0.0	-	0.0	-	-	-	-	-	-	-
123.0	42.0	-	0.0	-	0.0	-	-	-	-	-	-	-

## Melamphaes spp.

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
60.0	140.0	0.0	-	-	-	-	-	-	-	-	-	-
60.0	160.0	3.2	-	-	-	-	-	-	-	-	-	-
60.0	200.0	3.1	-	-	-	-	-	-	-	-	-	-
63.0	80.0	-	3.3	-	-	-	-	-	-	-	-	-
67.0	55.0	3.3	-	-	-	-	-	-	-	-	-	-
70.0	52.0	0.0	-	-	-	-	-	-	-	-	-	-
70.0	53.0	0.0	-	-	-	-	-	-	-	-	-	-
70.0	70.0	0.0	-	-	-	-	-	-	-	-	-	-
70.0	80.0	2.8	-	-	-	-	-	-	-	-	-	-

TABLE 4. (cont.)

*Melamphaes* spp. (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
70.0	90.0	2.9	-	-	-	-	-	0.0	-	-	-	-
70.0	100.0	0.0	-	-	-	-	-	2.8	-	-	-	-
70.0	120.0	0.0	-	-	-	-	-	5.5	-	-	-	-
70.0	160.0	3.0	-	-	-	-	-	0.0	-	-	-	-
70.0	200.0	6.1	-	-	-	-	-	2.8	-	-	-	-
73.0	53.0	0.0	-	0.0	0.0	-	-	0.0	-	-	-	-
73.0	60.0	0.0	-	0.0	0.0	-	-	2.8	-	-	-	-
73.0	65.0	-	-	-	-	-	-	-	-	-	-	-
73.0	70.0	-	-	-	-	-	-	-	-	-	-	-
73.0	80.0	-	-	-	-	-	-	-	-	-	-	-
77.0	47.6	0.3	-	-	-	-	-	-	-	-	-	-
77.0	55.0	0.0	-	-	-	-	-	-	-	-	-	-
77.0	57.0	0.0	-	-	-	-	-	-	-	-	-	-
77.0	80.0	-	-	-	-	-	-	-	-	-	-	-
80.0	60.0	0.0	-	-	-	-	-	-	-	-	-	-
80.0	65.0	0.0	-	-	-	-	-	-	-	-	-	-
80.0	70.0	0.0	-	-	-	-	-	-	-	-	-	-
80.0	80.0	0.0	-	-	-	-	-	-	-	-	-	-
80.0	90.0	0.0	-	-	-	-	-	-	-	-	-	-
80.0	100.0	0.0	-	-	-	-	-	-	-	-	-	-
80.0	140.0	2.7	-	-	-	-	-	-	-	-	-	-
80.0	160.0	8.8	-	-	-	-	-	-	-	-	-	-
80.0	200.0	0.0	-	-	-	-	-	-	-	-	-	-
83.0	80.0	3.5	-	-	-	-	-	-	-	-	-	-
83.0	90.0	3.0	-	-	-	-	-	-	-	-	-	-
87.0	55.0	0.0	-	-	-	-	-	-	-	-	-	-
87.0	60.0	0.0	-	-	-	-	-	-	-	-	-	-
87.0	65.0	0.0	-	-	-	-	-	-	-	-	-	-
87.0	70.0	0.0	-	-	-	-	-	-	-	-	-	-
87.0	75.0	-	-	-	-	-	-	-	-	-	-	-
87.0	80.0	3.0	-	-	-	-	-	-	-	-	-	-
87.0	90.0	0.0	-	-	-	-	-	-	-	-	-	-
87.0	120.0	0.0	-	-	-	-	-	-	-	-	-	-
90.0	200.0	2.8	-	-	-	-	-	-	-	-	-	-
93.0	30.0	0.0	-	0.0	0.0	-	-	0.0	-	-	-	-
93.0	55.0	0.0	-	-	-	-	-	-	-	-	-	-
93.0	65.0	3.2	-	-	-	-	-	-	-	-	-	-
93.0	70.0	2.9	-	-	-	-	-	-	-	-	-	-
93.0	80.0	0.0	-	-	-	-	-	-	-	-	-	-
93.0	90.0	0.0	-	-	-	-	-	-	-	-	-	-
93.0	100.0	-	-	-	-	-	-	-	-	-	-	-
97.0	35.0	3.1	-	-	-	-	-	-	-	-	-	-
97.0	40.0	0.0	-	-	-	-	-	-	-	-	-	-

TABLE 4. (cont.)

*Melamphaes* spp. (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
97.0	45.0	0.0	3.5	—	0.0	0.0	—	—	0.0	0.0	0.0	0.0
97.0	50.0	0.0	0.0	—	0.0	3.3	—	—	0.0	0.0	3.0	0.0
97.0	65.0	0.0	0.0	—	0.0	3.0	—	—	0.0	0.0	0.0	0.0
97.0	70.0	3.0	—	—	0.0	5.8	—	—	0.0	0.0	0.0	0.0
97.0	80.0	3.3	—	—	3.6	—	—	2.8	—	0.0	0.0	0.0
97.0	95.0	—	—	—	9.7	—	—	0.0	—	0.0	0.0	0.0
100.0	40.0	0.0	0.0	—	6.0	—	—	0.0	—	0.0	0.0	0.0
100.0	50.0	0.0	0.0	—	3.3	—	—	0.0	—	0.0	0.0	0.0
100.0	60.0	0.0	0.0	—	3.2	—	—	0.0	—	0.0	0.0	0.0
100.0	65.0	0.0	0.0	—	3.2	—	—	0.0	—	0.0	0.0	0.0
100.0	70.0	0.0	0.0	—	0.0	—	—	0.0	—	0.0	0.0	0.0
100.0	75.0	—	—	—	3.4	—	—	0.0	—	0.0	0.0	0.0
103.0	45.0	0.0	0.0	—	0.0	—	—	0.0	—	0.0	0.0	0.0
103.0	60.0	0.0	0.0	—	0.0	—	—	0.0	—	0.0	0.0	0.0
103.0	70.0	0.0	0.0	—	12.4	—	—	0.0	—	0.0	0.0	0.0
103.0	85.0	—	—	—	2.9	—	—	0.0	—	0.0	0.0	0.0
107.0	55.0	—	—	—	0.0	—	—	0.0	—	0.0	0.0	0.0
107.0	60.0	—	—	—	3.0	—	—	0.0	—	0.0	0.0	0.0
107.0	70.0	0.0	0.0	—	0.0	—	—	0.0	—	0.0	0.0	0.0
107.0	75.0	—	—	—	3.1	—	—	0.0	—	0.0	0.0	0.0
107.0	80.0	0.0	0.0	—	0.0	—	—	0.0	—	0.0	0.0	0.0
110.0	40.0	0.0	0.0	—	2.9	—	—	0.0	—	0.0	0.0	0.0
110.0	45.0	0.0	0.0	—	0.0	—	—	0.0	—	0.0	0.0	0.0
110.0	65.0	0.0	0.0	—	2.7	—	—	0.0	—	0.0	0.0	0.0
110.0	70.0	—	—	—	2.8	—	—	0.0	—	0.0	0.0	0.0
110.0	75.0	—	—	—	0.0	—	—	0.0	—	0.0	0.0	0.0
110.0	80.0	0.0	0.0	—	2.6	—	—	0.0	—	0.0	0.0	0.0
110.0	90.0	0.0	0.0	—	2.6	—	—	0.0	—	0.0	0.0	0.0
110.0	140.0	—	—	—	2.7	—	—	0.0	—	0.0	0.0	0.0
113.0	35.0	—	—	—	0.0	—	—	0.0	—	0.0	0.0	0.0
113.0	45.0	—	—	—	0.0	—	—	0.0	—	0.0	0.0	0.0
113.0	50.0	—	—	—	0.0	—	—	0.0	—	0.0	0.0	0.0
113.0	55.0	—	—	—	0.0	—	—	0.0	—	0.0	0.0	0.0
117.0	50.0	—	—	—	0.0	—	—	0.0	—	0.0	0.0	0.0
117.0	55.0	—	—	—	0.0	—	—	0.0	—	0.0	0.0	0.0
117.0	70.0	—	—	—	0.0	—	—	0.0	—	0.0	0.0	0.0
117.0	85.0	—	—	—	0.0	—	—	0.0	—	0.0	0.0	0.0
117.0	90.0	—	—	—	0.0	—	—	0.0	—	0.0	0.0	0.0
120.0	50.0	—	—	—	0.0	—	—	0.0	—	0.0	0.0	0.0
120.0	60.0	—	—	—	0.0	—	—	0.0	—	0.0	0.0	0.0
120.0	65.0	—	—	—	0.0	—	—	0.0	—	0.0	0.0	0.0
120.0	50.0	—	—	—	2.9	—	—	0.0	—	0.0	0.0	0.0
123.0	50.0	—	—	—	0.0	—	—	0.0	—	0.0	0.0	0.0
123.0	55.0	—	—	—	0.0	—	—	0.0	—	0.0	0.0	0.0
123.0	60.0	—	—	—	0.0	—	—	0.0	—	0.0	0.0	0.0
123.0	80.0	—	—	—	0.0	—	—	0.0	—	0.0	0.0	0.0
127.0	40.0	—	—	—	1.5	—	—	0.0	—	0.0	0.0	0.0

TABLE 4. (cont.)

*Melamphaes* spp. (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
127.0	50.0	-	2.9	-	0.0	-	-	0.0	-	-	0.0	-
127.0	65.0	-	0.0	-	0.0	-	-	8.6	-	-	-	-
127.0	70.0	-	0.0	-	11.5	-	-	2.8	-	-	-	-
127.0	80.0	-	0.0	-	-	-	-	5.5	-	-	-	-
130.0	40.0	-	0.0	-	0.0	-	-	-	-	-	7.9	0.0
130.0	45.0	-	5.6	-	3.0	-	-	-	-	-	2.8	3.0
130.0	55.0	-	0.0	-	0.0	-	-	-	-	-	0.0	3.1
130.0	60.0	-	0.0	-	3.1	-	-	-	-	-	0.0	0.0
130.0	70.0	-	0.0	-	0.0	-	-	-	-	-	2.5	-
130.0	90.0	-	2.7	-	-	-	-	-	-	-	8.1	-
130.0	100.0	-	0.0	-	-	-	-	-	-	-	2.8	-
130.0	120.0	-	0.0	-	-	-	-	-	-	-	2.7	-
133.0	35.0	-	0.0	-	3.1	-	-	0.0	-	-	0.0	-
133.0	40.0	-	0.0	-	3.1	-	-	2.9	-	-	0.0	-
133.0	45.0	-	0.0	-	6.0	-	-	0.0	-	-	0.0	-
137.0	40.0	-	0.0	-	3.0	-	-	-	-	-	0.0	-
137.0	55.0	-	0.0	-	0.0	-	-	-	-	-	2.8	-
137.0	70.0	-	3.0	-	3.2	-	-	-	-	-	0.0	-

*Poromitra* spp.

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
60.0	140.0	0.0	-	-	-	-	-	-	-	-	2.6	-
60.0	160.0	3.2	-	-	-	-	-	-	-	-	2.9	-
60.0	200.0	3.1	-	-	-	-	-	-	-	-	0.0	-
70.0	120.0	3.0	-	-	-	-	-	-	-	-	0.0	-
70.0	180.0	3.0	-	-	-	-	-	-	-	-	8.0	-
70.0	200.0	6.1	-	-	-	-	-	-	-	-	0.0	-
80.0	200.0	2.8	-	-	-	-	-	-	-	-	6.2	-
87.0	90.0	0.0	-	-	0.0	-	-	-	-	-	0.0	-
90.0	160.0	5.2	-	-	-	-	-	-	-	-	5.6	-
90.0	180.0	0.2	-	-	-	-	-	-	-	-	5.5	-
90.0	200.0	0.0	-	-	-	-	-	-	-	-	0.0	-
93.0	45.0	0.0	-	-	-	-	-	-	-	-	3.3	-
93.0	80.0	0.0	-	-	-	-	-	-	-	-	0.0	-
93.0	100.0	-	-	-	-	-	-	-	-	-	2.8	-
97.0	50.0	0.0	-	-	-	-	-	-	-	-	3.3	-
97.0	80.0	0.0	-	-	-	-	-	-	-	-	0.0	-
100.0	35.0	0.0	-	-	-	-	-	-	-	-	0.0	-
100.0	55.0	0.0	-	-	-	-	-	-	-	-	3.2	-
100.0	85.0	-	-	-	-	-	-	-	-	-	3.5	-
100.0	90.0	3.0	-	-	-	-	-	-	-	-	0.0	-
103.0	65.0	2.8	-	-	-	-	-	-	-	-	0.0	-
103.0	90.0	-	-	-	-	-	-	-	-	-	0.0	-
107.0	55.0	-	-	-	-	-	-	-	-	-	3.0	-

TABLE 4. (cont.)

*Poromitra* spp. (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
107.0	60.0	-	6.1	-	0.0	-	0.0	-	0.0	-	-	-
110.0	60.0	-	0.0	-	0.0	-	-	-	0.0	3.0	-	-
123.0	75.0	-	-	-	3.1	-	-	-	-	-	-	-
123.0	80.0	-	0.0	-	3.1	-	-	0.0	-	-	-	-
130.0	60.0	-	2.8	-	0.0	-	-	0.0	0.0	0.0	-	-
133.0	45.0	-	2.9	-	0.0	-	-	0.0	0.0	-	-	-
133.0	65.0	-	0.0	-	3.1	-	-	-	-	-	-	-

*Scopeloberyx robustus*

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
60.0	200.0	0.0	-	-	-	-	-	-	-	-	-	-
70.0	200.0	0.0	-	-	-	-	-	-	-	-	-	-
80.0	160.0	8.8	-	-	-	-	-	-	-	-	-	-
80.0	200.0	0.0	-	-	-	-	-	-	-	-	-	-
90.0	120.0	0.0	-	-	-	-	-	-	-	-	-	-
90.0	140.0	0.0	-	-	-	-	-	-	-	-	-	-
90.0	180.0	0.0	-	-	-	-	-	-	-	-	-	-

*Scopelogadus bispinosus*

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
60.0	140.0	0.0	-	-	-	-	-	-	-	-	-	-
70.0	160.0	3.0	-	-	-	-	-	-	-	-	-	-
80.0	80.0	3.0	-	-	-	-	-	-	-	-	-	-
90.0	70.0	0.0	-	-	-	-	-	-	-	-	-	-
90.0	100.0	0.0	-	-	-	-	-	-	-	-	-	-
100.0	65.0	0.0	-	-	-	-	-	-	-	-	-	-
103.0	65.0	0.0	-	-	-	-	-	-	-	-	-	-
110.0	40.0	-	0.0	-	-	-	-	-	-	-	-	-
110.0	120.0	-	5.2	-	-	-	-	-	-	-	-	-
120.0	100.0	-	3.0	-	-	-	-	-	-	-	-	-
127.0	80.0	-	0.0	-	-	-	-	-	-	-	-	-
130.0	70.0	-	0.0	-	-	-	-	-	-	-	-	-
130.0	120.0	-	0.0	-	-	-	-	-	-	-	-	-

*Macrorhamphosus gracilis*

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
117.0	90.0	-	0.0	-	0.0	-	-	3.3	-	-	-	-
120.0	50.0	-	0.0	-	3.0	-	-	-	0.0	0.0	-	-
120.0	60.0	-	0.0	-	2.9	-	-	-	0.0	0.0	-	-
123.0	50.0	-	2.9	-	0.0	-	-	-	0.0	0.0	-	-

TABLE 4. (cont.)

*Macroramphosus gracilis* (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
127.0	65.0	-	0.0	-	2.9	-	0.0	-	-	-	-	-
133.0	45.0	-	0.0	-	3.0	-	0.0	-	0.0	-	-	-
133.0	50.0	-	0.0	-	5.9	-	0.0	-	0.0	-	-	-

*Syngnathus* spp.

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
67.0	47.2	-	-	-	-	-	-	0.7	-	-	-	-
80.0	65.0	0.0	-	-	0.0	-	0.0	5.9	-	-	-	-
82.0	47.0	0.0	-	-	3.1	-	0.0	-	0.0	-	-	-
93.0	28.0	0.0	-	0.0	-	0.0	-	-	-	-	-	-
93.0	70.0	0.0	-	0.0	-	0.0	-	3.3	-	-	-	-
107.0	30.0	-	0.0	-	0.0	-	1.1	-	-	-	-	-
113.0	30.0	-	1.2	-	0.0	-	0.0	-	1.6	-	-	-
123.0	36.0	-	0.0	-	2.4	-	-	-	-	-	-	-
130.0	25.5	-	0.0	-	0.0	-	-	-	0.0	-	-	-
130.0	30.0	-	0.0	-	0.0	-	-	-	2.0	-	-	-
140.0	28.8	-	0.7	-	-	-	-	0.0	-	-	-	-

## Agonidae

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
67.0	50.0	0.0	-	-	2.7	-	-	0.0	-	-	-	-
70.0	50.0	0.0	-	-	0.0	-	-	3.1	-	-	-	-
80.0	50.2	0.0	-	-	13.3	-	-	0.4	0.0	-	-	-
80.0	52.0	0.0	-	-	2.8	-	-	0.0	0.0	-	-	-
83.0	43.0	11.8	-	-	4.1	-	-	-	-	-	-	-
87.0	40.0	0.0	-	-	0.0	-	-	-	-	-	-	-
93.0	27.0	0.0	-	-	0.0	-	-	-	-	-	-	-
96.8	28.7	2.4	-	-	0.0	-	-	0.0	-	-	-	-
97.0	29.0	0.0	-	-	2.4	-	-	0.0	-	-	-	-
97.0	32.0	0.0	-	-	3.1	-	-	0.0	-	-	-	-
103.0	28.5	2.5	-	-	0.0	-	-	0.0	-	-	-	-
103.0	29.0	0.0	-	-	0.0	-	-	1.7	-	-	-	-
107.0	30.0	-	0.0	-	-	-	-	3.3	-	-	-	-
110.0	32.0	-	2.0	-	-	-	-	-	-	-	-	-
113.0	28.5	-	0.0	-	-	-	-	-	-	-	-	-
113.0	29.0	-	0.0	-	-	-	-	-	-	-	-	-
113.0	30.0	-	1.2	-	-	-	-	-	-	-	-	-
115.0	35.0	-	0.0	-	-	-	-	-	-	-	-	-
120.0	25.0	-	0.0	-	-	-	-	2.7	-	-	-	-

TABLE 4. (cont.)

## Cottidae

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
60.0	50.0	2.6	-	-	-	-	-	0.0	-	-	-	-
63.0	49.7	4.1	-	-	0.9	-	-	-	-	-	-	-
63.0	50.0	4.1	-	-	0.0	-	-	0.0	-	-	-	-
63.0	50.0	2.6	-	-	0.0	-	-	0.0	-	-	-	-
63.0	51.0	-	-	-	0.0	-	-	0.0	-	-	-	-
63.0	52.0	17.4	-	-	4.1	-	-	0.0	-	-	-	-
63.0	55.0	0.0	-	-	0.0	-	-	2.8	-	-	-	-
67.0	47.3	0.7	-	-	0.0	-	-	-	-	-	-	-
67.0	49.0	0.0	-	-	0.0	-	-	3.1	-	-	-	-
70.0	50.0	0.0	-	-	42.4	-	-	0.0	-	-	-	-
73.0	49.4	0.0	-	-	0.7	-	-	0.0	-	-	-	-
73.0	50.0	0.0	-	-	2.8	-	-	0.0	-	-	-	-
77.0	47.6	0.3	-	-	3.2	-	-	0.0	-	-	-	-
77.0	48.0	1.2	-	-	0.6	-	-	0.0	-	-	-	-
77.0	49.0	-	-	-	0.7	-	-	7.0	-	-	-	-
83.0	51.0	0.0	-	-	2.6	-	-	15.8	-	-	-	-
83.0	70.0	0.0	-	-	0.0	-	-	0.0	-	-	-	-
87.0	40.0	0.0	-	-	0.0	-	-	3.3	-	-	-	-
87.0	50.0	3.9	-	-	0.0	-	-	0.0	-	-	-	-
90.0	27.5	0.6	-	-	0.0	-	-	0.0	-	-	-	-
90.0	28.0	0.0	-	-	0.0	-	-	4.7	-	-	-	-
93.0	27.0	0.0	-	-	0.0	-	-	0.0	-	-	-	-
96.8	28.7	3.5	-	-	0.0	-	-	1.0	-	-	-	-
97.0	30.0	0.0	-	-	7.6	-	-	0.0	-	-	-	-
97.0	32.0	0.0	-	-	18.6	-	-	0.0	-	-	-	-
100.0	35.0	0.0	-	-	-	-	-	-	-	-	-	-
103.0	28.5	1.3	-	-	5.4	-	-	2.8	-	-	-	-
113.0	29.0	-	-	-	0.0	-	-	0.0	-	-	-	-
117.0	24.8	-	-	-	0.0	-	-	0.0	-	-	-	-
130.0	25.5	-	-	-	0.6	-	-	-	-	-	-	-

*Scorpaenichthys marmoratus*

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
60.0	51.0	1.6	-	-	-	-	-	0.0	-	-	-	-
63.0	50.0	0.8	-	-	0.0	-	-	0.0	-	-	-	-
63.0	51.0	2.6	-	-	0.0	-	-	0.0	-	-	-	-
67.0	49.0	3.2	-	-	0.0	-	-	0.0	-	-	-	-
70.0	50.0	2.9	-	-	0.0	-	-	0.0	-	-	-	-
70.0	51.0	0.0	-	-	2.8	-	-	0.0	-	-	-	-
73.0	49.4	0.3	-	-	0.0	-	-	0.0	-	-	-	-
73.0	50.0	4.6	-	-	0.0	-	-	0.0	-	-	-	-
73.0	51.0	6.2	-	-	0.0	-	-	0.0	-	-	-	-
73.0	53.0	5.5	-	-	0.0	-	-	0.0	-	-	-	-
77.0	47.6	0.6	-	-	0.0	-	-	0.0	-	-	-	-
77.0	48.0	0.3	-	-	-	-	-	-	-	-	-	-

TABLE 4. (cont.)

*Scorpaenichthys marmoratus* (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
77.0	51.0	5.9	-	-	0.0	-	-	0.0	-	-	-	-
80.0	50.2	1.8	-	-	0.0	-	-	0.0	-	-	-	-
83.0	55.0	0.0	-	-	3.2	-	-	0.0	-	0.0	-	-
93.0	26.5	2.4	-	-	-	-	-	0.0	-	0.0	-	-
97.0	30.0	2.2	-	0.0	-	-	-	0.0	-	0.0	-	-
100.0	29.0	3.8	-	-	0.0	-	-	0.0	-	0.0	-	-
110.0	35.0	-	2.8	-	0.0	-	-	0.0	-	0.0	-	-
117.0	30.0	-	0.0	-	-	-	-	0.0	-	7.8	-	-

## Cyclopteridae

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
60.0	51.0	1.6	-	-	-	-	-	0.0	-	-	-	-
63.0	49.7	3.5	-	-	0.0	-	-	-	-	-	-	-
63.0	50.0	0.8	-	-	0.0	-	-	0.0	-	-	-	-
67.0	47.2	-	-	-	-	-	-	0.7	-	-	-	-
67.0	70.0	-	-	-	-	2.8	-	-	-	-	-	-
70.0	50.0	0.0	-	-	-	2.2	-	0.0	-	-	-	-
77.0	48.0	0.0	-	-	-	0.0	-	-	0.6	-	-	-
80.0	50.2	0.0	-	-	-	0.0	-	-	0.4	0.0	-	-
83.0	51.0	0.0	-	-	-	2.6	-	-	-	0.0	-	-
96.8	28.7	0.0	-	-	0.0	-	-	3.0	-	0.0	-	-
103.0	28.5	1.3	-	-	-	0.0	-	0.0	-	0.0	-	-
103.0	29.0	0.0	-	-	-	0.0	-	1.7	-	0.0	-	-
107.0	30.0	-	0.0	-	-	0.0	-	1.1	-	0.0	-	-
113.0	29.0	-	0.0	-	-	1.2	-	1.8	-	0.0	-	-

## Hexagrammidae

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
77.0	48.0	0.6	-	-	0.0	-	-	0.0	-	-	-	-

## Oxylebius pictus

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
63.0	52.0	2.9	-	-	0.0	-	-	0.0	-	-	-	-
67.0	48.0	0.0	-	-	1.3	-	-	0.0	-	-	-	-
70.0	50.0	5.9	-	-	0.0	-	-	0.0	-	-	-	-
77.0	47.6	0.0	-	-	0.6	-	-	0.3	-	-	-	-
83.0	55.0	0.0	-	-	0.0	-	-	3.3	-	0.0	-	-
87.0	34.0	0.0	-	-	2.5	-	-	0.0	-	0.0	-	-
87.0	50.0	7.9	-	-	0.0	-	-	0.0	-	0.0	-	-
93.0	40.0	3.0	-	-	0.0	-	-	0.0	-	0.0	-	-

TABLE 4. (cont.)

*Oxylebius pictus* (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
100.0	35.0	0.0	-	0.0	-	-	2.8	-	0.0	-	-	-
107.0	31.0	-	2.2	-	0.0	-	0.0	-	0.0	-	-	-
110.0	40.0	-	2.8	-	0.0	-	-	0.0	0.0	-	-	-
117.0	24.8	-	1.0	-	0.0	-	-	0.0	0.0	-	-	-

*Zaniolepis* spp.

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
60.0	52.0	2.3	-	-	-	-	-	0.0	-	-	-	-
77.0	48.0	0.0	-	-	0.6	-	-	0.0	-	-	-	-
93.0	30.0	3.0	-	0.0	-	-	0.0	-	-	-	-	-
97.0	29.0	0.0	-	0.0	-	-	2.6	-	-	0.0	-	-
97.0	30.0	2.2	-	0.0	-	-	0.0	-	-	0.0	-	-
120.0	35.0	-	3.0	-	0.0	-	-	0.0	-	0.0	-	-
123.0	36.0	-	5.3	-	0.0	-	-	0.0	-	0.0	-	-

*Scorpaena* spp.

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
103.0	40.0	0.0	-	0.0	-	0.0	-	-	-	3.1	-	-
110.0	35.0	-	0.0	0.0	-	-	-	-	2.8	0.0	-	-
110.0	60.0	-	0.0	0.0	-	-	-	-	8.3	0.0	-	-
113.0	45.0	-	0.0	0.0	-	0.0	-	-	-	2.5	-	-
115.0	35.0	-	0.0	-	-	-	-	-	-	2.6	3.1	-
118.0	39.0	-	0.0	0.0	-	-	-	0.0	-	-	2.7	-
119.0	33.0	-	0.0	0.0	-	-	-	-	23.9	0.0	-	-
120.0	35.0	-	0.0	0.0	-	-	-	0.0	-	-	2.7	-
120.0	40.0	-	0.0	0.0	-	-	-	2.6	-	-	0.0	-
120.0	45.0	-	0.0	0.0	-	-	-	-	2.9	0.0	-	-
120.0	50.0	-	0.0	0.0	-	-	-	-	15.1	0.0	-	-
123.0	42.0	-	0.0	0.0	-	-	-	-	3.2	0.0	-	-
127.0	40.0	-	0.0	0.0	-	-	-	-	2.8	5.5	0.0	-
130.0	26.0	-	0.0	0.0	-	-	-	-	0.0	0.0	2.3	-
130.0	30.0	-	0.0	0.0	-	-	-	-	0.0	0.0	2.6	-
133.0	30.0	-	0.0	0.0	-	-	-	-	3.0	-	2.9	-
133.0	40.0	-	0.0	0.0	-	-	-	-	2.9	-	0.0	-
133.0	45.0	-	0.0	0.0	-	-	-	-	8.0	-	0.0	-
133.0	50.0	-	0.0	0.0	-	-	-	-	10.5	-	0.0	-
137.0	23.0	-	0.0	0.0	-	-	-	-	1.9	2.8	0.0	-
137.0	40.0	-	0.0	0.0	-	-	-	-	-	2.8	0.0	-

TABLE 4. (cont.)

*Sebastes spp.*

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
60.0	49.5	1.7	-	-	-	-	-	-	-	-	-	-
60.0	50.0	6.5	-	-	-	-	-	-	-	-	-	-
60.0	51.0	129.6	-	-	-	-	-	-	-	-	-	-
60.0	52.0	101.3	-	-	-	-	-	-	-	-	-	-
60.0	55.0	970.3	-	-	-	-	-	-	-	-	-	-
60.0	60.0	10.6	-	-	-	-	-	-	-	-	-	-
60.0	70.0	3.2	-	-	-	-	-	-	-	-	-	-
60.0	80.0	10.2	-	-	-	-	-	-	-	-	-	-
60.0	90.0	6.7	-	-	-	-	-	-	-	-	-	-
60.0	120.0	0.0	-	-	-	-	-	-	-	-	-	-
63.0	49.7	7.5	-	-	-	-	-	-	-	-	-	-
63.0	50.0	59.0	-	-	-	-	-	-	-	-	-	-
63.0	51.0	81.5	-	-	-	-	-	-	-	-	-	-
63.0	52.0	745.3	-	-	-	-	-	-	-	-	-	-
63.0	55.0	1328.2	-	-	-	-	-	-	-	-	-	-
63.0	60.0	6.5	-	-	-	-	-	-	-	-	-	-
63.0	65.0	-	-	-	-	-	-	-	-	-	-	-
63.0	70.0	-	-	-	-	-	-	-	-	-	-	-
63.0	80.0	-	-	-	-	-	-	-	-	-	-	-
67.0	48.0	2.5	-	-	-	-	-	-	-	-	-	-
67.0	49.0	268.1	-	-	-	-	-	-	-	-	-	-
67.0	50.0	669.8	-	-	-	-	-	-	-	-	-	-
67.0	55.0	527.9	-	-	-	-	-	-	-	-	-	-
67.0	60.0	72.2	-	-	-	-	-	-	-	-	-	-
67.0	65.0	-	-	-	-	-	-	-	-	-	-	-
67.0	70.0	-	-	-	-	-	-	-	-	-	-	-
67.0	80.0	-	-	-	-	-	-	-	-	-	-	-
70.0	50.0	1535.3	-	-	-	-	-	-	-	-	-	-
70.0	51.0	586.4	-	-	-	-	-	-	-	-	-	-
70.0	52.0	315.0	-	-	-	-	-	-	-	-	-	-
70.0	53.0	171.1	-	-	-	-	-	-	-	-	-	-
70.0	55.0	66.7	-	-	-	-	-	-	-	-	-	-
70.0	60.0	8.6	-	-	-	-	-	-	-	-	-	-
70.0	65.0	-	-	-	-	-	-	-	-	-	-	-
70.0	70.0	51.5	-	-	-	-	-	-	-	-	-	-
70.0	80.0	0.0	-	-	-	-	-	-	-	-	-	-
73.0	49.4	41.2	-	-	-	-	-	-	-	-	-	-
73.0	50.0	1122.9	-	-	-	-	-	-	-	-	-	-
73.0	51.0	522.2	-	-	-	-	-	-	-	-	-	-
73.0	53.0	728.9	-	-	-	-	-	-	-	-	-	-
73.0	60.0	139.5	-	-	-	-	-	-	-	-	-	-
73.0	65.0	-	-	-	-	-	-	-	-	-	-	-
73.0	70.0	-	-	-	-	-	-	-	-	-	-	-
73.0	80.0	-	-	-	-	-	-	-	-	-	-	-
77.0	47.6	4.8	-	-	-	-	-	-	-	-	-	-
77.0	48.0	2.3	-	-	-	-	-	-	-	-	-	-

TABLE 4. (cont.)

*Sebastes spp.* (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
77.0	49.0	-	-	-	1.3	-	-	0.0	-	-	-	-
77.0	51.0	793.0	-	-	21.1	-	-	172.2	-	-	-	-
77.0	55.0	100.8	-	-	47.1	-	-	18.6	-	-	-	-
77.0	57.0	14.9	-	-	53.9	-	-	2.8	-	-	-	-
77.0	60.0	-	-	-	15.7	-	-	-	-	-	-	-
77.0	65.0	-	-	-	5.1	-	-	-	-	-	-	-
77.0	70.0	-	-	-	2.3	-	-	-	-	-	-	-
80.0	50.2	29.2	-	-	9.8	-	-	0.0	-	-	-	-
80.0	51.0	214.9	-	-	17.5	-	-	0.0	-	-	-	-
80.0	52.0	329.0	-	-	113.2	-	-	6.5	-	-	-	-
80.0	55.0	80.9	-	-	12.8	-	-	4.8	-	-	-	-
80.0	60.0	12.8	-	-	13.6	-	-	5.6	-	-	-	-
80.0	65.0	18.7	-	-	3.4	-	-	0.0	-	-	-	-
80.0	70.0	0.0	-	-	10.2	-	-	6.1	-	-	-	-
80.0	80.0	3.0	-	-	9.7	-	-	0.0	-	-	-	-
82.0	47.0	2.9	-	-	316.2	-	-	27.8	-	-	-	-
83.0	40.0	34.4	-	-	42.0	-	-	5.4	-	-	-	-
83.0	43.0	85.8	-	-	0.0	-	-	1.5	-	-	-	-
83.0	51.0	68.4	-	-	42.0	-	-	3.2	-	-	-	-
83.0	55.0	147.9	-	-	67.3	-	-	18.9	-	-	-	-
83.0	60.0	9.1	-	-	56.9	-	-	0.0	-	-	-	-
83.0	65.0	28.7	-	-	3.0	-	-	0.0	-	-	-	-
83.0	70.0	25.5	-	-	0.0	-	-	6.6	-	-	-	-
83.0	80.0	0.0	-	-	0.0	-	-	20.2	-	-	-	-
87.0	32.5	4.7	-	-	11.5	-	-	4.1	-	-	-	-
87.0	33.0	6.0	-	-	11.7	-	-	1.7	-	-	-	-
87.0	34.0	157.5	-	-	42.1	-	-	2.7	-	-	-	-
87.0	35.0	132.3	-	-	7.4	-	-	39.6	-	-	-	-
87.0	40.0	279.3	-	-	27.3	-	-	15.3	-	-	-	-
87.0	45.0	153.0	-	-	45.5	-	-	0.0	-	-	-	-
87.0	50.0	851.0	-	-	17.1	-	-	3.3	-	-	-	-
87.0	55.0	98.3	-	-	300.3	-	-	116.5	-	-	-	-
87.0	60.0	0.0	-	-	2.8	-	-	6.5	-	-	-	-
87.0	65.0	0.0	-	-	2.4	-	-	3.3	-	-	-	-
87.0	80.0	0.0	-	-	2.7	-	-	0.0	-	-	-	-
90.0	27.5	2.3	-	-	0.0	-	-	9.8	-	-	-	-
90.0	28.0	41.8	-	-	15.3	-	-	0.0	-	-	-	-
90.0	32.0	72.9	-	-	18.3	-	-	0.0	-	-	-	-
90.0	37.0	65.8	-	-	12.4	-	-	2.1	-	-	-	-
90.0	45.0	30.5	-	-	97.0	-	-	0.0	-	-	-	-
90.0	53.0	22.6	-	-	3.3	-	-	5.7	-	-	-	-
90.0	60.0	3.0	-	-	0.0	-	-	40.7	-	-	-	-
90.0	70.0	0.0	-	-	0.0	-	-	2.8	-	-	-	-
90.0	90.0	9.0	-	-	0.0	-	-	0.0	-	-	-	-
93.0	26.5	7.2	-	-	-	-	-	0.0	-	-	-	-
93.0	27.0	39.6	-	-	-	-	-	11.5	-	-	-	-
		7.9	-	-	-	-	-	-	-	-	-	-

TABLE 4. (cont.)

*Sebastes* spp. (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
93.0	28.0	30.9	-	23.2	-	-	16.1	-	-	13.4	-	-
93.0	30.0	60.8	-	14.3	-	-	3.1	-	-	0.0	-	-
93.0	35.0	15.0	-	2.9	-	-	0.0	-	-	0.0	-	-
93.0	40.0	3.0	-	0.0	-	-	12.6	-	-	0.0	-	-
93.0	45.0	12.2	-	0.0	-	-	26.2	-	-	0.0	-	-
93.0	50.0	12.4	-	0.0	-	-	62.7	-	-	0.0	-	-
93.0	55.0	70.6	-	0.0	-	-	16.5	-	-	0.0	-	-
93.0	60.0	21.4	-	10.1	-	-	10.0	-	-	0.0	-	-
93.0	60.0	0.0	-	23.3	-	-	3.3	-	-	0.0	-	-
93.0	70.0	-	-	-	-	-	0.0	-	-	2.9	-	-
96.8	28.7	3.5	-	0.0	-	-	0.0	-	-	0.0	-	-
97.0	29.0	39.3	-	4.8	-	-	0.0	-	-	2.3	-	-
97.0	30.0	27.0	-	25.3	-	-	0.0	-	-	0.0	-	-
97.0	32.0	0.0	-	372.0	-	-	10.1	-	-	0.0	-	-
97.0	40.0	0.0	-	6.4	-	-	22.9	-	-	0.0	-	-
97.0	45.0	3.0	-	0.0	-	-	0.0	-	-	0.0	-	-
97.0	50.0	113.4	-	3.2	-	-	0.0	-	-	3.0	-	-
97.0	55.0	36.0	-	-	-	-	0.0	-	-	0.0	-	-
97.0	60.0	17.8	-	-	-	-	0.0	-	-	0.0	-	-
97.0	65.0	21.5	-	3.5	-	-	0.0	-	-	2.8	-	-
97.0	70.0	5.9	-	-	-	-	0.0	-	-	0.0	-	-
97.0	75.0	-	-	-	-	-	31.1	-	-	0.0	-	-
100.0	29.0	615.0	-	-	-	-	101.9	-	-	0.0	-	-
100.0	30.0	25.3	-	-	-	-	70.6	-	-	5.0	-	-
100.0	35.0	-	-	-	-	-	0.0	-	-	0.0	-	-
100.0	45.0	3.1	-	-	-	-	0.0	-	-	0.0	-	-
100.0	55.0	3.0	-	-	-	-	0.0	-	-	0.0	-	-
100.0	60.0	3.0	-	-	-	-	6.7	-	-	0.0	-	-
100.0	65.0	3.0	-	-	-	-	12.8	-	-	0.0	-	-
100.0	70.0	0.0	-	-	-	-	3.3	-	-	0.0	-	-
103.0	28.5	16.3	-	-	-	-	18.1	-	-	1.0	-	-
103.0	29.0	160.3	-	-	-	-	9.5	-	-	0.0	-	-
103.0	30.0	130.4	-	-	-	-	532.4	-	-	2.6	-	-
103.0	35.0	70.6	-	-	-	-	105.7	-	-	0.0	-	-
103.0	40.0	17.5	-	-	-	-	15.3	-	-	0.0	-	-
103.0	45.0	3.0	-	-	-	-	12.1	-	-	3.2	-	-
103.0	55.0	2.9	-	-	-	-	0.0	-	-	0.0	-	-
103.0	60.0	0.0	-	-	-	-	13.4	-	-	0.0	-	-
103.0	65.0	2.8	-	-	-	-	0.0	-	-	0.0	-	-
103.0	70.0	3.0	-	-	-	-	0.0	-	-	0.0	-	-
107.0	30.0	-	-	-	-	-	16.5	-	-	0.0	-	-
107.0	31.0	-	-	-	-	-	131.2	-	-	0.0	-	-
107.0	32.0	-	-	-	-	-	349.2	-	-	3.4	-	-
107.0	35.0	-	-	-	-	-	166.4	-	-	6.9	-	-
107.0	40.0	-	-	-	-	-	0.0	-	-	3.3	-	-
107.0	80.0	-	-	-	-	-	0.0	-	-	0.0	-	-

TABLE 4. (cont.)

*Sebastes spp.* (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
110.0	32.0	-	22.1	-	135.0	-	-	-	0.0	0.0	-	-
110.0	40.0	-	2.8	-	8.6	-	-	-	0.0	0.0	-	-
110.0	45.0	-	0.0	-	3.2	-	-	-	0.0	0.0	-	-
113.0	28.5	-	1.8	-	2.2	-	0.0	-	-	-	-	-
113.0	29.0	-	4.3	-	17.1	-	0.0	-	-	-	-	-
113.0	30.0	-	30.1	-	36.1	-	3.0	-	0.0	-	-	-
113.0	35.0	-	0.0	-	0.0	-	3.3	-	-	-	-	-
113.0	40.0	-	2.9	-	0.0	-	0.0	-	-	-	-	-
113.0	45.0	-	3.1	-	0.0	-	0.0	-	-	-	-	-
113.0	50.0	-	3.0	-	0.0	-	0.0	-	-	-	-	-
113.0	80.0	-	2.8	-	0.0	-	0.0	-	-	-	-	-
115.0	35.0	-	11.4	-	-	-	0.0	-	0.0	-	-	-
117.0	24.8	-	3.0	-	0.0	-	0.0	-	0.0	-	-	-
117.0	25.0	-	5.2	-	8.4	-	0.0	-	0.0	-	-	-
117.0	26.0	-	154.0	-	3.0	-	0.0	-	0.0	-	-	-
117.0	30.0	-	26.0	-	6.5	-	0.0	-	0.0	-	-	-
117.0	35.0	-	9.4	-	16.3	-	0.0	-	0.0	-	-	-
117.0	40.0	-	21.4	-	36.7	-	0.0	-	0.0	-	-	-
117.0	45.0	-	8.9	-	8.7	-	3.4	-	0.0	-	-	-
117.0	50.0	-	3.2	-	11.6	-	3.3	-	0.0	-	-	-
117.0	55.0	-	0.0	-	0.0	-	3.3	-	0.0	-	-	-
118.0	39.0	-	9.3	-	37.1	-	0.0	-	0.0	-	-	-
119.0	33.0	-	8.0	-	2.9	-	0.0	-	0.0	-	-	-
120.0	30.0	-	34.8	-	0.0	-	0.0	-	0.0	-	-	-
120.0	35.0	-	24.4	-	0.0	-	0.0	-	0.0	-	-	-
120.0	45.0	-	20.0	-	3.1	-	0.0	-	0.0	-	-	-
120.0	50.0	-	21.5	-	3.0	-	0.0	-	0.0	-	-	-
120.0	65.0	-	2.8	-	0.0	-	1.5	-	0.0	-	-	-
123.0	36.0	-	5.3	-	14.5	-	0.0	-	0.0	-	-	-
123.0	37.0	-	62.9	-	17.2	-	0.0	-	0.0	-	-	-
123.0	42.0	-	6.3	-	0.0	-	0.0	-	0.0	-	-	-
123.0	45.0	-	0.0	-	5.8	-	0.0	-	0.0	-	-	-
123.0	65.0	-	0.0	-	0.0	-	2.8	-	0.0	-	-	-
123.5	35.6	-	2.2	-	17.1	-	0.0	-	0.0	-	-	-
127.0	32.6	-	6.0	-	3.3	-	0.0	-	0.0	-	-	-
127.0	33.0	-	34.2	-	16.1	-	0.0	-	0.0	-	-	-
127.0	34.0	-	10.5	-	22.0	-	2.4	-	0.0	-	-	-
127.0	40.0	-	0.0	-	62.4	-	0.0	-	0.0	-	-	-
127.0	45.0	-	0.0	-	0.0	-	3.2	-	0.0	-	-	-
130.0	28.0	-	0.0	-	2.7	-	0.0	-	0.0	-	-	-
130.0	35.0	-	3.3	-	0.0	-	0.0	-	0.0	-	-	-
130.0	40.0	-	0.0	-	31.5	-	0.0	-	0.0	-	-	-
133.0	21.0	-	2.0	-	0.0	-	0.0	-	0.0	-	-	-
133.0	23.0	-	0.0	-	2.6	-	0.0	-	0.0	-	-	-
133.0	25.0	-	6.1	-	6.1	-	9.3	-	0.0	-	-	-
133.0	30.0	-	0.0	-	0.0	-	0.0	-	0.0	-	-	-

TABLE 4. (cont.)

*Sebastes* spp. (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
133.0	35.0	-	11.5	-	24.7	-	-	0.0	-	0.0	-	-
133.0	40.0	-	3.0	-	0.0	-	-	0.0	-	0.0	-	-
133.0	45.0	-	14.7	-	0.0	-	-	0.0	-	0.0	-	-
133.0	50.0	-	2.8	-	0.0	-	-	0.0	-	0.0	-	-
137.0	23.0	-	0.0	-	2.6	-	-	0.0	-	0.0	-	-
137.0	30.0	-	0.0	-	15.5	-	-	0.0	-	0.0	-	-
137.0	35.0	-	0.0	-	9.0	-	-	0.0	-	0.0	-	-

*Sebastolobus* spp.

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
60.0	90.0	0.0	-	-	-	-	-	8.3	-	-	-	-
60.0	100.0	0.0	-	-	-	-	-	2.6	-	-	-	-
60.0	120.0	0.0	-	-	-	-	-	2.7	-	-	-	-
67.0	48.0	0.0	-	-	1.3	-	-	0.0	-	-	-	-
67.0	49.0	0.0	-	-	5.7	-	-	0.0	-	-	-	-
67.0	55.0	0.0	-	-	2.7	-	-	2.9	-	-	-	-
67.0	60.0	0.0	-	-	2.8	-	-	0.0	-	-	-	-
67.0	65.0	-	-	-	14.6	-	-	-	-	-	-	-
67.0	70.0	-	-	-	8.5	-	-	-	-	-	-	-
70.0	53.0	0.0	-	-	0.0	-	-	2.9	-	-	-	-
70.0	60.0	0.0	-	-	5.7	-	-	0.0	-	-	-	-
70.0	65.0	-	-	-	2.7	-	-	-	-	-	-	-
77.0	65.0	-	-	-	2.5	-	-	-	-	-	-	-
77.0	70.0	-	-	-	2.3	-	-	-	-	-	-	-
80.0	70.0	0.0	-	-	0.0	-	-	0.0	-	2.7	-	-
80.0	80.0	0.0	-	-	6.4	-	-	0.0	-	0.0	-	-
80.0	120.0	0.0	-	-	-	-	-	3.0	-	0.0	-	-
83.0	70.0	0.0	-	-	3.2	-	-	0.0	-	0.0	-	-
87.0	80.0	0.0	-	-	0.0	-	-	3.3	-	-	-	-

*Prionotus* spp.

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
117.0	26.0	-	0.0	-	0.0	-	-	0.0	-	2.6	-	-
118.0	39.0	-	0.0	-	0.0	-	-	0.0	-	5.4	-	-
119.0	33.0	-	0.0	-	0.0	-	-	6.5	-	14.3	-	-
120.0	23.0	-	0.0	-	0.0	-	-	0.0	-	7.5	-	-
120.0	24.0	-	0.0	-	0.0	-	-	0.0	-	7.1	-	-
120.0	25.0	-	0.0	-	0.0	-	-	0.0	-	2.3	-	-
120.0	30.0	-	0.0	-	0.0	-	-	0.0	-	3.0	-	-
120.0	40.0	-	0.0	-	0.0	-	-	0.0	-	5.5	-	-
130.0	25.5	-	0.0	-	0.0	-	-	0.0	-	0.0	-	-
130.0	26.0	-	0.0	-	0.0	-	-	0.0	-	6.6	-	-

TABLE 4. (cont.)

*Prionotus* spp. (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
130.0	28.0	-	0.0	-	0.0	-	-	-	0.0	4.7	-	-
130.0	30.0	-	0.0	-	0.0	-	-	-	0.0	10.3	-	-
133.0	21.0	-	0.0	-	0.0	-	-	-	0.0	52.7	-	-
133.0	23.0	-	0.0	-	0.0	-	-	-	0.0	45.0	-	-
133.0	25.0	-	0.0	-	0.0	-	-	-	0.0	20.6	-	-
133.0	30.0	-	0.0	-	0.0	-	-	-	0.0	17.2	-	-
133.0	35.0	-	0.0	-	0.0	-	-	-	0.0	8.9	-	-
133.0	35.0	-	0.0	-	0.0	-	-	-	0.0	3.8	-	-
137.0	20.0	-	0.0	-	0.0	-	-	-	20.5	21.6	-	-
137.0	21.0	-	0.0	-	0.0	-	-	-	194.7	42.8	-	-
137.0	22.0	-	0.0	-	0.0	-	-	-	53.2	74.5	-	-
137.0	23.0	-	0.0	-	0.0	-	-	-	0.0	79.3	-	-
137.0	30.0	-	0.0	-	0.0	-	-	-	0.0	7.7	-	-
140.0	28.8	-	0.0	-	0.0	-	-	-	201.9	-	-	-
140.0	29.0	-	0.0	-	0.0	-	-	-	2.8	-	-	-
140.0	30.0	-	0.0	-	0.0	-	-	-	-	-	-	-

*Blennioidei*

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
60.0	55.0	6.2	-	-	-	-	-	0.0	-	-	-	-
77.0	47.6	0.0	-	-	6.3	-	-	0.0	-	-	-	-
77.0	48.0	0.0	-	-	5.6	-	-	0.0	-	-	-	-
97.0	30.0	2.2	-	0.0	-	-	0.0	-	-	0.0	-	-

*Hypsoblennius* spp.

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
77.0	47.6	0.0	-	-	0.0	-	-	0.3	-	-	-	-
77.0	48.0	0.0	-	-	0.0	-	-	1.2	-	-	-	-
80.0	51.0	0.0	-	-	0.0	-	-	1.6	0.0	-	19.3	-
83.0	39.0	0.0	-	-	-	-	-	0.0	-	-	2.4	-
83.0	40.0	0.0	-	-	0.0	-	-	1.5	-	-	5.8	-
83.0	43.0	0.0	-	-	0.0	-	-	0.0	-	-	16.6	-
87.0	32.5	0.0	-	-	0.0	-	-	0.0	-	-	5.2	-
87.0	33.0	0.0	-	-	0.0	-	-	0.0	-	-	21.1	-
87.0	34.0	0.0	-	-	0.0	-	-	0.0	-	-	6.0	-
87.0	35.0	0.0	-	-	0.0	-	-	0.0	-	-	12.2	-
87.0	40.0	0.0	-	-	0.0	-	-	0.0	-	-	6.4	-
87.0	45.0	0.0	-	-	0.0	-	-	0.0	-	-	16.9	-
90.0	27.5	0.0	-	-	0.0	-	-	-	12.2	-	-	-
90.0	28.0	0.0	-	-	0.0	-	-	-	7.0	-	0.0	-
90.0	37.0	0.0	-	-	0.0	-	-	-	0.0	-	2.9	-
93.0	26.5	0.0	-	-	0.0	-	-	-	-	-	31.6	-
93.0	27.0	0.0	-	-	-	-	-	8.6	-	-	0.0	-

TABLE 4. (cont.)

*Hypsoblennius* spp. (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
93.0	28.0	0.0	-	0.0	-	-	22.5	-	-	13.4	-	-
93.0	30.0	0.0	-	0.0	-	-	3.1	-	-	0.0	-	-
96.8	28.7	0.0	-	0.0	-	-	0.0	-	-	16.7	-	-
97.0	29.0	0.0	-	0.0	-	-	0.0	-	-	12.2	-	-
97.0	30.0	0.0	-	0.0	-	-	0.0	-	-	11.4	-	-
100.0	29.0	0.0	-	0.0	-	-	0.0	-	-	4.2	-	-
103.0	28.5	0.0	-	0.0	-	-	0.0	-	-	5.0	-	-
103.0	29.0	0.0	-	0.0	-	-	0.0	-	-	5.1	-	-
107.0	30.0	0.0	-	0.0	-	-	0.0	-	-	0.6	-	-
107.0	31.0	0.0	-	0.0	-	-	1.9	-	-	2.4	-	-
110.0	32.0	0.0	-	0.0	-	-	-	-	0.0	0.9	-	-
110.0	35.0	0.0	-	0.0	-	-	-	-	2.8	0.0	-	-
110.0	40.0	0.0	-	0.0	-	-	-	-	0.0	3.1	-	-
113.0	28.5	0.0	-	0.0	-	-	1.3	-	-	5.6	-	-
113.0	29.0	0.0	-	0.0	-	-	5.4	-	-	1.8	-	-
117.0	24.8	0.0	-	0.0	-	-	-	-	0.0	1.1	-	-
117.0	26.0	0.0	-	0.0	-	-	-	-	0.0	5.3	-	-
117.0	35.0	0.0	-	0.0	-	-	-	-	0.0	5.8	-	-
118.0	39.0	0.0	-	0.0	-	-	-	-	0.0	5.4	-	-
119.0	33.0	0.0	-	0.0	-	-	-	-	-	0.0	0.0	-
120.0	22.5	0.0	-	0.0	-	-	2.0	-	1.0	-	0.0	-
120.0	24.0	0.0	-	0.0	-	-	0.0	-	1.2	-	2.4	-
120.0	40.0	0.0	-	0.0	-	-	0.0	-	0.0	-	2.7	-
123.0	37.0	0.0	-	0.0	-	-	-	-	2.3	-	0.0	-
127.0	32.6	0.0	-	0.0	-	-	-	-	1.4	-	0.9	-
127.0	33.0	0.0	-	0.0	-	-	-	-	0.0	-	2.6	-
127.0	40.0	0.0	-	0.0	-	-	-	-	8.5	0.0	0.0	-
130.0	25.5	0.0	-	0.0	-	-	3.6	-	-	8.1	7.7	-
130.0	26.0	0.0	-	0.0	-	-	4.8	-	-	0.8	11.6	-
130.0	30.0	0.0	-	0.0	-	-	-	-	-	2.0	7.7	-
133.0	19.5	0.0	-	0.0	-	-	-	-	0.0	-	13.3	-
133.0	21.0	0.0	-	0.0	-	-	-	-	4.0	-	4.6	-
133.0	23.0	0.0	-	0.0	-	-	-	-	0.0	-	2.5	-
133.0	25.0	0.0	-	2.7	-	-	-	-	0.0	-	0.0	-
133.0	30.0	0.0	-	0.0	-	-	-	-	3.0	-	0.0	-
137.0	20.0	0.0	-	3.6	-	-	-	-	0.0	-	0.0	-
137.0	21.0	0.0	-	0.0	-	-	-	-	4.3	-	2.2	-
137.0	22.0	0.0	-	0.0	-	-	-	-	1.9	-	0.0	-
137.0	23.0	0.0	-	0.0	-	-	-	-	-	1.4	-	-
140.0	28.8	0.0	-	-	-	-	-	-	-	-	-	-

## Clinidae

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
60.0	50.0	2.6	-	-	-	-	-	-	0.0	-	-	-

TABLE 4. (cont.)

## Clinidae (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
63.0	49.7	11.0	-	-	0.9	-	-	-	-	-	-	-
63.0	50.0	21.3	-	-	1.6	-	-	-	-	-	-	-
63.0	51.0	26.3	-	-	0.0	-	-	-	-	-	-	-
67.0	48.0	1.2	-	-	1.3	-	-	-	-	-	-	-
70.0	50.0	0.0	-	-	2.2	-	-	-	-	-	-	-
73.0	49.4	0.3	-	-	2.1	-	-	-	-	-	-	-
77.0	47.6	0.3	-	-	1.3	-	-	-	-	-	-	-
77.0	48.0	0.3	-	-	4.3	-	-	-	-	-	-	-
77.0	49.0	-	-	-	0.7	-	-	-	-	-	-	-
80.0	50.2	0.0	-	-	0.8	-	-	-	-	-	-	-
80.0	51.0	0.0	-	-	3.5	-	-	-	-	-	-	-
83.0	43.0	0.0	-	-	2.8	-	-	-	-	-	-	-
83.0	51.0	0.0	-	-	0.0	-	-	-	-	-	-	-
83.0	55.0	0.0	-	-	0.0	-	-	-	-	-	-	-
87.0	40.0	5.7	-	-	0.0	-	-	-	-	-	-	-
87.0	50.0	7.9	-	-	0.0	-	-	-	-	-	-	-
93.0	26.5	0.0	-	-	-	-	-	-	-	-	-	-
97.0	29.0	0.0	-	-	0.0	-	-	-	-	-	-	-
97.0	30.0	0.0	-	-	2.5	-	-	-	-	-	-	-
100.0	29.0	0.0	-	-	-	-	-	-	-	-	-	-
103.0	28.5	2.5	-	-	-	-	-	-	-	-	-	-
103.0	29.0	2.3	-	-	-	-	-	-	-	-	-	-
103.0	30.0	2.5	-	-	-	-	-	-	-	-	-	-
103.0	45.0	0.0	-	-	-	-	-	-	-	-	-	-
107.0	30.0	-	-	-	0.1	-	-	-	-	-	-	-
107.0	32.0	-	-	-	2.0	-	-	-	-	-	-	-
110.0	32.0	-	-	-	0.0	-	-	-	-	-	-	-
113.0	28.5	-	-	-	0.0	-	-	-	-	-	-	-
113.0	29.0	-	-	-	6.5	-	-	-	-	-	-	-
113.0	30.0	-	-	-	3.7	-	-	-	-	-	-	-
117.0	26.0	-	-	-	2.8	-	-	-	-	-	-	-
120.0	24.0	-	-	-	0.0	-	-	-	-	-	-	-
120.0	40.0	-	-	-	5.3	-	-	-	-	-	-	-
123.0	36.0	-	-	-	0.0	-	-	-	-	-	-	-
123.0	37.0	-	-	-	0.0	-	-	-	-	-	-	-
123.5	35.6	-	-	-	2.2	-	-	-	-	-	-	-
127.0	32.6	-	-	-	6.1	-	-	-	-	-	-	-
127.0	33.0	-	-	-	0.0	-	-	-	-	-	-	-
127.0	34.0	-	-	-	7.2	-	-	-	-	-	-	-
130.0	25.5	-	-	-	2.5	-	-	-	-	-	-	-
130.0	26.0	-	-	-	6.1	-	-	-	-	-	-	-
133.0	19.5	-	-	-	0.0	-	-	-	-	-	-	-
137.0	22.0	-	-	-	0.0	-	-	-	-	-	-	-
137.0	30.0	-	-	-	0.0	-	-	-	-	-	-	-
140.0	28.8	-	-	-	0.0	-	-	-	-	-	-	-
140.0	29.0	-	-	-	0.0	-	-	-	-	-	-	-

TABLE 4. (cont.)

## Gobiidae

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
60.0	49.5	1.7	-	-	-	-	-	-	-	-	-	-
60.0	50.0	14.3	-	-	-	-	-	-	-	-	-	-
60.0	51.0	19.4	-	-	-	-	-	-	-	-	-	-
60.0	52.0	6.8	-	-	-	-	-	-	-	-	-	-
63.0	50.0	0.0	-	-	-	-	-	-	-	-	-	-
63.0	51.0	0.0	-	-	-	-	-	-	-	-	-	-
63.0	52.0	0.0	-	-	-	-	-	-	-	-	-	-
67.0	48.0	1.2	-	-	-	-	-	-	-	-	-	-
67.0	50.0	2.3	-	-	-	-	-	-	-	-	-	-
67.0	60.0	0.0	-	-	-	-	-	-	-	-	-	-
67.0	65.0	-	-	-	-	-	-	-	-	-	-	-
70.0	51.0	0.0	-	-	-	-	-	-	-	-	-	-
70.0	52.0	0.0	-	-	-	-	-	-	-	-	-	-
70.0	53.0	0.0	-	-	-	-	-	-	-	-	-	-
70.0	55.0	2.9	-	-	-	-	-	-	-	-	-	-
70.0	60.0	0.0	-	-	-	-	-	-	-	-	-	-
70.0	65.0	-	-	-	-	-	-	-	-	-	-	-
73.0	51.0	0.0	-	-	-	-	-	-	-	-	-	-
73.0	53.0	2.7	-	-	-	-	-	-	-	-	-	-
80.0	65.0	0.0	-	-	-	-	-	-	-	-	-	-
82.0	47.0	0.0	-	-	-	-	-	-	-	-	-	-
83.0	43.0	0.0	-	-	-	-	-	-	-	-	-	-
83.0	51.0	0.0	-	-	-	-	-	-	-	-	-	-
83.0	70.0	0.0	-	-	-	-	-	-	-	-	-	-
87.0	33.0	2.0	-	-	-	-	-	-	-	-	-	-
87.0	34.0	0.0	-	-	-	-	-	-	-	-	-	-
87.0	50.0	0.0	-	-	-	-	-	-	-	-	-	-
90.0	27.5	0.6	-	-	-	-	-	-	-	-	-	-
90.0	37.0	0.0	-	-	-	-	-	-	-	-	-	-
90.0	60.0	0.0	-	-	-	-	-	-	-	-	-	-
93.0	26.5	0.0	-	-	-	-	-	-	-	-	-	-
93.0	27.0	0.0	-	-	-	-	-	-	-	-	-	-
93.0	28.0	0.0	-	-	-	-	-	-	-	-	-	-
93.0	50.0	0.0	-	-	-	-	-	-	-	-	-	-
97.0	29.0	0.0	-	-	-	-	-	-	-	-	-	-
97.0	30.0	0.0	-	-	-	-	-	-	-	-	-	-
97.0	32.0	0.0	-	-	-	-	-	-	-	-	-	-
97.0	40.0	0.0	-	-	-	-	-	-	-	-	-	-
97.0	45.0	0.0	-	-	-	-	-	-	-	-	-	-
100.0	50.0	0.0	-	-	-	-	-	-	-	-	-	-
103.0	28.5	2.5	-	-	-	-	-	-	-	-	-	-
103.0	29.0	0.0	-	-	-	-	-	-	-	-	-	-
103.0	30.0	7.4	-	-	-	-	-	-	-	-	-	-
103.0	35.0	5.9	-	-	-	-	-	-	-	-	-	-
103.0	45.0	0.0	-	-	-	-	-	-	-	-	-	-
107.0	31.0	0.0	-	-	-	-	-	-	-	-	-	-

TABLE 4. (cont.)

## Gobiidae (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
107.0	32.0	-	6.2	-	9.2	-	3.4	-	-	0.0	-	-
107.0	35.0	-	3.2	-	0.0	-	0.0	-	-	0.0	-	-
107.0	70.0	-	0.0	-	0.0	-	0.0	-	-	3.0	-	-
110.0	32.0	-	0.0	-	0.0	-	0.0	-	-	0.0	2.6	-
113.0	29.0	-	0.0	-	0.0	-	0.0	-	-	1.8	-	-
113.0	50.0	-	3.0	-	0.0	-	0.0	-	-	0.0	-	-
115.0	35.0	-	0.0	-	-	-	-	-	-	0.0	3.1	-
117.0	35.0	-	0.0	-	0.0	-	0.0	-	-	2.9	-	-
117.0	40.0	-	0.0	-	0.0	-	0.0	-	-	3.1	-	-
118.0	39.0	-	0.0	-	0.0	-	0.0	-	-	2.7	-	-
120.0	22.5	-	0.0	-	0.0	-	0.0	-	-	15.7	-	-
120.0	23.0	-	0.0	-	2.8	-	0.0	-	-	0.0	-	-
120.0	24.0	-	47.7	-	0.0	-	0.0	-	-	4.8	-	-
120.0	25.0	-	7.3	-	22.7	-	0.0	-	-	0.0	-	-
120.0	35.0	-	0.0	-	0.0	-	0.0	-	-	2.6	-	-
120.0	40.0	-	0.0	-	0.0	-	0.0	-	-	10.9	-	-
120.0	60.0	-	0.0	-	0.0	-	0.0	-	-	0.0	-	-
123.5	35.6	-	0.0	-	0.0	-	0.0	-	-	4.9	-	-
127.0	32.6	-	0.0	-	1.7	-	0.0	-	-	58.3	-	-
127.0	33.0	-	2.8	-	0.0	-	0.0	-	-	2.6	-	-
127.0	34.0	-	0.0	-	0.0	-	0.0	-	-	3.0	-	-
130.0	25.5	-	0.0	-	0.0	-	0.0	-	-	4.9	-	-
130.0	35.6	-	0.0	-	1.7	-	0.0	-	-	0.0	2.4	-
130.0	28.0	-	0.0	-	0.0	-	0.0	-	-	0.0	5.2	-
130.0	30.0	-	0.0	-	0.0	-	0.0	-	-	0.0	6.1	-
133.0	19.5	-	30.6	-	0.0	-	2.5	-	-	18.3	-	-
133.0	21.0	-	8.0	-	0.0	-	2.0	-	-	2.5	-	-
133.0	23.0	-	0.0	-	0.0	-	0.0	-	-	0.0	5.7	-
133.0	28.0	-	0.0	-	0.0	-	0.0	-	-	0.0	8.9	-
133.0	35.0	-	0.0	-	0.0	-	0.0	-	-	1.3	3.8	-
137.0	20.0	-	0.0	-	0.0	-	0.0	-	-	2.7	0.0	-
137.0	21.0	-	0.0	-	0.0	-	0.0	-	-	0.0	5.5	-
137.0	23.0	-	0.0	-	0.0	-	0.0	-	-	0.0	9.1	-
137.0	30.0	-	0.0	-	0.0	-	0.0	-	-	2.9	-	-
140.0	28.8	-	3.4	-	-	-	-	-	-	0.0	-	-
140.0	29.0	-	2.3	-	-	-	-	-	-	0.0	-	-

## Labridae

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
133.0	30.0	-	0.0	-	0.0	-	0.0	-	-	11.4	-	-
133.0	35.0	-	0.0	-	0.0	-	0.0	-	-	3.0	-	-
137.0	22.0	-	0.0	-	0.0	-	0.0	-	-	2.0	-	-
137.0	23.0	-	0.0	-	0.0	-	0.0	-	-	0.0	2.8	-
137.0	30.0	-	0.0	-	0.0	-	0.0	-	-	0.0	6.1	-
137.0	35.0	-	0.0	-	0.0	-	0.0	-	-	0.0	9.3	-

TABLE 4. (cont.)

## Labridae (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
	-	-	-	-	0.0	-	-	-	0.0	3.0	-	-
137.0	40.0	-	0.0	-	-	-	-	-	-	-	-	-

*Halichoeres spp.*

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
	-	-	-	-	-	-	-	-	-	-	-	-
117.0	24.8	-	0.0	-	0.0	-	-	-	5.7	-	0.0	-
117.0	26.0	-	0.0	-	0.0	-	-	-	0.0	-	7.9	-
117.0	40.0	-	0.0	-	0.0	-	-	-	0.0	-	9.3	-
117.0	45.0	-	0.0	-	0.0	-	-	-	0.0	-	34.7	-
118.0	39.0	-	0.0	-	0.0	-	-	-	5.6	-	136.0	-
120.0	22.5	-	0.0	-	0.0	-	-	-	17.5	-	4.7	-
120.0	23.0	-	0.0	-	0.0	-	-	-	10.3	-	5.6	-
120.0	24.0	-	0.0	-	0.0	-	-	-	1.2	-	4.8	-
120.0	25.0	-	0.0	-	0.0	-	-	-	0.0	-	7.0	-
120.0	30.0	-	0.0	-	0.0	-	-	-	0.0	-	9.0	-
120.0	35.0	-	0.0	-	0.0	-	-	-	2.6	-	2.7	-
120.0	40.0	-	0.0	-	0.0	-	-	-	5.2	-	2.7	-
123.5	35.6	-	0.0	-	0.0	-	-	-	0.0	-	4.3	-
127.0	32.6	-	0.0	-	0.0	-	-	-	0.0	-	1.7	-
127.0	33.0	-	0.0	-	0.0	-	-	-	0.0	-	5.2	-
127.0	45.0	-	0.0	-	0.0	-	-	-	3.2	-	0.0	-
130.0	25.5	-	0.0	-	0.0	-	-	-	8.1	-	0.0	-
130.0	26.0	-	0.0	-	0.0	-	-	-	11.6	-	27.8	-
130.0	28.0	-	0.0	-	0.0	-	-	-	5.0	-	4.7	-
130.0	30.0	-	0.0	-	0.0	-	-	-	0.0	-	18.1	-
130.0	45.0	-	0.0	-	0.0	-	-	-	0.0	-	3.0	-
130.0	55.0	-	0.0	-	0.0	-	-	-	0.0	-	3.1	-
133.0	21.0	-	0.0	-	0.0	-	-	-	0.0	-	18.3	-
133.0	23.0	-	0.0	-	0.0	-	-	-	0.0	-	12.5	-
133.0	25.0	-	0.0	-	0.0	-	-	-	0.0	-	29.4	-
137.0	21.0	-	0.0	-	0.0	-	-	-	1.4	-	6.5	-
137.0	22.0	-	0.0	-	0.0	-	-	-	2.1	-	0.0	-

*Oxyjulis californica*

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
	-	-	-	-	-	-	-	-	-	-	-	-
80.0	60.0	0.0	-	-	0.0	-	-	-	0.0	8.1	-	-
80.0	65.0	0.0	-	-	0.0	-	-	-	0.0	3.0	-	-
82.0	47.0	0.0	-	-	0.0	-	-	-	2.7	5.1	-	-
83.0	39.0	0.0	-	-	-	-	-	-	1.3	-	0.0	-
83.0	43.0	0.0	-	-	0.0	-	-	-	34.7	-	20.2	-
83.0	51.0	0.0	-	-	0.0	-	-	-	3.2	-	0.0	-
83.0	55.0	0.0	-	-	0.0	-	-	-	0.0	-	3.1	-
87.0	33.0	0.0	-	-	-	-	-	-	0.0	-	2.7	-

TABLE 4. (cont.)

*Oxyjuliss californica* (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
87.0	35.0	0.0	-	-	0.0	-	3.1	-	-	0.0	-	-
87.0	40.0	0.0	-	-	0.0	-	0.0	-	-	6.1	-	-
87.0	45.0	0.0	-	-	0.0	-	0.0	-	-	3.2	-	-
87.0	50.0	0.0	-	-	0.0	-	0.0	-	-	2.7	-	-
90.0	27.5	0.0	-	-	1.3	-	-	0.0	-	0.0	-	-
90.0	28.0	0.0	-	-	3.1	-	-	0.0	-	0.0	-	-
90.0	37.0	0.0	-	-	0.0	-	-	4.2	-	0.0	-	-
90.0	53.0	0.0	-	-	0.0	-	-	2.9	-	0.0	-	-
93.0	26.5	0.0	-	-	-	-	0.0	-	-	7.9	-	-
93.0	28.0	0.0	-	-	0.0	-	19.3	-	-	0.0	-	-
93.0	30.0	0.0	-	-	0.0	-	12.3	-	-	0.0	-	-
97.0	29.0	0.0	-	-	0.0	-	15.1	-	-	0.0	-	-
97.0	35.0	0.0	-	-	0.0	-	-	3.3	-	0.0	-	-
97.0	40.0	0.0	-	-	0.0	-	-	13.1	-	0.0	-	-
97.0	45.0	0.0	-	-	0.0	-	-	0.0	-	3.0	-	-
100.0	29.0	0.0	-	-	-	-	0.0	-	-	4.2	-	-
103.0	28.5	0.0	-	-	0.0	-	-	1.0	-	0.0	-	-
103.0	29.0	0.0	-	-	0.0	-	-	0.0	-	1.7	-	-
103.0	45.0	0.0	-	-	0.0	-	-	40.6	-	0.0	-	-
117.0	30.0	-	0.0	-	0.0	-	-	2.9	-	0.0	-	-
120.0	30.0	-	0.0	-	0.0	-	-	2.7	-	0.0	-	-

*Semicossyphus pulcher*

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
82.0	47.0	0.0	-	-	0.0	-	5.4	-	-	0.0	-	-
83.0	43.0	0.0	-	-	0.0	-	0.0	-	-	11.5	-	-
87.0	35.0	0.0	-	-	0.0	-	0.0	-	-	3.0	-	-
87.0	50.0	0.0	-	-	0.0	-	2.7	-	-	2.7	-	-
90.0	37.0	0.0	-	-	0.0	-	-	4.2	-	0.0	-	-
90.0	45.0	0.0	-	-	0.0	-	-	2.6	-	0.0	-	-
93.0	53.0	0.0	-	-	0.0	-	-	5.7	-	0.0	-	-
93.0	30.0	0.0	-	-	0.0	-	3.1	-	-	0.0	-	-
93.0	45.0	0.0	-	-	0.0	-	3.3	-	-	0.0	-	-
103.0	29.0	0.0	-	-	0.0	-	0.0	-	-	1.7	-	-
110.0	32.0	0.0	-	-	0.0	-	-	0.0	-	0.9	-	-
113.0	28.5	0.0	-	-	0.0	-	-	-	-	2.2	-	-
113.0	29.0	0.0	-	-	0.0	-	-	0.0	-	1.8	-	-
113.0	30.0	0.0	-	-	0.0	-	3.0	-	-	0.0	-	-
115.0	35.0	0.0	-	-	-	-	-	-	-	6.3	-	-
117.0	30.0	0.0	-	-	0.0	-	-	2.9	-	0.0	-	-
117.0	35.0	0.0	-	-	0.0	-	-	0.0	-	2.9	-	-
117.0	45.0	0.0	-	-	0.0	-	-	3.4	-	2.9	-	-
118.0	39.0	0.0	-	-	0.0	-	-	5.6	-	2.7	-	-
119.0	33.0	0.0	-	-	-	-	-	-	-	8.7	0.0	-

TABLE 4. (cont.)

*Semicossyphus pulcher* (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
120.0	22.5	-	0.0	-	0.0	-	-	1.0	-	0.0	-	-
120.0	35.0	-	0.0	-	0.0	-	-	5.2	-	0.0	-	-
127.0	50.0	-	0.0	-	0.0	-	-	2.9	-	0.0	-	-
130.0	26.0	-	0.0	-	0.0	-	-	0.0	-	2.3	-	-

## Pomacentridae

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
110.0	32.0	-	0.0	-	0.0	-	-	-	35.7	0.0	-	-
117.0	25.0	-	0.0	-	0.0	-	-	0.0	-	2.6	-	-
120.0	40.0	-	0.0	-	0.0	-	-	0.0	-	27.3	-	-
123.5	35.6	-	0.0	-	0.0	-	-	0.0	-	18.6	-	-
127.0	34.0	-	0.0	-	0.0	-	-	0.0	-	6.0	-	-
133.0	35.0	-	0.0	-	0.0	-	-	0.0	-	3.0	-	-
137.0	30.0	-	0.0	-	0.0	-	-	-	44.2	0.0	-	-
137.0	35.0	-	0.0	-	0.0	-	-	-	2.7	0.0	-	-

*Chromis punctipinnis*

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
83.0	43.0	0.0	-	0.0	-	0.0	-	-	-	37.4	-	-
87.0	35.0	0.0	-	0.0	-	0.0	-	-	-	6.0	-	-
87.0	40.0	0.0	-	0.0	-	0.0	-	-	-	18.3	-	-
90.0	37.0	0.0	-	0.0	-	-	-	18.8	-	0.0	-	-
93.0	26.5	0.0	-	0.0	-	0.0	-	-	-	6.3	-	-
93.0	28.0	0.0	-	0.0	-	0.0	-	-	-	16.0	-	-
93.0	30.0	0.0	-	0.0	-	0.0	-	-	-	29.5	-	-
97.0	32.0	0.0	-	0.0	-	0.0	-	-	-	37.0	-	-
97.0	40.0	0.0	-	0.0	-	0.0	-	-	-	3.0	-	-
97.0	45.0	0.0	-	0.0	-	0.0	-	-	-	3.0	-	-
97.0	75.0	-	0.0	-	0.0	-	-	-	-	6.5	-	-
100.0	30.0	0.0	-	0.0	-	0.0	-	-	-	44.0	-	-
100.0	35.0	0.0	-	0.0	-	0.0	-	-	-	6.1	-	-
100.0	45.0	0.0	-	0.0	-	0.0	-	-	-	3.1	-	-
103.0	29.0	0.0	-	0.0	-	0.0	-	-	-	1.7	-	-
103.0	40.0	0.0	-	0.0	-	0.0	-	-	-	21.9	-	-
107.0	32.0	-	0.0	-	0.0	-	-	-	-	18.0	-	-
110.0	32.0	-	0.0	-	0.0	-	-	-	-	3.4	-	-
110.0	35.0	-	0.0	-	0.0	-	-	-	-	8.6	-	-
113.0	30.0	-	0.0	-	0.0	-	-	-	-	31.2	-	-
113.0	35.0	-	0.0	-	0.0	-	-	-	-	3.0	-	-
113.0	40.0	-	0.0	-	0.0	-	-	-	-	3.1	-	-
117.0	26.0	-	0.0	-	0.0	-	-	-	-	7.9	-	-
117.0	35.0	-	0.0	-	0.0	-	-	-	-	8.8	-	-

TABLE 4. (cont.)

*Chromis punctipinnis* (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
117.0	45.0	-	0.0	-	0.0	-	-	0.0	-	2.9	-	-
117.0	60.0	-	0.0	-	0.0	-	-	3.8	-	0.0	-	-
118.0	39.0	-	0.0	-	3.1	-	-	0.0	-	16.3	-	-
119.0	33.0	-	0.0	-	0.0	-	-	0.0	-	6.5	-	-
120.0	40.0	-	0.0	-	0.0	-	-	0.0	-	49.1	-	-
123.0	36.0	-	0.0	-	0.0	-	-	0.0	-	35.0	-	-
123.0	37.0	-	0.0	-	0.0	-	-	0.0	-	164.4	-	-
123.0	42.0	-	0.0	-	0.0	-	-	0.0	-	0.0	-	-
127.0	32.6	-	0.0	-	0.0	-	-	3.2	-	2.6	-	-
127.0	33.0	-	0.0	-	0.0	-	-	0.0	-	52.0	-	-

*Hypsypops rubicundus*

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
93.0	26.5	0.0	-	-	-	-	-	0.0	-	-	600.4	-
93.0	27.0	0.0	-	0.0	-	-	-	0.0	-	-	17.4	-
93.0	28.0	0.0	-	0.0	-	-	-	0.0	-	-	21.4	-
93.0	30.0	0.0	-	0.0	-	-	-	0.0	-	-	8.9	-
97.0	32.0	0.0	-	0.0	-	-	-	0.0	-	-	12.3	-
107.0	30.0	-	0.0	-	0.0	-	-	0.0	-	-	0.6	-
107.0	31.0	-	0.0	-	0.0	-	-	0.0	-	-	7.1	-
110.0	32.0	-	0.0	-	0.0	-	-	0.0	-	-	12.0	-

*Mugil* spp.

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
133.0	35.0	-	0.0	-	0.0	-	-	0.0	-	3.0	-	-

*Howella brodiei*

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
60.0	180.0	3.0	-	-	-	-	-	0.0	-	-	-	-
90.0	120.0	0.0	-	-	-	-	-	2.7	-	-	-	-
90.0	140.0	0.0	-	-	-	-	-	2.7	-	-	-	-
90.0	160.0	0.0	-	-	-	-	-	2.6	-	-	-	-

*Brama* spp.

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
60.0	140.0	0.0	-	-	-	-	-	2.6	-	-	-	-
60.0	200.0	15.5	-	-	-	-	-	0.0	-	-	-	-
80.0	160.0	5.8	-	-	-	-	-	-	-	-	-	-

TABLE 4. (cont.)

*Brama spp.* (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
80.0	180.0	2.7	-	-	-	-	-	-	-	-	-	-
90.0	160.0	2.6	-	-	-	-	-	2.6	-	-	-	-
90.0	180.0	2.7	-	-	-	-	-	0.0	-	-	-	-
90.0	200.0	2.8	-	-	-	-	-	2.8	-	-	-	-

## Carangidae

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
120.0	22.5	-	0.0	-	0.0	-	-	19.6	-	0.0	-	-
120.0	23.0	-	0.0	-	0.0	-	-	14.7	-	0.0	-	-
130.0	25.5	-	0.0	-	0.0	-	-	12.1	34.6	-	-	-
130.0	26.0	-	0.0	-	0.0	-	-	16.6	55.7	-	-	-
130.0	28.0	-	0.0	-	0.0	-	-	0.0	71.1	-	-	-
130.0	30.0	-	0.0	-	0.0	-	-	0.0	5.2	-	-	-
130.0	35.0	-	0.0	-	0.0	-	-	0.0	6.1	-	-	-
133.0	19.5	-	0.0	-	0.0	-	-	0.0	1.2	-	-	-
133.0	21.0	-	0.0	-	0.0	-	-	8.1	41.2	-	-	-
133.0	23.0	-	0.0	-	0.0	-	-	0.0	2.5	-	-	-
133.0	25.0	-	0.0	-	0.0	-	-	0.0	17.6	-	-	-
133.0	30.0	-	0.0	-	0.0	-	-	3.0	2.9	-	-	-
137.0	20.0	-	0.0	-	0.0	-	-	44.1	0.0	-	-	-
137.0	21.0	-	0.0	-	0.0	-	-	102.8	49.7	-	-	-
137.0	22.0	-	0.0	-	0.0	-	-	81.3	53.0	-	-	-
137.0	23.0	-	0.0	-	0.0	-	-	26.6	0.0	-	-	-
140.0	28.8	-	0.0	-	-	-	-	0.5	-	-	-	-
140.0	29.0	-	0.0	-	-	-	-	25.5	-	-	-	-
140.0	30.0	-	0.0	-	-	-	-	2.8	-	-	-	-

*Seriola lalandi*

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
80.0	60.0	0.0	-	-	0.0	-	-	19.6	0.0	-	-	-
87.0	33.0	0.0	-	-	0.0	-	-	0.0	-	2.6	-	-
120.0	35.0	-	0.0	-	0.0	-	-	0.0	-	2.7	-	-
120.0	40.0	-	0.0	-	0.0	-	-	0.0	-	5.5	-	-
120.0	45.0	-	0.0	-	0.0	-	-	0.0	-	3.0	-	-
120.0	50.0	-	0.0	-	0.0	-	-	0.0	-	0.0	-	-
123.0	36.0	-	0.0	-	0.0	-	-	0.0	-	9.5	-	-
123.0	45.0	-	0.0	-	0.0	-	-	0.0	-	0.0	-	-
123.0	50.0	-	0.0	-	0.0	-	-	0.0	-	0.0	-	-
123.5	35.6	-	0.0	-	0.0	-	-	0.0	-	22.9	-	-
130.0	50.0	-	0.0	-	0.0	-	-	0.0	-	0.0	-	-
133.0	70.0	-	0.0	-	3.0	-	-	0.0	-	0.4	0.0	-
137.0	20.0	-	0.0	-	-	-	-	-	-	-	-	-

TABLE 4. (cont.)

*Trachurus symmetricus*

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
60.0	55.0	0.0									5.2	
60.0	70.0	0.0									14.1	
60.0	80.0	0.0									12.2	
60.0	120.0	0.0									2.7	
60.0	140.0	0.0									5.2	
70.0	60.0	0.0			0.0						13.3	
70.0	70.0	0.0			0.0						30.7	
70.0	80.0	0.0			0.0						25.3	
70.0	90.0	0.0			0.0						2.8	
70.0	100.0	0.0			0.0						5.6	
80.0	65.0	0.0			0.0						118.8	0.0
80.0	70.0	0.0			0.0						125.9	0.0
83.0	60.0	0.0			0.0						0.0	
83.0	65.0	0.0			0.0						13.2	
83.0	90.0	0.0			0.0						19.3	
87.0	33.0	0.0			0.0						2.7	
87.0	34.0	0.0			0.0						5.3	
87.0	40.0	0.0			0.0						0.0	
87.0	45.0	0.0			0.0						0.0	
87.0	55.0	0.0			0.0						6.5	
87.0	60.0	0.0			0.0						13.0	
87.0	65.0	0.0			0.0						32.0	
87.0	70.0	0.0			0.0						443.0	24.3
87.0	80.0	0.0			0.0						24.1	19.6
87.0	90.0	0.0			0.0						2.8	12.5
87.0	100.0	0.0			0.0						3.3	
90.0	60.0	0.0			16.1						0.0	
90.0	65.0	0.0			8.8						0.0	
90.0	70.0	0.0			34.2						0.0	
90.0	80.0	0.0			35.4						2.8	
90.0	85.0	0.0			44.8						2.6	
90.0	90.0	0.0			15.6						0.0	
93.0	28.0	0.0			0.0						6.4	
93.0	30.0	0.0			0.0						0.0	
93.0	50.0	0.0			0.0						16.5	
93.0	55.0	0.0			0.0						66.0	
93.0	60.0	0.0			0.0						99.9	
93.0	65.0	0.0			34.7						19.5	
93.0	70.0	0.0			8.8						10.0	
93.0	75.0	0.0			10.0						3.3	
93.0	80.0	3.0			-						-	
93.0	85.0	-			139.5						6.3	
93.0	90.0	-			-						-	
93.0	100.0	-			137.8						25.1	
97.0	35.0	0.0			-						101.9	
97.0	-	-			0.0						-	32.5

TABLE 4. (cont.)

*Trachurus symmetricus* (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
97.0	40.0	0.0	-	3.2	-	-	196.2	-	-	0.0	-	-
97.0	45.0	0.0	-	3.5	-	-	25.4	-	-	0.0	-	-
97.0	50.0	0.0	-	0.0	-	-	33.2	-	-	8.9	-	-
97.0	55.0	0.0	-	-	-	-	9.1	-	-	0.0	-	-
97.0	60.0	0.0	-	-	-	-	6.3	-	-	0.0	-	-
97.0	65.0	0.0	-	13.8	-	-	32.5	-	-	0.0	-	-
97.0	70.0	0.0	-	-	-	-	21.6	-	-	2.9	-	-
97.0	75.0	-	-	52.9	-	-	-	-	-	0.0	-	-
97.0	80.0	0.0	-	-	-	-	25.8	-	-	92.8	-	-
97.0	85.0	-	-	526.3	-	-	-	-	-	-	-	-
97.0	90.0	0.0	-	-	-	-	12.4	-	-	-	-	-
97.0	95.0	-	-	32.7	-	-	-	-	-	-	-	-
97.0	100.0	-	-	28.0	-	-	-	-	-	0.0	-	-
100.0	129.0	0.0	-	-	-	-	2.9	-	-	0.0	-	-
100.0	30.0	0.0	-	-	-	-	3.2	-	-	0.0	-	-
100.0	35.0	0.0	-	-	-	-	6.3	-	-	2.8	-	-
100.0	40.0	0.0	-	-	-	-	6.4	-	-	19.3	-	-
100.0	45.0	0.0	-	-	-	-	43.1	-	-	2.9	-	-
100.0	50.0	0.0	-	-	-	-	12.0	-	-	8.8	-	-
100.0	55.0	0.0	-	-	-	-	39.3	-	-	0.0	-	-
100.0	60.0	0.0	-	26.8	-	-	23.4	-	-	0.0	-	-
100.0	65.0	20.9	-	-	-	-	98.9	-	-	0.0	-	-
100.0	70.0	37.0	-	-	-	-	230.5	-	-	2.3	-	-
100.0	75.0	-	-	-	-	-	225.8	-	-	-	-	-
100.0	80.0	0.0	-	-	-	-	63.5	-	-	0.0	-	-
100.0	85.0	-	-	-	-	-	607.2	-	-	-	-	-
100.0	90.0	0.0	-	-	-	-	360.8	-	-	2.9	-	-
100.0	100.0	-	-	0.0	-	-	-	-	-	3.0	-	-
100.0	120.0	-	-	0.0	-	-	-	-	-	-	-	-
103.0	40.0	11.7	-	-	-	-	6.1	-	-	0.0	-	-
103.0	45.0	0.0	-	-	-	-	42.4	-	-	0.0	-	-
103.0	50.0	5.8	-	-	-	-	26.3	-	-	29.4	-	-
103.0	55.0	57.4	-	-	-	-	0.0	-	-	6.7	-	-
103.0	60.0	158.1	-	-	-	-	6.7	-	-	0.0	-	-
103.0	65.0	99.0	-	-	-	-	25.2	-	-	6.6	-	-
103.0	70.0	117.4	-	-	-	-	177.3	-	-	3.4	-	-
103.0	75.0	-	-	-	-	-	140.4	-	-	-	-	-
103.0	80.0	-	-	0.0	-	-	101.0	-	-	0.0	-	-
103.0	85.0	-	-	-	-	-	353.8	-	-	-	-	-
103.0	90.0	-	-	0.0	-	-	170.1	-	-	10.0	-	-
107.0	30.0	-	-	0.0	-	-	21.4	-	-	0.0	-	-
107.0	31.0	-	-	0.0	-	-	5.7	-	-	0.0	-	-
107.0	32.0	-	-	0.0	-	-	3.1	-	-	0.0	-	-
107.0	40.0	-	-	2.9	-	-	0.0	-	-	0.0	-	-
107.0	45.0	-	-	0.0	-	-	41.6	-	-	3.5	-	-
107.0	50.0	-	-	2.9	-	-	24.1	-	-	0.0	-	-

TABLE 4. (cont.)

*Trachurus symmetricus* (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
107.0	55.0	-	0.0	-	28.6	-	3.3	-	-	6.1	-	-
107.0	60.0	-	54.5	-	6.5	-	0.0	-	-	0.0	-	-
107.0	65.0	-	5.9	-	3.1	-	3.2	-	-	3.3	-	-
107.0	70.0	-	232.1	-	33.7	0.0	-	-	-	9.1	-	-
107.0	75.0	-	-	-	9.2	-	-	-	-	-	-	-
107.0	80.0	-	0.0	-	8.9	0.0	-	-	-	-	-	-
107.0	85.0	-	-	-	55.3	-	-	-	-	-	-	-
107.0	90.0	-	0.0	-	327.4	13.4	-	-	-	-	-	-
110.0	32.0	-	0.0	-	2.4	-	-	-	-	0.0	0.0	-
110.0	40.0	-	0.0	-	2.8	-	-	-	-	16.9	0.0	-
110.0	45.0	-	0.0	-	0.0	-	-	-	-	7.7	0.0	-
110.0	50.0	-	5.6	-	5.6	-	-	-	-	0.0	0.0	-
110.0	55.0	-	49.1	-	3.0	-	-	-	-	0.0	0.0	-
110.0	60.0	-	0.0	-	9.1	-	-	-	-	0.0	0.0	-
110.0	65.0	-	37.0	-	0.0	-	-	-	-	0.0	0.0	-
110.0	70.0	-	92.4	-	0.0	-	-	-	-	0.0	0.0	-
110.0	80.0	-	23.1	-	0.0	-	-	-	-	0.0	-	-
110.0	85.0	-	-	-	9.2	-	-	-	-	0.0	-	-
110.0	90.0	-	0.0	-	15.4	-	-	-	-	0.0	-	-
113.0	40.0	-	0.0	-	6.5	0.0	-	-	-	0.0	-	-
113.0	45.0	-	0.0	-	24.9	0.0	-	-	-	0.0	-	-
113.0	50.0	-	0.0	-	21.6	3.2	-	-	-	0.0	-	-
113.0	55.0	-	0.0	-	3.6	14.8	-	-	-	0.0	-	-
113.0	60.0	-	0.0	-	3.2	0.0	-	-	-	0.0	-	-
113.0	65.0	-	0.0	-	36.2	6.1	-	-	-	0.0	-	-
113.0	70.0	-	0.0	-	40.0	10.2	-	-	-	0.0	-	-
113.0	75.0	-	-	-	3.3	-	-	-	-	-	-	-
113.0	80.0	-	14.2	-	9.3	0.0	-	-	-	-	-	-
113.0	85.0	-	-	-	6.0	-	-	-	-	-	-	-
113.0	90.0	-	2.8	-	17.0	0.0	-	-	-	6.7	-	-
117.0	45.0	-	0.0	-	0.0	-	-	-	-	0.0	-	-
117.0	55.0	-	18.4	-	0.0	-	-	-	-	3.8	-	-
117.0	60.0	-	0.0	-	0.0	-	-	-	-	3.1	-	-
117.0	70.0	-	3.1	-	3.0	-	-	-	-	-	-	-
117.0	75.0	-	-	-	27.5	3.3	-	-	-	-	-	-
117.0	80.0	-	3.0	-	-	-	-	-	-	11.2	-	-
117.0	85.0	-	-	-	8.7	-	-	-	-	0.0	-	-
117.0	90.0	-	2.9	-	16.9	0.0	-	-	-	0.0	-	-
118.0	39.0	-	0.0	-	-	-	-	-	-	2.8	-	-
120.0	45.0	-	0.0	-	3.1	-	-	-	-	0.0	-	-
120.0	60.0	-	0.0	-	0.0	-	-	-	-	0.0	-	-
120.0	65.0	-	0.0	-	3.1	6.6	-	-	-	0.0	-	-
120.0	70.0	-	0.0	-	0.0	-	-	-	-	0.0	-	-
120.0	75.0	-	-	-	-	-	-	-	-	-	-	-

TABLE 4. (cont.)

*Caristius macropus*

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
103.0	65.0	0.0	-	-	0.0	-	0.0	-	-	3.2	-	-

*Coryphaena hippurus*

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
117.0	26.0	-	0.0	-	0.0	-	-	2.7	-	0.0	-	-
127.0	40.0	-	0.0	-	0.0	-	-	2.8	0.0	0.0	-	-
133.0	30.0	-	0.0	-	0.0	-	-	0.0	-	2.9	-	-
133.0	35.0	-	0.0	-	0.0	-	-	0.0	-	8.9	-	-
137.0	22.0	-	0.0	-	0.0	-	-	-	2.1	0.0	-	-
137.0	23.0	-	0.0	-	2.6	-	-	0.0	-	0.0	-	-
137.0	30.0	-	0.0	-	0.0	-	-	-	0.0	3.0	-	-
137.0	35.0	-	0.0	-	0.0	-	-	-	0.0	3.1	-	-
137.0	50.0	-	0.0	-	0.0	-	-	-	2.6	0.0	-	-
140.0	50.0	-	0.0	-	-	-	-	-	2.3	-	-	-

## Gerreidae

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
127.0	32.6	-	0.0	-	0.0	-	-	0.0	-	0.9	-	-
130.0	25.5	-	0.0	-	0.0	-	-	-	34.0	0.0	-	-
130.0	26.0	-	0.0	-	0.0	-	-	-	18.3	0.0	-	-
133.0	21.0	-	0.0	-	0.0	-	-	0.0	-	45.8	-	-
133.0	23.0	-	0.0	-	0.0	-	-	0.0	-	7.5	-	-
137.0	21.0	-	0.6	-	0.0	-	-	-	24.7	30.2	-	-
137.0	22.0	-	0.0	-	0.0	-	-	-	30.0	24.5	-	-
137.0	23.0	-	0.0	-	0.0	-	-	-	13.3	8.3	-	-
140.0	28.8	-	0.0	-	-	-	-	-	2.4	-	-	-
140.0	29.0	-	0.0	-	-	-	-	-	9.8	-	-	-

## Haemulidae

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
120.0	22.5	-	0.0	-	0.0	-	-	0.0	-	6.3	-	-
130.0	25.5	-	0.0	-	0.0	-	-	-	868.3	0.0	-	-
130.0	26.0	-	0.0	-	0.0	-	-	-	410.0	0.0	-	-
130.0	30.0	-	0.0	-	0.0	-	-	-	0.0	2.6	-	-
133.0	21.0	-	0.0	-	0.0	-	-	0.0	-	6.9	-	-
137.0	20.0	-	0.0	-	0.0	-	-	-	11.3	0.0	-	-
137.0	21.0	-	0.0	-	0.0	-	-	-	263.0	30.2	-	-
137.0	22.0	-	0.0	-	0.0	-	-	-	297.5	6.1	-	-
137.0	23.0	-	0.0	-	0.0	-	-	-	72.2	0.0	-	-

TABLE 4. (cont.)

<i>Girella nigricans</i>												
STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
93.0	26.5	0.0	-	-	-	-	1.3	-	0.0	-	-	-
117.0	26.0	-	0.0	-	0.0	-	-	2.7	-	0.0	-	-
120.0	40.0	-	0.0	-	0.0	-	-	0.0	-	2.7	-	-
<i>Medialuna californiensis</i>												
STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
83.0	90.0	0.0	-	-	-	-	3.2	-	-	-	-	-
100.0	40.0	0.0	-	-	0.0	-	-	2.8	-	0.0	-	-
103.0	65.0	0.0	-	-	0.0	-	0.0	-	-	3.2	-	-
107.0	55.0	-	0.0	-	0.0	-	0.0	-	-	3.1	-	-
133.0	50.0	-	0.0	-	0.0	-	-	2.6	-	0.0	-	-
<i>Caulolatilus princeps</i>												
STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
107.0	40.0	-	0.0	-	0.0	-	0.0	-	-	-	-	-
117.0	70.0	-	0.0	-	0.0	-	3.1	-	-	-	-	-
127.0	50.0	-	0.0	-	0.0	-	-	5.8	-	0.0	-	-
127.0	55.0	-	0.0	-	0.0	-	-	2.7	-	0.0	-	-
130.0	35.0	-	0.0	-	0.0	-	-	-	2.5	0.0	-	-
130.0	45.0	-	0.0	-	0.0	-	-	-	2.8	0.0	-	-
137.0	21.0	-	0.0	-	0.0	-	-	-	1.4	0.0	-	-
<i>Sciaenidae</i>												
STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
60.0	49.5	19.8	-	-	-	-	-	-	-	-	-	-
60.0	50.0	13.0	-	-	-	-	-	-	-	0.0	-	-
60.0	51.0	55.1	-	-	-	-	-	-	-	0.0	-	-
60.0	52.0	29.3	-	-	-	-	-	-	-	0.0	-	-
60.0	55.0	6.2	-	-	-	-	-	-	-	0.0	-	-
63.0	49.7	45.8	-	-	-	-	4.4	-	-	-	-	-
63.0	50.0	169.7	-	-	-	-	0.8	-	-	0.0	-	-
63.0	51.0	76.3	-	-	-	-	0.0	-	-	0.0	-	-
63.0	52.0	26.1	-	-	-	-	0.0	-	-	0.0	-	-
67.0	47.3	14.3	-	-	-	-	0.0	-	-	-	-	-
67.0	50.0	2.3	-	-	-	-	0.0	-	-	0.0	-	-
67.0	55.0	3.3	-	-	-	-	0.0	-	-	0.0	-	-
73.0	49.4	0.6	-	-	-	-	0.0	-	-	0.0	-	-
73.0	50.0	16.2	-	-	-	-	5.5	-	-	0.0	-	-
73.0	51.0	15.4	-	-	-	-	0.0	-	-	0.0	-	-
73.0	53.0	2.7	-	-	-	-	0.0	-	-	0.0	-	-

TABLE 4. (cont.)

## Sciaenidae (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
77.0	47.6	46.8	-	-	52.3	-	-	0.0	-	-	-	-
77.0	48.0	138.9	-	-	49.0	-	-	0.0	-	-	-	-
77.0	49.0	-	-	-	0.7	-	-	0.0	-	-	-	-
77.0	55.0	0.0	-	-	7.2	-	-	0.0	-	-	-	-
77.0	57.0	0.0	-	-	2.5	-	-	0.0	-	-	-	-
80.0	50.2	28.8	-	-	23.3	-	-	0.0	0.0	-	-	-
80.0	51.0	34.8	-	-	84.2	-	-	0.0	0.0	-	-	-
80.0	52.0	0.0	-	-	149.8	-	-	0.0	2.7	-	-	-
80.0	55.0	0.0	-	-	9.6	-	-	0.0	0.0	-	-	-
83.0	39.0	36.8	-	-	-	0.0	-	-	-	3.9	-	-
83.0	40.0	43.9	-	-	0.8	-	-	6.2	-	-	-	-
83.0	43.0	3.0	-	-	0.0	-	-	0.0	-	-	-	-
83.0	51.0	2.4	-	-	2.6	-	-	0.0	-	-	-	-
87.0	32.5	214.9	-	-	6.6	-	-	1.7	-	-	-	-
87.0	33.0	1158.2	-	-	7.9	-	-	10.8	-	-	-	-
87.0	34.0	181.6	-	-	0.0	-	-	10.6	-	-	-	-
87.0	35.0	29.4	-	-	0.0	-	-	0.0	-	-	-	-
87.0	40.0	0.0	-	-	0.0	-	-	0.0	-	-	-	-
90.0	27.5	138.5	-	-	7.1	-	-	0.8	-	-	-	-
90.0	28.0	535.7	-	-	3.1	-	-	0.0	-	-	-	-
90.0	32.0	132.3	-	-	0.0	-	-	0.0	-	-	-	-
90.0	37.0	26.9	-	-	-	-	-	0.0	-	-	-	-
93.0	26.5	69.6	-	-	15.8	-	-	2.5	-	-	-	-
93.0	27.0	14.1	-	-	3.3	-	-	60.3	-	-	-	-
93.0	28.0	0.0	-	-	5.7	-	-	12.8	-	-	-	-
93.0	30.0	0.0	-	-	-	-	-	6.1	-	-	-	-
96.8	28.7	228.9	-	-	-	-	-	2.6	-	-	-	-
97.0	29.0	170.2	-	-	0.0	-	-	0.0	-	-	-	-
97.0	30.0	19.4	-	-	7.6	-	-	0.0	-	-	-	-
97.0	32.0	0.0	-	-	24.8	-	-	0.0	-	-	-	-
100.0	29.0	33.8	-	-	5.8	-	-	0.0	-	-	-	-
100.0	30.0	89.9	-	-	0.0	-	-	0.0	-	-	-	-
103.0	28.5	13.8	-	-	6.5	-	-	5.7	-	-	-	-
103.0	29.0	11.4	-	-	3.1	-	-	0.0	-	-	-	-
103.0	30.0	2.5	-	-	3.2	-	-	5.3	-	-	-	-
107.0	30.0	-	-	-	10.5	-	-	0.0	-	-	-	-
107.0	31.0	-	-	-	6.5	-	-	0.0	-	-	-	-
107.0	32.0	-	-	-	3.1	-	-	0.0	-	-	-	-
107.0	35.0	-	-	-	3.2	-	-	0.0	-	-	-	-
110.0	32.0	-	-	-	4.7	-	-	0.0	-	-	-	-
113.0	28.5	-	-	-	3.5	-	-	6.7	-	-	-	-
113.0	29.0	-	-	-	19.4	-	-	0.0	-	-	-	-
117.0	24.8	-	-	-	1.0	-	-	0.0	-	-	-	-
117.0	25.0	-	-	-	0.0	-	-	0.0	-	-	-	-
117.0	26.0	-	-	-	0.0	-	-	0.0	-	-	-	-
118.0	39.0	-	-	-	3.1	-	-	5.6	-	-	-	-

TABLE 4. (cont.)

## Sciaenidae (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
120.0	23.0	-	2.5	-	0.0	-	-	0.0	-	1.9	-	-
120.0	24.0	-	310.1	-	0.0	-	-	0.0	-	0.0	-	-
120.0	25.0	-	65.9	-	0.0	-	-	2.0	-	0.0	-	-
120.0	35.0	-	0.0	-	0.0	-	-	2.6	-	0.0	-	-
120.0	40.0	-	0.0	-	0.0	-	-	18.1	-	0.0	-	-
123.0	37.0	-	0.0	-	7.4	-	-	0.0	-	0.0	-	-
123.5	35.6	-	0.0	-	0.0	-	-	2.5	-	0.0	-	-
127.0	32.6	-	0.0	-	0.0	-	-	1.4	-	0.0	-	-
127.0	40.0	-	1.4	-	0.0	-	-	0.0	-	0.0	-	-
130.0	25.5	-	0.0	-	0.0	-	-	652.9	-	5.8	-	-
130.0	26.0	-	0.0	-	0.0	-	-	-	-	21.6	-	-
130.0	28.0	-	0.0	-	0.0	-	-	-	-	12.4	-	-
130.0	30.0	-	0.0	-	2.9	-	-	-	-	4.0	-	-
133.0	19.5	-	1.7	-	0.0	-	-	0.0	-	2.4	-	-
133.0	21.0	-	4.0	-	0.0	-	-	14.1	-	9.2	-	-
133.0	23.0	-	0.0	-	2.6	-	-	0.0	-	0.0	-	-
133.0	30.0	-	0.0	-	0.0	-	-	8.9	-	0.0	-	-
133.0	50.0	-	0.0	-	0.0	-	-	2.6	-	0.0	-	-
133.0	55.0	-	0.0	-	0.0	-	-	5.3	-	-	-	-
137.0	20.0	-	0.3	-	0.0	-	-	15.1	-	0.0	-	-
137.0	21.0	-	0.0	-	0.0	-	-	-	-	242.5	-	-
137.0	22.0	-	0.0	-	0.0	-	-	-	-	143.4	-	-
137.0	23.0	-	0.0	-	5.3	-	-	-	-	41.8	-	-
140.0	28.8	-	0.7	-	-	-	-	-	-	5.5	-	-
140.0	29.0	-	2.3	-	-	-	-	-	-	0.5	-	-

## Serraniidae

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
80.0	52.0	0.0	-	-	0.0	-	-	0.0	-	5.3	-	-
83.0	39.0	0.0	-	-	0.0	-	-	0.0	-	1.9	-	-
83.0	43.0	0.0	-	-	0.0	-	-	-	-	14.4	-	-
83.0	55.0	0.0	-	-	0.0	-	-	-	-	3.1	-	-
87.0	32.5	0.0	-	-	0.0	-	-	0.0	-	10.0	-	-
87.0	34.0	0.0	-	-	0.0	-	-	0.0	-	5.3	-	-
87.0	35.0	0.0	-	-	0.0	-	-	0.0	-	6.0	-	-
87.0	40.0	0.0	-	-	0.0	-	-	0.0	-	6.1	-	-
90.0	45.0	0.0	-	-	0.0	-	-	-	-	0.0	-	-
93.0	26.5	0.0	-	-	-	-	-	-	-	15.4	-	-
93.0	27.0	0.0	-	-	-	-	-	-	-	23.9	-	-
93.0	28.0	0.0	-	-	-	-	-	-	-	37.3	-	-
93.0	30.0	0.0	-	-	-	-	-	-	-	0.0	-	-
97.0	30.0	0.0	-	-	-	-	-	-	-	6.1	-	-
113.0	28.5	-	0.0	-	-	-	-	-	-	2.8	-	-
117.0	30.0	-	0.0	-	-	-	-	-	-	0.0	-	-

TABLE 4. (cont.)

## Serranidae (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
117.0	35.0	-	0.0	-	0.0	-	-	0.0	-	2.9	-	-
118.0	39.0	-	0.0	-	0.0	-	-	0.0	-	16.3	-	-
119.0	33.0	-	0.0	-	0.0	-	-	0.0	-	4.7	-	-
120.0	22.5	-	0.0	-	0.0	-	-	0.0	-	5.6	-	-
120.0	23.0	-	0.0	-	0.0	-	-	0.0	-	2.4	-	-
120.0	24.0	-	0.0	-	0.0	-	-	0.0	-	19.1	-	-
120.0	40.0	-	0.0	-	0.0	-	-	0.0	-	3.2	-	-
123.0	36.0	-	0.0	-	0.0	-	-	0.0	-	0.0	-	-
123.0	42.0	-	0.0	-	0.0	-	-	0.0	-	0.0	-	-
123.5	35.6	-	0.0	-	0.0	-	-	0.0	-	4.3	-	-
127.0	45.0	-	0.0	-	0.0	-	-	0.0	-	0.0	-	-
127.0	50.0	-	0.0	-	0.0	-	-	2.9	-	0.0	-	-
130.0	25.5	-	0.0	-	0.0	-	-	1157.5	-	11.5	-	-
130.0	26.0	-	0.0	-	0.0	-	-	105.4	-	4.6	-	-
130.0	28.0	-	0.0	-	0.0	-	-	5.0	-	9.5	-	-
130.0	30.0	-	0.0	-	0.0	-	-	0.0	-	41.3	-	-
133.0	21.0	-	0.0	-	0.0	-	-	0.0	-	116.8	-	-
133.0	23.0	-	0.0	-	0.0	-	-	0.0	-	10.0	-	-
133.0	25.0	-	0.0	-	0.0	-	-	0.0	-	2.9	-	-
133.0	30.0	-	0.0	-	0.0	-	-	0.0	-	94.4	-	-
133.0	35.0	-	0.0	-	0.0	-	-	0.0	-	8.9	-	-
133.0	45.0	-	2.9	-	0.0	-	-	0.0	-	0.0	-	-
133.0	50.0	-	2.8	-	0.0	-	-	0.0	-	3.0	-	-
137.0	21.0	-	0.0	-	0.0	-	-	515.1	-	32.4	-	-
137.0	22.0	-	0.0	-	0.0	-	-	532.9	-	14.3	-	-
137.0	23.0	-	0.0	-	0.0	-	-	112.1	-	13.8	-	-
137.0	30.0	-	0.0	-	0.0	-	-	0.0	-	12.2	-	-
137.0	35.0	-	0.0	-	0.0	-	-	0.0	-	6.2	-	-
137.0	40.0	-	0.0	-	0.0	-	-	0.0	-	3.0	-	-
140.0	28.8	-	0.0	-	0.0	-	-	1.0	-	-	-	-
140.0	29.0	-	0.0	-	0.0	-	-	68.6	-	-	-	-
140.0	30.0	-	0.0	-	0.0	-	-	17.1	-	-	-	-
140.0	35.0	-	0.0	-	0.0	-	-	2.7	-	-	-	-

## Gempylidae

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
70.0	200.0	0.0	-	-	-	-	-	8.0	-	-	-	-
80.0	160.0	2.9	-	-	-	-	-	2.8	-	-	-	-
80.0	200.0	0.0	-	-	-	-	-	0.0	-	-	-	-
90.0	200.0	2.8	-	-	-	-	-	0.0	-	-	-	-
93.0	90.0	3.2	-	-	-	-	-	0.0	-	-	0.0	3.0
137.0	45.0	-	0.0	-	0.0	-	-	-	-	2.8	-	-
140.0	45.0	-	0.0	-	0.0	-	-	-	-	2.3	-	-
140.0	50.0	-	0.0	-	0.0	-	-	-	-	-	-	-

TABLE 4. (cont.)

*Scombridae*

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
120.0	22.5	-	0.0	-	0.0	-	-	7.2	-	0.0	-	-
120.0	50.0	-	0.0	-	0.0	-	-	3.0	-	0.0	-	-
133.0	30.0	-	0.0	-	0.0	-	-	3.0	-	0.0	-	-
133.0	55.0	-	0.0	-	0.0	-	-	2.7	-	0.0	-	-

*Auxis spp.*

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
130.0	35.0	-	0.0	-	3.1	-	-	5.2	0.0	3.0	-	-
133.0	50.0	-	0.0	-	0.0	-	-	-	56.2	0.0	-	-
137.0	21.0	-	0.0	-	0.0	-	-	-	100.6	0.0	-	-
137.0	22.0	-	0.0	-	0.0	-	-	-	22.8	0.0	-	-
137.0	23.0	-	0.0	-	0.0	-	-	-	0.5	-	-	-
140.0	28.8	-	0.0	-	-	-	-	-	3.9	-	-	-
140.0	29.0	-	0.0	-	-	-	-	-	-	-	-	-

*Sarda chiliensis*

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
120.0	50.0	-	0.0	-	23.8	-	-	0.0	0.0	-	-	-
123.0	45.0	-	0.0	-	8.7	-	-	0.0	0.0	-	-	-
130.0	40.0	-	0.0	-	6.3	-	-	0.0	0.0	-	-	-
133.0	35.0	-	0.0	-	12.4	-	-	0.0	0.0	-	-	-
133.0	50.0	-	0.0	-	8.9	-	-	0.0	0.0	-	-	-
133.0	55.0	-	0.0	-	3.0	-	-	0.0	0.0	-	-	-
137.0	40.0	-	0.0	-	21.0	-	-	0.0	0.0	-	-	-
137.0	60.0	-	0.0	-	3.1	-	-	0.0	0.0	-	-	-
140.0	29.0	-	0.0	-	-	-	-	-	7.8	-	-	-

*Scomber japonicus*

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
87.0	33.0	0.0	-	-	0.0	-	-	0.0	-	2.6	-	-
87.0	34.0	0.0	-	-	0.0	-	-	0.0	-	2.6	-	-
87.0	35.0	0.0	-	-	0.0	-	-	3.1	-	18.1	-	-
87.0	40.0	0.0	-	-	0.0	-	-	0.0	-	3.0	-	-
93.0	26.5	0.0	-	-	-	-	-	8.8	-	0.0	-	-
93.0	27.0	0.0	-	-	0.0	-	-	5.7	-	0.0	-	-
93.0	30.0	0.0	-	-	0.0	-	-	3.1	-	0.0	-	-
103.0	80.0	-	0.0	-	3.0	-	-	0.0	-	2.6	-	-
115.0	35.0	-	0.0	-	-	-	-	-	-	11.8	-	-
117.0	30.0	-	0.0	-	0.0	-	-	0.0	-	3.4	-	-
117.0	45.0	-	0.0	-	-	-	-	-	-	-	-	-

TABLE 4. (cont.)

*Scomber japonicus* (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
117.0	60.0	-	0.0	-	0.0	-	15.4	-	0.0	-	-	-
118.0	39.0	-	0.0	-	0.0	-	-	0.0	-	2.7	-	-
119.0	33.0	-	0.0	-	0.0	-	-	-	23.9	0.0	-	-
120.0	23.0	-	0.0	-	0.0	-	-	0.0	-	16.9	-	-
120.0	24.0	-	0.0	-	0.0	-	-	0.0	-	9.5	-	-
120.0	25.0	-	0.0	-	0.0	-	-	10.0	-	0.0	-	-
120.0	30.0	-	0.0	-	0.0	-	-	0.0	-	9.0	-	-
120.0	35.0	-	0.0	-	0.0	-	-	2.6	-	0.0	-	-
120.0	40.0	-	0.0	-	0.0	-	-	0.0	-	21.8	-	-
120.0	45.0	-	0.0	-	0.0	-	-	-	14.3	0.0	-	-
123.0	36.0	-	0.0	-	0.0	-	-	0.0	-	9.5	-	-
123.0	37.0	-	0.0	-	0.0	-	-	0.0	-	3.0	-	-
123.0	45.0	-	0.0	-	0.0	-	-	13.3	-	0.0	-	-
123.5	35.6	-	0.0	-	0.0	-	-	0.0	-	2.9	-	-
127.0	45.0	-	0.0	-	0.0	-	-	6.5	-	0.0	-	-
130.0	26.0	-	0.0	-	0.0	-	-	-	15.8	0.0	-	-
130.0	28.0	-	0.0	-	0.0	-	-	-	0.0	7.1	-	-
130.0	30.0	-	0.0	-	0.0	-	-	-	0.0	5.2	-	-
133.0	21.0	-	0.0	-	0.0	-	-	-	-	18.3	-	-
133.0	23.0	-	0.0	-	0.0	-	-	-	0.0	2.5	-	-
137.0	21.0	-	0.0	-	0.0	-	-	-	-	1.4	0.0	-
137.0	22.0	-	0.0	-	0.0	-	-	-	-	6.4	0.0	-

*Scomberomorus* spp.

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
137.0	20.0	-	0.0	-	0.0	-	-	-	36.1	0.0	-	-
137.0	21.0	-	0.0	-	0.0	-	-	-	8.2	0.0	-	-
137.0	22.0	-	0.0	-	0.0	-	-	-	113.4	0.0	-	-
137.0	23.0	-	0.0	-	0.0	-	-	-	20.9	0.0	-	-
140.0	29.0	-	0.0	-	-	-	-	-	2.0	-	-	-

## Trichiuridae

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
103.0	45.0	0.0	-	-	3.0	-	-	0.0	-	0.0	-	-
113.0	45.0	-	0.0	-	0.0	-	-	0.0	-	2.5	-	-
113.0	60.0	-	2.9	-	0.0	-	-	0.0	-	0.0	-	-
115.0	35.0	-	0.0	-	-	-	-	-	-	2.6	-	-
117.0	40.0	-	0.0	-	0.0	-	-	0.0	-	3.1	-	-
117.0	45.0	-	3.0	-	0.0	-	-	0.0	-	2.9	-	-
118.0	39.0	-	3.1	-	0.0	-	-	0.0	-	2.7	-	-
120.0	45.0	-	0.0	-	0.0	-	-	-	-	22.9	0.0	-
120.0	50.0	-	0.0	-	0.0	-	-	-	-	3.0	0.0	-

TABLE 4. (cont.)

## Trichuridae (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
120.0	55.0	-	0.0	-	0.0	-	-	0.0	5.9	-	-	-
123.0	42.0	-	0.0	-	0.0	-	-	6.3	0.0	3.0	-	-
123.0	45.0	-	0.0	-	0.0	-	-	0.0	-	3.0	-	-
123.0	55.0	-	0.0	-	0.0	-	-	2.8	-	0.0	-	-
123.0	60.0	-	0.0	-	0.0	-	-	0.0	-	3.1	-	-
123.0	65.0	-	0.0	-	0.0	-	-	2.8	-	-	-	-
127.0	34.0	-	0.0	-	0.0	-	-	2.4	-	0.0	-	-
127.0	40.0	-	1.4	-	0.0	-	-	0.0	0.0	0.0	-	-
127.0	50.0	-	0.0	-	0.0	-	-	0.0	-	3.0	-	-
130.0	35.0	-	0.0	-	0.0	-	-	-	2.5	3.0	-	-
133.0	40.0	-	3.0	-	0.0	-	-	0.0	-	0.0	-	-
133.0	45.0	-	8.8	-	0.0	-	-	0.0	-	0.0	-	-
137.0	20.0	-	0.0	-	0.0	-	-	-	0.0	1.9	-	-
137.0	30.0	-	3.1	-	-	-	-	-	0.0	-	-	-

*Sphyraena argentea*

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
87.0	32.5	0.0	-	0.0	-	0.0	-	-	-	3.3	-	-
87.0	35.0	0.0	-	0.0	-	0.0	-	-	-	3.0	-	-
90.0	32.0	0.0	-	0.0	-	-	-	2.5	-	0.0	-	-
93.0	26.5	0.0	-	-	-	-	26.5	-	0.0	-	-	-
93.0	27.0	0.0	-	0.0	-	-	45.9	-	0.0	-	-	-
93.0	28.0	0.0	-	0.0	-	-	9.6	-	0.0	-	-	-
93.0	30.0	0.0	-	0.0	-	-	27.6	-	-	5.9	-	-
93.0	35.0	0.0	-	0.0	-	-	3.0	-	-	0.0	-	-
115.0	35.0	-	0.0	-	-	-	-	-	2.6	0.0	-	-
117.0	24.8	-	3.0	-	0.0	-	-	0.0	-	0.0	-	-
120.0	23.0	-	0.0	-	0.0	-	-	0.0	-	1.9	-	-
120.0	40.0	-	0.0	-	0.0	-	-	2.6	-	0.0	-	-
120.0	45.0	-	0.0	-	0.0	-	-	-	2.9	0.0	-	-
127.0	40.0	-	0.0	-	0.0	-	-	8.5	0.0	0.0	-	-
127.0	45.0	-	0.0	-	0.0	-	-	3.2	-	0.0	-	-
127.0	45.0	-	0.0	-	0.0	-	-	-	28.4	0.0	-	-
130.0	25.5	-	0.0	-	0.0	-	-	-	-	9.1	-	-
130.0	26.0	-	0.0	-	0.0	-	-	-	-	15.3	-	-
130.0	35.0	-	0.0	-	0.0	-	-	-	-	8.9	-	-
133.0	30.0	-	0.0	-	0.0	-	-	-	-	0.4	-	-
137.0	20.0	-	0.0	-	0.0	-	-	-	-	28.8	0.0	-
137.0	21.0	-	0.0	-	0.0	-	-	-	-	55.6	0.0	-
137.0	22.0	-	0.0	-	0.0	-	-	-	-	39.9	0.0	-
137.0	23.0	-	0.0	-	0.0	-	-	-	-	5.9	-	-
140.0	29.0	-	0.0	-	0.0	-	-	-	-	-	-	-

TABLE 4. (cont.)

*Icichthys lockingtoni*

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
60.0	60.0	0.0	-	-	-	-	-	-	-	-	-	-
60.0	70.0	3.2	-	-	-	-	-	-	-	-	-	-
60.0	80.0	0.0	-	-	-	-	-	-	-	-	-	-
60.0	120.0	0.0	-	-	-	-	-	-	-	-	-	-
63.0	60.0	0.0	-	-	-	-	-	-	-	-	-	-
67.0	55.0	0.0	-	-	-	-	-	-	-	-	-	-
67.0	65.0	-	-	-	-	-	-	-	-	-	-	-
67.0	70.0	-	-	-	-	-	-	-	-	-	-	-
70.0	52.0	6.3	-	-	-	-	-	-	-	-	-	-
70.0	53.0	3.0	-	-	-	-	-	-	-	-	-	-
70.0	55.0	3.0	-	-	-	-	-	-	-	-	-	-
70.0	60.0	8.7	-	-	-	-	-	-	-	-	-	-
70.0	70.0	0.0	-	-	-	-	-	-	-	-	-	-
70.0	90.0	0.0	-	-	-	-	-	-	-	-	-	-
73.0	49.4	0.0	-	-	-	-	-	-	-	-	-	-
73.0	53.0	2.7	-	-	-	-	-	-	-	-	-	-
73.0	60.0	0.0	-	-	-	-	-	-	-	-	-	-
73.0	70.0	-	-	-	-	-	-	-	-	-	-	-
73.0	80.0	-	-	-	-	-	-	-	-	-	-	-
73.0	80.0	2.6	-	-	-	-	-	-	-	-	-	-
77.0	55.0	2.9	-	-	-	-	-	-	-	-	-	-
77.0	57.0	3.0	-	-	-	-	-	-	-	-	-	-
77.0	65.0	-	-	-	-	-	-	-	-	-	-	-
80.0	65.0	6.2	-	-	-	-	-	-	-	-	-	-
80.0	70.0	0.0	-	-	-	-	-	-	-	-	-	-
80.0	80.0	6.0	-	-	-	-	-	-	-	-	-	-
83.0	65.0	0.0	-	-	-	-	-	-	-	-	-	-
83.0	70.0	0.0	-	-	-	-	-	-	-	-	-	-
83.0	80.0	0.0	-	-	-	-	-	-	-	-	-	-
83.0	90.0	0.0	-	-	-	-	-	-	-	-	-	-
87.0	55.0	0.0	-	-	-	-	-	-	-	-	-	-
87.0	60.0	0.0	-	-	-	-	-	-	-	-	-	-
87.0	65.0	0.0	-	-	-	-	-	-	-	-	-	-
87.0	70.0	0.0	-	-	-	-	-	-	-	-	-	-
87.0	75.0	-	-	-	-	-	-	-	-	-	-	-
87.0	80.0	3.0	-	-	-	-	-	-	-	-	-	-
87.0	90.0	0.0	-	-	-	-	-	-	-	-	-	-
90.0	60.0	0.0	-	-	-	-	-	-	-	-	-	-
90.0	65.0	0.0	-	-	-	-	-	-	-	-	-	-
93.0	50.0	0.0	-	-	-	-	-	-	-	-	-	-
93.0	70.0	0.0	-	-	-	-	-	-	-	-	-	-
97.0	40.0	0.0	-	-	-	-	-	-	-	-	-	-
100.0	35.0	0.0	-	-	-	-	-	-	-	-	-	-
100.0	60.0	3.0	-	-	-	-	-	-	-	-	-	-
103.0	65.0	2.8	-	-	-	-	-	-	-	-	-	-
103.0	70.0	3.0	-	-	-	-	-	-	-	-	-	-

TABLE 4. (cont.)

Nomeidae												
STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
113.0	40.0	-	0.0	-	0.0	-	0.0	-	12.4	-	-	-
<i>Peprius simillimus</i>												
STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
77.0	49.0	-	-	-	0.7	-	-	0.0	-	-	-	-
80.0	50.2	0.0	-	-	1.5	-	-	0.0	0.0	-	-	-
80.0	51.0	0.0	-	-	3.5	-	-	0.0	0.0	-	-	-
80.0	52.0	0.0	-	-	6.7	-	-	0.0	0.0	-	-	-
83.0	40.0	0.0	-	-	0.8	-	-	0.0	0.0	-	-	-
83.0	43.0	0.0	-	-	0.0	-	-	15.8	-	-	-	-
83.0	51.0	0.0	-	-	2.6	-	-	0.0	-	-	-	-
87.0	32.5	0.0	-	-	0.0	-	-	1.7	-	-	-	-
87.0	33.0	0.0	-	-	0.0	-	-	16.1	-	-	-	-
87.0	34.0	0.0	-	-	0.0	-	-	10.6	-	-	-	-
90.0	37.0	0.0	-	-	0.0	-	-	2.1	-	-	-	-
93.0	26.5	0.0	-	-	-	-	-	2.5	-	-	-	-
93.0	27.0	0.0	-	-	0.0	-	-	14.3	-	-	-	-
93.0	30.0	0.0	-	-	-	-	-	3.1	-	-	-	-
103.0	30.0	0.0	-	-	-	-	-	2.6	-	-	-	-
117.0	25.0	-	0.0	-	0.0	-	-	0.0	-	-	-	-
117.0	35.0	-	0.0	-	0.0	-	-	3.3	-	-	-	-
117.0	60.0	-	0.0	-	0.0	-	-	3.8	-	-	-	-
119.0	33.0	-	0.0	-	2.9	-	-	-	32.6	-	-	-
120.0	23.0	-	0.0	-	2.5	-	-	-	2.9	-	-	-
120.0	24.0	-	0.0	-	55.7	-	-	-	0.0	-	-	-
120.0	25.0	-	0.0	-	3.7	-	-	-	6.0	-	-	-
120.0	35.0	-	0.0	-	6.1	-	-	-	18.2	-	-	-
120.0	40.0	-	0.0	-	0.0	-	-	-	5.2	-	-	-
123.0	45.0	-	0.0	-	2.9	-	-	-	0.0	-	-	-
127.0	45.0	-	0.0	-	0.0	-	-	-	3.2	-	-	-
133.0	21.0	-	4.0	-	0.0	-	-	-	2.0	-	-	-
133.0	23.0	-	4.0	-	18.6	-	-	-	0.0	-	-	-
133.0	25.0	-	40.5	-	40.5	-	-	-	0.0	-	-	-
133.0	30.0	-	2.8	-	2.8	-	-	-	0.0	-	-	-
133.0	35.0	-	2.9	-	2.9	-	-	-	0.0	-	-	-
133.0	40.0	-	3.0	-	3.0	-	-	-	0.0	-	-	-
133.0	45.0	-	2.9	-	2.9	-	-	-	0.0	-	-	-
137.0	20.0	-	0.3	-	0.3	-	-	-	0.0	-	-	-
137.0	21.0	-	12.0	-	12.0	-	-	-	0.0	-	-	-
137.0	22.0	-	40.3	-	40.3	-	-	-	0.0	-	-	-
137.0	23.0	-	3.7	-	3.7	-	-	-	0.0	-	-	-
140.0	28.8	-	2.7	-	2.7	-	-	-	0.0	-	-	-
140.0	30.0	-	13.6	-	13.6	-	-	-	0.0	-	-	-

TABLE 4. (cont.)

*Tetragonurus cuvieri*

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
60.0	140.0	0.0	-	-	-	-	-	2.6	-	-	-	-
60.0	180.0	0.0	-	-	-	-	-	2.7	-	-	-	-
70.0	120.0	3.0	-	-	-	-	-	0.0	-	-	-	-
80.0	100.0	3.0	-	-	-	-	-	0.0	-	-	-	-
80.0	140.0	2.7	-	-	-	-	-	-	-	-	-	-
87.0	65.0	0.0	-	0.0	0.0	0.0	-	-	-	-	2.9	-
87.0	90.0	0.0	-	0.0	0.0	3.1	-	2.5	-	-	0.0	-
90.0	32.0	0.0	-	0.0	0.0	-	-	2.5	-	-	20.8	-
90.0	70.0	0.0	-	0.0	0.0	-	-	0.0	-	-	24.6	-
90.0	90.0	0.0	-	0.0	0.0	-	-	0.0	-	-	-	-
90.0	100.0	5.5	-	0.0	0.0	-	-	0.0	-	-	-	-
90.0	120.0	0.0	-	-	-	-	-	5.4	-	-	-	-
90.0	140.0	0.0	-	-	-	-	-	2.7	-	-	-	-
93.0	75.0	-	-	-	-	-	-	-	-	-	2.9	-
93.0	80.0	0.0	-	0.0	0.0	-	-	0.0	-	-	29.4	-
93.0	85.0	-	-	-	-	-	-	-	-	-	29.9	-
93.0	90.0	0.0	-	0.0	0.0	-	-	0.0	-	-	14.3	-
97.0	80.0	0.0	-	-	-	-	-	0.0	-	-	11.6	-
97.0	100.0	-	-	-	-	-	-	-	-	-	-	-
100.0	100.0	-	3.0	-	-	-	-	0.0	-	-	-	-
100.0	120.0	-	0.0	-	-	-	-	3.0	-	-	-	-
103.0	60.0	0.0	-	-	-	-	-	0.0	-	-	3.1	-
103.0	90.0	-	6.1	-	-	-	-	0.0	-	-	-	-
107.0	65.0	-	0.0	-	-	-	-	0.0	-	-	3.3	-
107.0	70.0	-	0.0	-	-	-	-	0.0	-	-	3.0	-
110.0	45.0	-	0.0	-	-	-	-	0.0	-	-	10.1	-
113.0	40.0	-	0.0	-	-	-	-	0.0	-	-	3.1	-
113.0	45.0	-	0.0	-	-	-	-	0.0	-	-	2.5	-
117.0	40.0	-	3.0	-	-	-	-	0.0	-	-	0.0	-
127.0	40.0	-	0.0	-	-	-	-	0.0	-	-	2.7	-
140.0	45.0	-	0.0	-	-	-	-	0.0	-	-	2.8	-

## Chiassmodontidae

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
90.0	80.0	0.0	-	-	-	-	-	0.0	-	-	0.0	-
93.0	80.0	0.0	-	-	-	-	-	0.0	-	-	0.0	-
97.0	45.0	0.0	-	-	3.5	-	-	0.0	-	-	0.0	-
97.0	50.0	0.0	-	-	0.0	-	-	3.3	-	-	0.0	-
97.0	85.0	-	-	-	-	-	-	-	-	-	-	-
100.0	85.0	-	-	-	-	-	-	7.2	-	-	-	-
100.0	90.0	0.0	-	-	-	-	-	3.5	-	-	0.0	-
103.0	50.0	0.0	-	-	-	-	-	3.3	-	-	0.0	-
103.0	75.0	-	-	-	-	-	-	6.4	-	-	0.0	-
107.0	80.0	-	0.0	-	-	-	-	0.0	-	-	3.2	-

TABLE 4. (cont.)

## Chiasmodontidae (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
110.0	75.0	-	-	3.2	-	-	-	-	-	-	-	-
110.0	80.0	-	2.6	0.0	-	-	-	-	-	-	-	-
113.0	65.0	-	0.0	3.3	-	0.0	-	-	-	-	-	-
113.0	70.0	-	0.0	3.1	-	0.0	-	-	-	-	-	-
115.0	35.0	-	0.0	-	-	-	-	-	2.6	0.0	-	-
117.0	65.0	-	0.0	2.8	-	0.0	-	-	-	-	-	-
117.0	85.0	-	-	2.9	0.0	-	-	-	-	-	-	-
117.0	90.0	-	2.9	0.0	-	-	-	-	-	-	-	-
120.0	65.0	-	0.0	3.1	-	-	-	-	-	-	-	-
120.0	70.0	-	0.0	6.6	-	-	-	-	-	-	-	-
120.0	75.0	-	-	3.1	-	-	-	-	-	-	-	-
123.0	42.0	-	0.0	2.9	-	-	-	-	-	-	-	-
123.0	50.0	-	0.0	0.0	-	-	-	-	-	-	-	-
123.0	70.0	-	0.0	6.2	-	-	-	-	-	-	-	-
127.0	45.0	-	0.0	6.2	-	-	-	-	-	-	-	-
127.0	50.0	-	0.0	6.1	-	-	-	-	-	-	-	-
127.0	70.0	-	0.0	0.0	-	-	-	-	-	-	-	-
127.0	80.0	-	2.9	-	-	-	-	-	-	-	-	-
130.0	35.0	-	0.0	3.1	-	-	-	-	-	-	-	-
130.0	45.0	-	0.0	0.0	-	-	-	-	-	-	-	-
130.0	50.0	-	0.0	0.0	-	-	-	-	-	-	-	-
130.0	55.0	-	2.9	0.0	-	-	-	-	-	-	-	-
130.0	60.0	-	0.0	0.0	-	-	-	-	-	-	-	-
130.0	90.0	-	0.0	-	-	-	-	-	-	-	-	-
130.0	100.0	-	2.8	-	-	-	-	-	-	-	-	-
133.0	55.0	-	0.0	5.9	-	-	-	-	-	-	-	-
133.0	65.0	-	0.0	3.1	-	-	-	-	-	-	-	-

## Pleuronectiformes

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
77.0	47.6	0.6	-	0.0	-	-	-	-	0.0	-	-	-
80.0	52.0	0.0	-	0.0	-	-	-	-	0.0	-	-	-
118.0	39.0	-	0.0	0.0	-	-	-	-	2.8	-	0.0	-
120.0	24.0	-	0.0	0.0	-	-	-	-	3.5	-	0.0	-

## Citharichthys spp.

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
60.0	49.5	1.7	-	-	-	-	-	-	-	-	-	-
60.0	51.0	11.3	-	-	-	-	-	-	-	-	0.0	-
60.0	52.0	6.8	-	-	-	-	-	-	-	-	0.0	-
60.0	55.0	9.3	-	-	-	-	-	-	-	-	0.0	-
60.0	70.0	6.5	-	-	-	-	-	-	-	-	-	-

TABLE 4. (cont.)

*Citharichthys* spp. (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
60.0	80.0	2.5	-	-	0.0	-	-	-	-	-	-	-
63.0	50.0	1.6	-	-	0.0	-	-	-	-	-	-	-
63.0	52.0	2.9	-	-	0.0	-	-	-	-	-	-	-
63.0	55.0	15.9	-	-	0.0	-	-	-	-	-	-	-
67.0	50.0	2.3	-	-	2.7	-	-	-	-	-	-	-
67.0	55.0	10.0	-	-	0.0	-	-	-	-	-	-	-
67.0	60.0	6.0	-	-	0.0	-	-	-	-	-	-	-
67.0	65.0	-	-	-	2.9	-	-	-	-	-	-	-
70.0	50.0	29.3	-	-	0.0	-	-	-	-	-	-	-
70.0	51.0	6.1	-	-	0.0	-	-	-	-	-	-	-
70.0	52.0	6.3	-	-	0.0	-	-	-	-	-	-	-
70.0	53.0	3.0	-	-	5.4	-	-	-	-	-	-	-
70.0	55.0	8.7	-	-	-	-	-	-	-	-	-	-
70.0	60.0	0.0	-	-	2.9	-	-	-	-	-	-	-
70.0	70.0	2.7	-	-	0.0	-	-	-	-	-	-	-
73.0	49.4	0.0	-	-	0.7	-	-	-	-	-	-	-
73.0	50.0	7.0	-	-	0.0	-	-	-	-	-	-	-
73.0	51.0	3.1	-	-	2.8	-	-	-	-	-	-	-
73.0	53.0	2.7	-	-	3.0	-	-	-	-	-	-	-
73.0	60.0	2.8	-	-	0.0	-	-	-	-	-	-	-
77.0	47.6	9.8	-	-	0.0	-	-	-	-	-	-	-
77.0	48.0	0.3	-	-	0.0	-	-	-	-	-	-	-
77.0	51.0	0.0	-	-	0.0	-	-	-	-	-	-	-
77.0	55.0	5.8	-	-	0.0	-	-	-	-	-	-	-
77.0	60.0	-	-	-	3.1	-	-	-	-	-	-	-
80.0	50.2	7.6	-	-	0.8	-	-	-	-	-	-	-
80.0	51.0	25.3	-	-	0.0	-	-	-	-	-	-	-
80.0	52.0	16.5	-	-	3.3	-	-	-	-	-	-	-
80.0	55.0	0.0	-	-	3.2	-	-	-	-	-	-	-
80.0	60.0	5.1	-	-	0.0	-	-	-	-	-	-	-
80.0	65.0	3.1	-	-	0.0	-	-	-	-	-	-	-
80.0	70.0	0.0	-	-	3.4	-	-	-	-	-	-	-
80.0	80.0	80.0	-	-	3.2	-	-	-	-	-	-	-
82.0	47.0	0.0	-	-	0.0	-	-	-	-	-	-	-
83.0	39.0	0.0	-	-	-	-	-	-	-	-	-	-
83.0	40.0	1.9	-	-	-	-	-	-	-	-	-	-
83.0	43.0	0.0	-	-	5.6	-	-	-	-	-	-	-
83.0	51.0	0.0	-	-	2.6	-	-	-	-	-	-	-
83.0	55.0	5.8	-	-	0.0	-	-	-	-	-	-	-
83.0	60.0	3.0	-	-	6.0	-	-	-	-	-	-	-
83.0	65.0	8.6	-	-	0.0	-	-	-	-	-	-	-
83.0	70.0	16.0	-	-	0.0	-	-	-	-	-	-	-
83.0	75.0	-	-	-	-	-	-	-	-	-	-	-
83.0	80.0	14.1	-	-	-	-	-	-	-	-	-	-
87.0	32.5	0.0	-	-	5.7	-	-	-	-	-	-	-
87.0	33.0	0.0	-	-	3.5	-	-	-	-	-	-	-
87.0	81.1	8.1	-	-	8.1	-	-	-	-	-	-	-

TABLE 4. (cont.)

*Citharichthys* spp. (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
87.0	34.0	26.7	-	-	0.0	-	-	-	-	7.9	-	-
87.0	35.0	17.6	-	-	0.0	-	9.2	-	-	0.0	-	-
87.0	40.0	5.7	-	-	0.0	-	0.0	-	-	3.0	-	-
87.0	45.0	9.2	-	-	0.0	-	3.3	-	-	0.0	-	-
87.0	50.0	7.9	-	-	0.0	-	0.0	-	-	0.0	-	-
87.0	55.0	11.9	-	-	0.0	-	3.3	-	-	0.0	-	-
87.0	65.0	3.1	-	-	0.0	-	0.0	-	-	0.0	-	-
87.0	70.0	3.1	-	-	0.0	-	0.0	-	-	6.1	-	-
87.0	75.0	-	-	-	-	-	-	-	-	3.2	-	-
87.0	80.0	3.0	-	-	3.0	-	0.0	-	-	0.0	-	-
87.0	80.0	0.6	-	-	0.0	-	0.0	-	-	0.0	-	-
90.0	27.5	16.7	-	-	0.0	-	0.0	-	-	0.0	-	-
90.0	28.0	21.6	-	-	0.0	-	0.0	-	-	0.0	-	-
90.0	32.0	12.0	-	-	0.0	-	0.0	-	-	0.0	-	-
90.0	37.0	6.5	-	-	0.0	-	0.0	-	-	0.0	-	-
90.0	53.0	6.5	-	-	0.0	-	0.0	-	-	0.0	-	-
90.0	60.0	0.0	-	-	0.0	-	0.0	-	-	0.0	-	-
90.0	90.0	3.0	-	-	0.0	-	0.0	-	-	3.0	-	-
90.0	100.0	0.0	-	-	3.1	-	-	-	-	0.0	-	-
93.0	26.5	0.0	-	-	-	-	2.5	-	-	0.0	-	-
93.0	27.0	2.8	-	-	0.0	-	2.9	-	-	15.3	-	-
93.0	28.0	3.1	-	-	0.0	-	0.0	-	-	26.7	-	-
93.0	30.0	6.1	-	-	0.0	-	0.0	-	-	0.0	-	-
93.0	40.0	0.0	-	-	0.0	-	0.0	-	-	0.0	-	-
93.0	55.0	3.1	-	-	0.0	-	0.0	-	-	0.0	-	-
93.0	80.0	3.0	-	-	0.0	-	0.0	-	-	0.0	-	-
96.8	28.7	14.2	-	-	0.0	-	0.0	-	-	0.0	-	-
97.0	29.0	43.0	-	-	4.8	-	0.0	-	-	0.0	-	-
97.0	30.0	4.3	-	-	7.6	-	5.6	-	-	0.0	-	-
97.0	32.0	6.2	-	-	31.0	-	6.7	-	-	0.0	-	-
97.0	35.0	0.0	-	-	6.3	-	0.0	-	-	2.9	-	-
97.0	40.0	0.0	-	-	0.0	-	3.3	-	-	21.2	-	-
97.0	45.0	0.0	-	-	0.0	-	0.0	-	-	15.2	-	-
97.0	70.0	5.9	-	-	-	-	-	-	-	0.0	-	-
100.0	29.0	3.8	-	-	-	-	-	-	-	5.5	-	-
100.0	30.0	14.1	-	-	-	-	-	-	-	30.4	-	-
100.0	35.0	0.0	-	-	-	-	-	-	-	3.1	-	-
100.0	40.0	0.0	-	-	0.0	-	0.0	-	-	0.0	-	-
100.0	70.0	3.1	-	-	0.0	-	0.0	-	-	0.0	-	-
103.0	28.5	1.3	-	-	-	-	-	-	-	3.1	-	-
103.0	30.0	2.5	-	-	6.1	-	0.0	-	-	0.0	-	-
103.0	35.0	0.0	-	-	3.1	-	0.0	-	-	0.0	-	-
103.0	40.0	0.0	-	-	0.0	-	0.0	-	-	0.0	-	-
103.0	45.0	0.0	-	-	-	-	-	-	-	6.8	-	-
107.0	30.0	-	-	-	-	-	-	-	-	0.0	-	-
107.0	31.0	-	-	-	-	-	-	-	-	1.5	-	-
107.0	32.0	-	-	-	-	-	-	-	-	2.8	-	-

TABLE 4. (cont.)

*Citharichthys spp.* (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
107.0	35.0	-	0.0	-	-	3.5	-	-	3.0	-	-	-
107.0	40.0	-	0.0	0.0	0.0	0.0	-	-	5.9	-	-	-
107.0	45.0	-	0.0	0.0	0.0	0.0	-	-	3.3	-	-	-
107.0	55.0	-	0.0	0.0	0.0	0.0	-	-	3.1	-	-	-
110.0	32.0	-	1.3	0.0	-	-	-	0.0	29.2	-	-	-
110.0	35.0	-	0.0	0.0	-	-	-	0.0	17.9	-	-	-
110.0	40.0	-	0.0	0.0	-	-	-	0.0	18.8	-	-	-
110.0	50.0	-	5.6	0.0	-	-	-	0.0	3.1	-	-	-
110.0	55.0	-	0.0	0.0	-	-	-	5.7	0.0	-	-	-
110.0	60.0	-	0.0	0.0	-	-	-	8.3	0.0	-	-	-
110.0	70.0	-	0.0	0.0	-	-	-	18.1	-	-	-	-
110.0	80.0	-	0.0	0.0	-	-	-	16.0	-	-	-	-
113.0	28.5	-	0.9	0.0	0.0	0.0	-	-	20.2	-	-	-
113.0	29.0	-	10.8	0.0	0.0	0.0	-	-	16.0	-	-	-
113.0	30.0	-	68.5	5.6	0.0	0.0	-	90.5	16.1	-	-	-
113.0	35.0	-	6.0	3.5	0.0	0.0	-	-	42.1	-	-	-
113.0	40.0	-	0.0	0.0	0.0	0.0	-	-	3.1	-	-	-
113.0	45.0	-	0.0	0.0	0.0	0.0	-	-	14.8	-	-	-
113.0	50.0	-	0.0	0.0	0.0	0.0	-	-	3.0	-	-	-
113.0	55.0	-	0.0	0.0	0.0	0.0	-	-	0.0	-	-	-
113.0	60.0	-	11.8	0.0	0.0	0.0	-	-	0.0	-	-	-
113.0	65.0	-	6.2	0.0	0.0	0.0	-	-	-	-	-	-
115.0	35.0	-	19.9	-	-	-	-	-	289.7	81.4	-	-
117.0	24.8	-	15.0	0.0	0.0	0.0	-	28.6	20.7	-	-	-
117.0	25.0	-	26.1	2.8	-	-	-	37.8	136.2	-	-	-
117.0	26.0	-	67.2	27.1	-	-	-	138.8	181.5	-	-	-
117.0	30.0	-	453.7	13.0	0.0	0.0	-	32.3	49.4	-	-	-
117.0	35.0	-	6.3	13.0	-	-	-	-	90.5	-	-	-
117.0	40.0	-	15.3	0.0	-	-	-	0.0	-	6.2	-	-
117.0	45.0	-	38.5	0.0	-	-	-	3.4	11.6	-	-	-
117.0	50.0	-	3.2	49.5	-	-	-	19.9	0.0	-	-	-
117.0	55.0	-	0.0	0.0	-	-	-	69.3	0.0	-	-	-
117.0	60.0	-	0.0	0.0	-	-	-	77.0	-	0.0	-	-
118.0	39.0	-	6.2	80.3	-	-	-	56.0	-	13.6	-	-
119.0	33.0	-	191.5	29.0	-	-	-	516.5	-	71.5	-	-
120.0	22.5	-	0.0	18.1	-	-	-	21.6	-	33.0	-	-
120.0	23.0	-	1.3	5.6	-	-	-	25.0	-	7.5	-	-
120.0	24.0	-	58.3	12.5	-	-	-	617.1	-	54.7	-	-
120.0	25.0	-	84.2	8.0	-	-	-	1986.0	-	81.5	-	-
120.0	30.0	-	107.3	812.3	-	-	-	1505.9	-	90.3	-	-
120.0	35.0	-	155.6	133.9	-	-	-	764.4	-	27.4	-	-
120.0	40.0	-	18.6	39.9	-	-	-	1029.4	-	13.7	-	-
120.0	45.0	-	5.8	12.4	-	-	-	68.6	0.0	-	-	-
120.0	50.0	-	24.6	0.0	-	-	-	0.0	0.0	-	-	-
120.0	55.0	-	2.8	0.0	-	-	-	2.6	-	8.9	-	-
120.0	65.0	-	2.3	0.0	-	-	-	2.7	-	-	-	-

TABLE 4. (cont.)

*Citharichthys* spp. (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
120.0	70.0	-	2.9	-	0.0	-	-	39.8	-	0.0	-	-
123.0	36.0	-	18.7	-	4.8	-	-	36.0	-	0.0	-	-
123.0	37.0	-	17.2	-	2.5	-	-	3.2	56.7	3.0	-	-
123.0	42.0	-	35.9	-	8.6	-	-	19.9	-	0.0	-	-
123.0	45.0	-	0.0	-	0.0	-	-	0.0	-	0.0	-	-
123.0	50.0	-	2.9	-	0.0	-	-	0.0	-	0.0	-	-
123.0	55.0	-	0.0	-	0.0	-	-	5.7	-	3.1	-	-
123.5	35.6	-	8.9	-	4.9	-	-	6.3	-	0.0	-	-
127.0	32.6	-	34.3	-	19.9	-	-	11.0	-	0.0	-	-
127.0	33.0	-	373.3	-	13.5	-	-	0.0	-	0.0	-	-
127.0	34.0	-	147.3	-	22.0	-	-	65.3	-	0.0	-	-
127.0	40.0	-	43.9	-	219.8	-	-	2.8	2.7	0.0	-	-
127.0	45.0	-	0.0	-	0.0	-	-	148.6	-	6.2	-	-
127.0	50.0	-	0.0	-	0.0	-	-	5.8	-	0.0	-	-
127.0	55.0	-	0.0	-	0.0	-	-	2.7	-	3.2	-	-
130.0	25.5	-	1.3	-	0.0	-	-	-	591.3	7.7	-	-
130.0	26.0	-	1.9	-	0.0	-	-	-	16.6	2.3	-	-
130.0	28.0	-	21.4	-	16.3	-	-	-	12.4	2.4	-	-
130.0	30.0	-	0.0	-	0.0	-	-	-	0.0	23.2	-	-
130.0	35.0	-	20.0	-	6.2	-	-	-	22.9	6.1	-	-
130.0	40.0	-	0.0	-	78.8	-	-	-	0.0	0.0	-	-
130.0	45.0	-	5.6	-	0.0	-	-	-	0.0	0.0	-	-
130.0	50.0	-	2.9	-	0.0	-	-	-	0.0	0.0	-	-
133.0	21.0	-	6.0	-	0.0	-	-	-	0.0	320.6	-	-
133.0	23.0	-	10.6	-	0.0	-	-	-	0.0	17.5	-	-
133.0	25.0	-	24.3	-	60.8	-	-	2.7	-	2.9	-	-
133.0	30.0	-	16.6	-	37.2	-	-	29.6	-	5.7	-	-
133.0	35.0	-	20.2	-	9.3	-	-	29.6	-	3.0	-	-
133.0	40.0	-	17.9	-	0.0	-	-	23.3	-	0.0	-	-
133.0	45.0	-	120.1	-	0.0	-	-	0.0	-	0.0	-	-
133.0	50.0	-	19.8	-	0.0	-	-	0.0	-	3.0	-	-
137.0	20.0	-	0.0	-	6.6	-	-	0.0	-	9.4	-	-
137.0	21.0	-	2.5	-	2.2	-	-	58.9	49.7	-	-	-
137.0	22.0	-	2.6	-	4.5	-	-	19.3	40.8	-	-	-
137.0	23.0	-	1.9	-	21.1	-	-	1.9	44.2	-	-	-
137.0	30.0	-	6.2	-	9.3	-	-	0.0	15.3	-	-	-
137.0	35.0	-	3.1	-	0.0	-	-	0.0	0.0	0.0	-	-
140.0	28.8	-	0.0	-	-	-	-	-	1.4	-	-	-
140.0	29.0	-	2.3	-	-	-	-	-	7.8	-	-	-

*Citharichthys stigmaeus*

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
60.0	60.0	2.7	-	-	-	-	-	-	0.0	-	-	-
60.0	80.0	0.0	-	-	-	-	-	-	3.1	-	-	-

TABLE 4. (cont.)

*Citharichthys stigmaeus* (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
67.0	49.0	0.0	-	-	0.0	-	-	3.1	-	-	-	-
67.0	55.0	0.0	-	-	2.7	-	-	0.0	-	-	-	-
67.0	70.0	-	-	-	2.8	-	-	-	-	-	-	-
67.0	80.0	-	3.2	-	2.7	-	-	-	-	-	-	-
70.0	52.0	-	-	-	0.0	-	-	-	-	-	-	-
70.0	65.0	-	-	-	2.7	-	-	-	-	-	-	-
77.0	65.0	-	-	-	5.1	-	-	-	-	-	-	-
80.0	55.0	0.0	-	-	3.2	-	-	0.0	-	-	-	-
80.0	60.0	0.0	-	-	3.4	-	-	0.0	-	-	-	-
80.0	65.0	3.1	-	-	0.0	-	-	0.0	-	-	-	-
80.0	70.0	2.8	-	-	3.4	-	-	0.0	-	-	-	-
80.0	75.0	-	-	-	-	-	-	10.8	-	8.2	-	-
80.0	80.0	0.0	-	-	3.2	-	-	0.0	-	0.0	-	-
80.0	120.0	3.0	-	-	-	-	-	0.0	-	-	-	-
83.0	55.0	2.9	-	-	0.0	-	-	0.0	-	0.0	-	-
83.0	60.0	0.0	-	-	0.0	-	-	6.9	-	2.7	-	-
83.0	65.0	5.7	-	-	0.0	-	-	0.0	-	0.0	-	-
83.0	70.0	0.0	-	-	0.0	-	-	15.8	-	2.8	-	-
87.0	34.0	-	-	-	0.0	-	-	0.0	-	0.0	-	-
87.0	45.0	0.0	-	-	0.0	-	-	0.0	-	3.2	-	-
87.0	50.0	0.0	-	-	0.0	-	-	0.0	-	2.7	-	-
87.0	70.0	3.1	-	-	0.0	-	-	0.0	-	0.0	-	-
87.0	80.0	0.0	-	-	0.0	-	-	0.0	-	3.0	-	-
90.0	37.0	-	-	-	0.0	-	-	6.3	-	0.0	-	-
90.0	65.0	-	-	-	0.0	-	-	8.0	-	0.0	-	-
90.0	70.0	0.0	-	-	0.0	-	-	5.5	-	0.0	-	-
90.0	90.0	0.0	-	-	0.0	-	-	0.0	-	3.1	-	-
90.0	100.0	0.0	-	-	0.0	-	-	2.7	-	0.0	-	-
93.0	127.0	2.8	-	-	0.0	-	-	0.0	-	0.0	-	-
93.0	35.0	0.0	-	-	5.8	-	-	3.0	-	0.0	-	-
93.0	40.0	0.0	-	-	0.0	-	-	3.2	-	0.0	-	-
93.0	50.0	0.0	-	-	0.0	-	-	0.0	-	2.6	-	-
93.0	80.0	3.0	-	-	0.0	-	-	6.5	-	0.0	-	-
97.0	29.0	0.0	-	-	0.0	-	-	6.5	-	0.0	-	-
97.0	35.0	0.0	-	-	0.0	-	-	0.0	-	3.0	-	-
97.0	40.0	0.0	-	-	0.0	-	-	0.0	-	4.2	-	-
97.0	50.0	3.2	-	-	0.0	-	-	0.0	-	2.8	-	-
100.0	29.0	0.0	-	-	0.0	-	-	0.0	-	0.0	-	-
100.0	35.0	0.0	-	-	0.0	-	-	0.0	-	3.2	-	-
100.0	45.0	0.0	-	-	0.0	-	-	0.0	-	0.0	-	-
100.0	55.0	0.0	-	-	0.0	-	-	0.0	-	0.0	-	-
103.0	30.0	-	-	-	0.0	-	-	0.0	-	0.0	-	-
103.0	35.0	0.0	-	-	0.0	-	-	3.1	-	0.0	-	-
103.0	40.0	0.0	-	-	0.0	-	-	0.0	-	0.0	-	-
107.0	32.0	-	-	-	9.0	-	-	0.0	-	0.0	-	-

TABLE 4. (cont.)

*Citharichthys stigmaeus* (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
107.0	40.0	-	0.0	-	0.0	-	6.7	-	-	0.0	-	-
110.0	35.0	-	0.0	-	0.0	-	-	-	0.0	6.0	-	-
110.0	40.0	-	0.0	-	2.8	-	-	-	0.0	0.0	-	-
110.0	45.0	-	0.0	-	0.0	-	-	-	2.6	0.0	-	-
110.0	70.0	-	0.0	-	0.0	-	-	-	3.0	-	-	-
110.0	80.0	-	0.0	-	0.0	-	-	10.7	-	0.0	-	-
123.0	42.0	-	0.0	-	0.0	-	-	0.0	3.2	0.0	-	-
127.0	45.0	-	0.0	-	0.0	-	-	6.5	-	0.0	-	-

*Hippoglossina stomachata*

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
80.0	65.0	0.0	-	-	0.0	-	-	2.7	0.0	3.0	-	-
82.0	47.0	0.0	-	-	0.0	-	-	6.3	-	0.0	8.6	-
83.0	43.0	0.0	-	-	2.8	-	-	0.0	-	0.0	-	-
83.0	55.0	2.9	-	-	0.0	-	-	2.7	-	0.0	-	-
87.0	33.0	0.0	-	-	0.0	-	-	0.0	-	2.6	-	-
87.0	34.0	0.0	-	-	0.0	-	-	2.3	0.0	0.0	-	-
87.0	50.0	0.0	-	-	0.0	-	-	0.0	-	0.0	-	-
90.0	28.0	0.0	-	-	0.0	-	-	0.0	-	0.0	-	-
93.0	26.5	0.0	-	-	0.0	-	-	0.0	-	0.0	-	-
97.0	29.0	0.0	-	-	0.0	-	-	0.0	-	0.0	-	-
97.0	30.0	0.0	-	-	0.0	-	-	0.0	-	0.0	-	-
117.0	24.8	-	0.0	-	0.0	-	-	1.9	-	2.4	-	-
117.0	25.0	-	0.0	-	0.0	-	-	0.0	-	2.3	-	-
117.0	26.0	-	0.0	-	0.0	-	-	0.0	-	1.6	-	-
117.0	30.0	-	0.0	-	0.0	-	-	0.0	-	0.0	-	-
117.0	35.0	-	0.0	-	0.0	-	-	0.0	-	0.0	-	-
118.0	39.0	-	0.0	-	0.0	-	-	0.0	-	0.0	-	-
119.0	33.0	-	0.0	-	0.0	-	-	0.0	-	0.0	-	-
120.0	23.0	-	0.0	-	0.0	-	-	0.0	-	0.0	-	-
120.0	25.0	-	0.0	-	0.0	-	-	0.0	-	0.0	-	-
120.0	30.0	-	0.0	-	0.0	-	-	0.0	-	0.0	-	-
120.0	35.0	-	0.0	-	0.0	-	-	0.0	-	0.0	-	-
120.0	40.0	-	0.0	-	0.0	-	-	0.0	-	0.0	-	-
120.0	45.0	-	0.0	-	0.0	-	-	0.0	-	0.0	-	-
120.0	55.0	-	0.0	-	0.0	-	-	0.0	-	0.0	-	-
123.0	42.0	-	0.0	-	0.0	-	-	0.0	-	0.0	-	-
123.0	34.0	-	0.0	-	0.0	-	-	0.0	-	0.0	-	-
127.0	34.0	-	0.0	-	0.0	-	-	0.0	-	0.0	-	-
130.0	28.0	-	0.0	-	0.0	-	-	0.0	-	0.0	-	-
130.0	35.0	-	0.0	-	0.0	-	-	0.0	-	0.0	-	-
133.0	19.5	-	0.0	-	0.0	-	-	0.0	-	0.0	-	-
133.0	23.0	-	0.0	-	0.0	-	-	0.0	-	0.0	-	-
133.0	25.0	-	0.0	-	0.0	-	-	0.0	-	0.0	-	-
133.0	30.0	-	0.0	-	0.0	-	-	0.0	-	0.0	-	-

TABLE 4. (cont.)

*Hippoglossina stomata* (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
133.0	35.0	-	0.0	-	0.0	-	-	3.0	-	3.0	-	-
137.0	23.0	-	1.9	-	0.0	-	-	-	11.4	0.0	-	-
137.0	30.0	-	0.0	-	0.0	-	-	-	0.0	21.4	-	-
140.0	30.0	-	0.0	-	-	-	-	-	2.8	-	-	-
140.0	35.0	-	3.0	-	-	-	-	0.0	-	-	-	-

*Paralichthys californicus*

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
63.0	49.7	1.2	-	-	0.0	-	-	-	-	-	-	-
63.0	50.0	0.8	-	-	0.0	-	-	-	0.0	-	-	-
67.0	47.3	0.7	-	-	0.0	-	-	-	-	-	-	-
77.0	48.0	0.6	-	-	1.2	-	-	-	0.0	-	-	-
77.0	49.0	-	-	-	0.7	-	-	-	0.0	-	-	-
77.0	51.0	0.0	-	-	3.0	-	-	-	0.0	-	-	-
80.0	50.2	1.4	-	-	5.3	-	-	-	0.4	0.0	-	-
80.0	51.0	0.0	-	-	31.6	-	-	-	1.6	0.0	-	-
80.0	52.0	0.0	-	-	0.0	-	-	-	2.2	0.0	-	-
83.0	39.0	1.6	-	-	-	-	6.3	-	-	-	-	-
83.0	40.0	1.9	-	-	-	-	7.8	-	-	-	-	-
83.0	43.0	0.0	-	-	0.0	-	6.3	-	-	-	-	-
83.0	51.0	0.0	-	-	0.0	-	3.2	-	-	-	-	-
87.0	32.5	0.0	-	-	1.5	-	3.5	-	-	-	-	-
87.0	33.0	4.0	-	-	2.6	-	0.0	-	-	-	-	-
87.0	34.0	0.0	-	-	2.5	-	0.0	-	-	-	-	-
87.0	40.0	0.0	-	-	0.0	-	0.0	-	-	-	-	-
90.0	27.5	1.1	-	-	4.5	-	-	-	0.0	-	-	-
90.0	28.0	2.8	-	-	0.0	-	-	-	0.0	-	-	-
90.0	32.0	8.1	-	-	0.0	-	-	-	0.0	-	-	-
90.0	37.0	3.0	-	-	0.0	-	-	-	0.0	-	-	-
93.0	26.5	12.0	-	-	-	-	34.0	-	-	-	-	-
93.0	27.0	0.0	-	-	7.9	-	8.6	-	-	-	-	-
96.8	28.7	4.7	-	-	0.0	-	4.0	-	-	-	-	-
97.0	29.0	18.7	-	-	7.3	-	5.1	-	-	-	-	-
97.0	30.0	0.0	-	-	2.5	-	0.0	-	-	-	-	-
97.0	32.0	0.0	-	-	3.1	-	0.0	-	-	-	-	-
100.0	29.0	7.5	-	-	-	-	-	-	0.0	-	-	-
100.0	30.0	2.8	-	-	0.0	-	-	-	0.0	-	-	-
103.0	28.5	2.5	-	-	3.6	-	2.1	-	-	-	-	-
103.0	29.0	2.3	-	-	1.6	-	0.0	-	-	-	-	-
107.0	30.0	-	-	-	3.0	-	9.9	-	-	-	-	-
107.0	31.0	-	-	-	0.0	-	0.0	-	-	-	-	-
107.0	35.0	-	-	-	3.2	-	0.0	-	-	-	-	-
110.0	32.0	-	-	-	2.0	-	0.0	-	-	-	-	-
113.0	28.5	-	-	-	3.5	-	0.0	-	-	-	-	-

TABLE 4. (cont.)

*Paralichthys californicus* (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
1113.0	29.0	-	10.8	-	1.2	-	0.0	-	-	0.0	-	-
1113.0	30.0	-	0.0	-	0.0	-	0.0	-	-	0.0	-	-
1115.0	35.0	-	2.8	-	0.0	-	-	-	-	0.0	-	-
1117.0	24.8	-	1.0	-	0.0	-	-	-	-	0.0	-	-
1117.0	25.0	-	2.6	-	2.8	-	0.0	-	-	5.1	-	-
1117.0	26.0	-	0.0	-	6.0	-	0.0	-	-	0.0	-	-
1118.0	39.0	-	0.0	-	9.3	-	0.0	-	-	0.0	-	-
1119.0	33.0	-	0.0	-	0.0	-	-	-	-	2.2	-	-
120.0	22.5	-	3.3	-	0.0	-	-	-	-	0.0	-	-
120.0	23.0	-	12.5	-	2.8	-	-	-	-	0.0	-	-
120.0	24.0	-	21.2	-	18.7	-	0.0	-	-	0.0	-	-
120.0	25.0	-	23.7	-	5.3	-	0.0	-	-	0.0	-	-
120.0	30.0	-	0.0	-	2.8	-	8.2	-	-	0.0	-	-
120.0	35.0	-	0.0	-	5.6	-	0.0	-	-	0.0	-	-
120.0	40.0	-	0.0	-	1.9	-	0.0	-	-	0.0	-	-
123.0	36.0	-	0.0	-	2.4	-	-	-	-	0.0	-	-
127.0	33.0	-	2.8	-	0.0	-	0.0	-	-	4.1	-	-
130.0	25.5	-	0.0	-	0.0	-	-	-	-	0.0	-	-
130.0	26.0	-	1.0	-	0.0	-	-	-	-	0.0	-	-
133.0	19.5	-	1.7	-	0.0	-	-	-	-	6.6	-	-
133.0	21.0	-	38.0	-	0.0	-	-	-	-	6.1	-	-
133.0	30.0	-	0.0	-	3.1	-	0.0	-	-	0.0	-	-
133.0	50.0	-	0.0	-	0.0	-	-	-	-	2.6	-	-
137.0	20.0	-	0.0	-	0.0	-	-	-	-	0.4	-	-
137.0	21.0	-	1.3	-	0.0	-	-	-	-	4.1	-	-
137.0	22.0	-	0.0	-	6.8	-	-	-	-	34.2	-	-
140.0	28.8	-	0.7	-	-	-	-	-	-	0.0	-	-
140.0	29.0	-	0.0	-	-	-	-	-	-	2.0	-	-

*Syacium ovale*

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
133.0	30.0	-	0.0	-	0.0	-	-	0.0	-	-	8.6	-

*Xystreurus liolepis*

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
87.0	33.0	0.0	-	-	0.0	-	0.0	-	-	5.2	-	-
87.0	34.0	0.0	-	-	0.0	-	0.0	-	-	10.6	-	-
113.0	29.0	-	0.0	-	0.0	-	0.0	-	-	1.8	-	-
118.0	39.0	-	0.0	-	0.0	-	-	-	-	2.8	-	-
120.0	24.0	-	15.9	-	0.0	-	-	-	-	1.2	-	-
120.0	25.0	-	3.7	-	0.0	-	-	-	-	0.0	-	-
130.0	28.0	-	0.0	-	-	-	-	-	-	2.5	0.0	-

TABLE 4. (cont.)

*Glyptocephalus zachirus*

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
60.0	60.0	0.0	-	-	-	-	-	5.7	-	-	-	-
60.0	70.0	0.0	-	-	-	-	-	5.6	-	-	-	-
63.0	52.0	0.0	-	-	2.0	-	-	0.0	-	-	-	-
70.0	51.0	0.0	-	-	0.0	-	-	2.9	-	-	-	-

*Hypsopsetta guttulata*

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
87.0	32.5	0.0	-	-	0.0	-	-	0.0	-	1.7	-	-
87.0	40.0	0.0	-	-	0.0	-	-	0.0	-	3.0	-	-
90.0	27.5	0.0	-	-	0.0	-	-	0.0	-	1.3	-	-
93.0	26.5	4.8	-	-	-	-	-	0.0	-	-	-	-
93.0	27.0	2.8	-	-	0.0	-	-	0.0	-	0.0	-	-
93.0	30.0	3.0	-	-	0.0	-	-	0.0	-	2.2	-	-
97.0	30.0	0.0	-	-	2.5	-	-	0.0	-	0.0	-	-
130.0	28.0	-	2.1	-	0.0	-	-	0.0	-	0.0	-	-
133.0	23.0	-	2.7	-	0.0	-	-	0.0	-	0.0	-	-

*Lepidopsetta bilineata*

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
77.0	49.0	-	-	-	0.0	-	-	5.4	3.5	-	-	-
87.0	50.0	0.0	-	-	0.0	-	-	0.0	0.0	-	-	-

*Lyopsetta exilis*

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
60.0	70.0	0.0	-	-	-	-	-	-	-	5.6	-	-
63.0	55.0	0.0	-	-	-	-	-	1.9	-	2.8	-	-
63.0	60.0	0.0	-	-	-	-	-	5.0	-	0.0	-	-
67.0	49.0	0.0	-	-	-	-	-	0.0	-	3.1	-	-
67.0	50.0	0.0	-	-	-	-	-	2.7	-	0.0	-	-
67.0	55.0	0.0	-	-	-	-	-	2.7	-	0.0	-	-
67.0	60.0	9.0	-	-	-	-	-	0.0	-	0.0	-	-
67.0	70.0	-	-	-	-	-	-	8.5	-	0.0	-	-
70.0	50.0	2.9	-	-	-	-	-	0.0	-	0.0	-	-
70.0	52.0	0.0	-	-	-	-	-	5.2	-	0.0	-	-
70.0	65.0	-	-	-	-	-	-	2.7	-	0.0	-	-
77.0	47.6	0.0	-	-	-	-	-	0.6	-	0.0	-	-
77.0	48.0	0.3	-	-	-	-	-	0.0	-	0.0	-	-
77.0	51.0	-	-	-	-	-	-	3.0	-	0.0	-	-
80.0	52.0	3.3	-	-	-	-	-	20.0	-	0.0	-	-
80.0	55.0	-	-	-	-	-	-	0.0	-	0.0	-	-

TABLE 4. (cont.)

*Lyopsetta exilis* (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
80.0	80.0	0.0	-	-	3.2	-	-	0.0	-	0.0	-	-
82.0	47.0	0.0	-	-	6.2	-	0.0	-	0.0	-	-	-
83.0	43.0	0.0	-	-	5.6	-	0.0	-	0.0	-	-	-
87.0	35.0	0.0	-	-	6.8	-	0.0	-	0.0	-	-	-
90.0	27.5	0.0	-	-	1.3	-	0.0	-	0.0	-	-	-
93.0	27.0	0.0	-	-	4.0	-	0.0	-	0.0	-	-	-
93.0	28.0	0.0	-	-	3.3	-	0.0	-	0.0	-	-	-
100.0	30.0	0.0	-	-	9.6	-	0.0	-	0.0	-	-	-
103.0	28.5	0.0	-	-	1.8	-	0.0	-	0.0	-	-	-
103.0	30.0	0.0	-	-	3.1	-	0.0	-	0.0	-	-	-
113.0	30.0	-	0.0	-	2.8	-	0.0	-	0.0	-	-	-
117.0	30.0	-	0.0	-	0.0	-	3.8	-	2.9	-	-	-
117.0	60.0	-	0.0	-	24.7	-	14.0	-	0.0	-	-	-
118.0	39.0	-	0.0	-	-	-	-	-	-	-	-	-

*Microstomus pacificus*

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
60.0	100.0	0.0	-	-	-	-	-	2.6	-	-	-	-
60.0	120.0	0.0	-	-	-	-	-	2.7	-	-	-	-
63.0	60.0	0.0	-	-	2.5	-	-	0.0	-	-	-	-
63.0	80.0	-	-	-	5.9	-	-	-	-	-	-	-
67.0	55.0	0.0	-	-	2.7	-	-	0.0	-	-	-	-
67.0	65.0	-	-	-	11.6	-	-	-	-	-	-	-
70.0	60.0	0.0	-	-	0.0	-	-	2.7	-	-	-	-
70.0	65.0	-	-	-	2.7	-	-	-	-	-	-	-
77.0	65.0	-	-	-	2.5	-	-	-	-	-	-	-
80.0	65.0	0.0	-	-	3.4	-	-	0.0	-	-	-	-
83.0	65.0	0.0	-	-	2.9	-	-	3.3	-	-	-	-
83.0	70.0	0.0	-	-	0.0	-	-	6.7	-	-	-	-
87.0	80.0	0.0	-	-	0.0	-	-	3.3	-	-	-	-
93.0	50.0	0.0	-	-	0.0	-	-	3.3	-	-	-	-
97.0	40.0	0.0	-	-	0.0	-	-	3.3	-	-	-	-

*Parophrys vetulus*

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
60.0	49.5	6.0	-	-	-	-	-	-	-	-	-	-
60.0	51.0	30.8	-	-	-	-	-	-	-	-	-	-
60.0	52.0	15.8	-	-	-	-	-	-	-	-	-	-
60.0	55.0	6.2	-	-	-	-	-	-	-	-	-	-
63.0	49.7	8.1	-	-	-	-	-	-	-	-	-	-
63.0	50.0	28.7	-	-	-	-	-	-	-	-	-	-
63.0	51.0	31.6	-	-	-	-	-	-	-	-	-	-

TABLE 4. (cont.)

*Parophrys vetulus* (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
63.0	52.0	0.0	-	-	6.2	-	-	0.0	-	-	-	-
63.0	55.0	0.0	-	-	1.9	-	-	0.0	-	-	-	-
70.0	51.0	6.1	-	-	2.8	-	-	0.0	-	-	-	-
70.0	52.0	6.3	-	-	0.0	-	-	0.0	-	-	-	-
70.0	53.0	3.0	-	-	0.0	-	-	0.0	-	-	-	-
73.0	49.4	0.3	-	-	0.0	-	-	0.0	-	-	-	-
73.0	51.0	0.0	-	-	5.6	-	-	0.0	-	-	-	-
77.0	47.6	0.3	-	-	0.0	-	-	0.0	-	-	-	-
77.0	48.0	0.6	-	-	0.0	-	-	0.0	-	-	-	-
77.0	49.0	-	-	-	4.7	-	-	0.0	-	-	-	-
77.0	51.0	3.0	-	-	33.1	-	-	0.0	-	-	-	-
77.0	50.2	1.8	-	-	0.8	-	-	0.9	-	-	-	-
80.0	51.0	15.8	-	-	24.6	-	-	1.6	-	-	-	-
80.0	52.0	0.0	-	-	146.5	-	-	0.0	-	-	-	-
80.0	55.0	3.1	-	-	9.6	-	-	0.0	-	-	-	-
80.0	60.0	0.0	-	-	6.8	-	-	0.0	-	-	-	-
80.0	65.0	0.0	-	-	3.4	-	-	0.0	-	-	-	-
82.0	47.0	8.6	-	-	0.0	-	-	2.7	-	-	-	-
83.0	40.0	1.9	-	-	0.0	-	-	0.0	-	-	-	-
83.0	43.0	23.7	-	-	8.4	-	-	0.0	-	-	-	-
83.0	51.0	4.7	-	-	0.0	-	-	0.0	-	-	-	-
87.0	32.5	0.0	-	-	2.2	-	-	0.0	-	-	-	-
87.0	33.0	0.0	-	-	5.3	-	-	2.7	-	-	-	-
87.0	34.0	24.0	-	-	2.5	-	-	0.0	-	-	-	-
87.0	35.0	20.6	-	-	17.1	-	-	0.0	-	-	-	-
87.0	40.0	2.8	-	-	0.0	-	-	0.0	-	-	-	-
87.0	55.0	8.9	-	-	0.0	-	-	0.0	-	-	-	-
90.0	27.5	1.7	-	-	0.6	-	-	0.0	-	-	-	-
90.0	28.0	0.0	-	-	6.1	-	-	0.0	-	-	-	-
90.0	32.0	0.0	-	-	0.0	-	-	0.0	-	-	-	-
90.0	45.0	2.5	-	-	0.0	-	-	0.0	-	-	-	-
93.0	27.0	0.0	-	-	11.9	-	-	2.9	-	-	-	-
93.0	28.0	3.1	-	-	3.3	-	-	0.0	-	-	-	-
93.0	35.0	0.0	-	-	0.0	-	-	3.0	-	-	-	-
96.8	28.7	1.2	-	-	0.0	-	-	0.0	-	-	-	-
97.0	29.0	1.9	-	-	12.1	-	-	0.0	-	-	-	-
97.0	30.0	2.2	-	-	5.1	-	-	3.4	-	-	-	-
97.0	32.0	0.0	-	-	18.6	-	-	2.9	-	-	-	-
100.0	29.0	7.5	-	-	-	-	-	0.0	-	-	-	-
100.0	30.0	2.8	-	-	-	-	-	3.2	-	-	-	-
103.0	35.0	0.0	-	-	-	-	-	0.0	-	-	-	-
110.0	40.0	-	-	-	-	-	-	2.8	-	-	-	-
113.0	28.5	-	-	-	-	-	-	1.8	-	-	-	-
113.0	29.0	-	-	-	-	-	-	9.0	-	-	-	-
113.0	30.0	-	-	-	-	-	-	12.2	-	-	-	-
117.0	25.0	-	-	-	-	-	-	0.0	-	-	-	-
		2.6	-	-	-	-	-	14.8	-	-	-	-
		0.0	-	-	-	-	-	14.8	-	-	-	-

TABLE 4. (cont.)

<i>Parophrys vetulus</i> (cont.)												
STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
117.0	26.0	-	8.4	-	0.0	-	-	0.0	-	0.0	-	-
120.0	30.0	-	92.8	-	2.8	-	-	0.0	-	0.0	-	-
120.0	35.0	-	24.4	-	2.8	-	-	0.0	-	0.0	-	-
127.0	32.6	-	0.0	-	3.3	-	-	0.0	-	0.0	-	-
<i>Pleuronichthys</i> spp.												
STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
137.0	30.0	-	0.0	-	3.1	-	-	0.0	0.0	-	-	-
<i>Pleuronichthys coenosus</i>												
STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
82.0	47.0	0.0	-	-	0.0	-	-	2.7	-	0.0	-	-
83.0	51.0	0.0	-	-	0.0	-	-	3.2	-	0.0	-	-
90.0	37.0	0.0	-	-	0.0	-	-	2.1	-	0.0	-	-
103.0	30.0	0.0	-	-	0.0	-	-	2.6	-	0.0	-	-
110.0	35.0	-	0.0	-	0.0	-	-	-	2.8	0.0	-	-
<i>Pleuronichthys decurrens</i>												
STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
60.0	70.0	3.2	-	-	-	-	-	0.0	-	-	-	-
63.0	52.0	2.9	-	-	0.0	-	-	0.0	-	-	-	-
80.0	80.0	0.0	-	-	3.2	-	-	0.0	-	0.0	-	-
87.0	55.0	0.0	-	-	0.0	-	-	3.3	-	-	-	-
<i>Pleuronichthys ritteri</i>												
STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
93.0	26.5	0.0	-	-	-	-	-	5.0	-	-	3.2	-
96.8	28.7	0.0	-	0.0	-	-	-	2.0	-	-	0.0	-
107.0	30.0	-	0.0	-	0.0	-	-	1.1	-	-	0.0	-
120.0	24.0	-	5.3	-	0.0	-	-	-	0.0	-	0.0	-
120.0	25.0	-	7.3	-	0.0	-	-	-	0.0	-	0.0	-
120.0	30.0	-	0.0	-	0.0	-	-	-	2.7	-	0.0	-
120.0	35.0	-	0.0	-	0.0	-	-	-	2.6	-	0.0	-
127.0	32.6	-	0.0	-	0.0	-	-	-	1.4	-	0.0	-

TABLE 4. (cont.)

*Pleuronichthys verticalis*

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
67.0	49.0	0.0	-	-	0.0	-	-	3.1	-	-	-	-
77.0	47.6	0.8	-	-	0.0	-	-	0.0	-	-	-	-
77.0	48.0	1.4	-	-	0.0	-	-	0.0	-	-	-	-
77.0	49.0	-	-	-	2.0	-	-	0.0	-	-	-	-
77.0	51.0	0.0	-	-	6.0	-	-	0.0	-	-	-	-
80.0	50.2	1.4	-	-	0.0	-	-	3.1	0.0	-	-	-
80.0	51.0	3.2	-	-	3.5	-	-	3.2	0.0	-	-	-
80.0	52.0	0.0	-	-	13.3	-	-	2.2	2.7	-	-	-
80.0	52.0	0.0	-	-	0.0	-	-	0.0	0.0	-	-	-
82.0	60.0	0.0	-	-	3.1	-	-	5.4	-	-	-	-
82.0	47.0	0.0	-	-	-	-	-	1.3	-	-	-	-
83.0	39.0	0.0	-	-	-	-	-	13.9	-	-	-	-
83.0	40.0	1.9	-	-	0.0	-	-	5.6	6.3	-	-	-
83.0	43.0	0.0	-	-	-	-	-	0.0	3.2	-	-	-
83.0	51.0	0.0	-	-	-	-	-	3.7	1.7	-	-	-
87.0	32.5	0.0	-	-	-	-	-	2.6	5.4	-	-	-
87.0	33.0	2.0	-	-	-	-	-	0.0	0.0	-	-	-
87.0	34.0	0.0	-	-	-	-	-	3.4	0.0	-	-	-
87.0	45.0	0.0	-	-	-	-	-	0.0	0.0	-	-	-
90.0	27.5	0.6	-	-	-	-	-	0.0	0.0	-	-	-
90.0	28.0	0.0	-	-	-	-	-	0.0	0.0	-	-	-
90.0	32.0	2.7	-	-	-	-	-	2.5	-	-	-	-
93.0	26.5	0.0	-	-	-	-	-	2.5	-	-	-	-
93.0	27.0	0.0	-	-	4.0	-	-	0.0	0.0	-	-	-
93.0	28.0	0.0	-	-	0.0	-	-	0.0	3.2	-	-	-
93.0	35.0	0.0	-	-	0.0	-	-	0.0	3.0	-	-	-
96.8	28.7	7.1	-	-	0.0	-	-	2.0	2.0	-	-	-
97.0	29.0	16.8	-	-	0.0	-	-	17.9	0.0	-	-	-
97.0	30.0	0.0	-	-	0.0	-	-	2.8	2.8	-	-	-
100.0	29.0	0.0	-	-	0.0	-	-	0.0	0.0	-	-	-
100.0	35.0	0.0	-	-	0.0	-	-	0.0	0.0	-	-	-
103.0	28.5	1.3	-	-	0.0	-	-	0.0	0.0	-	-	-
103.0	29.0	0.0	-	-	1.5	-	-	1.5	5.0	-	-	-
107.0	30.0	-	-	-	0.0	-	-	0.0	1.1	-	-	-
107.0	31.0	-	-	-	0.0	-	-	0.0	3.8	-	-	-
107.0	32.0	-	-	-	3.1	-	-	0.0	0.0	-	-	-
110.0	32.0	-	-	-	0.0	-	-	0.0	0.0	-	-	-
113.0	28.5	-	-	-	0.0	-	-	0.0	0.0	-	-	-
113.0	29.0	-	-	-	2.2	-	-	3.7	1.8	-	-	-
113.0	30.0	-	-	-	0.0	-	-	0.0	0.0	-	-	-
117.0	24.8	-	-	-	0.0	-	-	0.0	0.0	-	-	-
117.0	25.0	-	-	-	0.0	-	-	0.0	0.0	-	-	-
117.0	26.0	-	-	-	0.0	-	-	0.0	0.0	-	-	-
117.0	30.0	-	-	-	0.0	-	-	0.0	0.0	-	-	-
118.0	39.0	-	-	-	0.0	-	-	0.0	0.0	-	-	-
119.0	33.0	-	-	-	0.0	-	-	2.9	2.9	-	-	-
120.0	23.0	-	-	-	0.0	-	-	0.0	0.0	-	-	-

TABLE 4. (cont.)

*Pleuronichthys verticalis* (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
120.0	24.0	-	0.0	-	0.0	-	-	1.2	-	0.0	-	-
120.0	25.0	-	0.0	-	2.7	-	-	6.0	-	0.0	-	-
120.0	30.0	-	2.9	-	0.0	-	-	2.7	-	0.0	-	-
120.0	35.0	-	6.1	-	0.0	-	-	2.6	-	0.0	-	-
120.0	40.0	-	0.0	-	0.0	-	-	2.6	-	0.0	-	-
123.0	36.0	-	0.0	-	4.8	-	-	1.5	-	0.0	-	-
123.5	35.6	-	0.0	-	1.2	-	-	0.0	-	0.0	-	-
127.0	34.0	-	0.0	-	0.0	-	-	2.4	-	0.0	-	-
133.0	30.0	-	0.0	-	0.0	-	-	3.0	-	0.0	-	-

*Psettichthys melanostictus*

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
60.0	49.5	0.9	-	-	-	-	-	-	-	-	-	-
63.0	49.7	0.6	-	-	-	-	-	-	-	-	-	-
63.0	50.0	2.5	-	-	0.0	-	-	-	-	-	-	-
63.0	51.0	5.3	-	-	3.2	-	-	0.0	-	-	-	-
63.0	52.0	0.0	-	-	0.0	-	-	0.0	-	-	-	-
73.0	51.0	6.2	-	-	2.0	-	-	0.0	-	-	-	-
77.0	47.6	0.3	-	-	0.0	-	-	0.0	-	-	-	-
77.0	48.0	3.2	-	-	0.0	-	-	0.0	-	-	-	-

*Sympfururus* spp.

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
83.0	43.0	0.0	-	-	0.0	-	-	0.0	-	-	31.7	-
87.0	32.5	0.0	-	-	0.0	-	-	0.0	-	-	1.7	-
87.0	34.0	0.0	-	-	0.0	-	-	0.0	-	-	7.9	-
87.0	35.0	0.0	-	-	0.0	-	-	0.0	-	-	9.1	-
87.0	40.0	0.0	-	-	0.0	-	-	0.0	-	-	15.3	-
87.0	45.0	0.0	-	-	0.0	-	-	0.0	-	-	51.2	-
87.0	50.0	0.0	-	-	0.0	-	-	0.0	-	-	10.6	-
93.0	30.0	0.0	-	-	0.0	-	-	0.0	-	-	3.0	-
97.0	40.0	0.0	-	-	0.0	-	-	0.0	-	-	3.0	-
110.0	35.0	-	0.0	-	0.0	-	-	0.0	-	-	3.0	-
110.0	55.0	-	0.0	-	0.0	-	-	0.0	-	-	2.8	0.0
113.0	28.5	-	0.0	-	0.0	-	-	0.0	-	-	1.1	-
113.0	29.0	-	0.0	-	0.0	-	-	0.0	-	-	39.2	-
113.0	30.0	-	0.0	-	0.0	-	-	0.0	-	-	35.9	0.0
113.0	35.0	-	0.0	-	0.0	-	-	0.0	-	-	9.0	-
113.0	50.0	-	0.0	-	0.0	-	-	0.0	-	-	3.0	-
113.0	55.0	-	0.0	-	0.0	-	-	0.0	-	-	3.0	-
115.0	35.0	-	0.0	-	0.0	-	-	0.0	-	-	10.4	3.1
117.0	24.8	-	0.0	-	0.0	-	-	0.0	-	-	0.0	-

TABLE 4. (cont.)

*Sympodus spp.* (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
117.0	26.0	-	0.0	-	0.0	-	-	-	0.0	-	128.9	-
117.0	30.0	-	0.0	-	0.0	-	-	-	0.0	-	62.4	-
117.0	35.0	-	0.0	-	0.0	-	-	-	0.0	-	46.7	-
117.0	40.0	-	0.0	-	0.0	-	-	-	0.0	-	3.1	-
117.0	45.0	-	0.0	-	0.0	-	-	-	0.0	-	31.8	-
117.0	50.0	-	0.0	-	0.0	-	-	-	0.0	-	2.8	-
118.0	39.0	-	0.0	-	0.0	-	-	-	0.0	-	100.6	-
119.0	33.0	-	0.0	-	0.0	-	-	-	0.0	-	211.6	-
120.0	22.5	-	0.0	-	0.0	-	-	-	0.0	-	1.6	-
120.0	23.0	-	0.0	-	0.0	-	-	-	0.0	-	15.0	-
120.0	24.0	-	0.0	-	0.0	-	-	-	0.0	-	61.9	-
120.0	25.0	-	0.0	-	0.0	-	-	-	0.0	-	53.6	-
120.0	30.0	-	0.0	-	0.0	-	-	-	0.0	-	93.3	-
120.0	35.0	-	0.0	-	0.0	-	-	-	0.0	-	30.1	-
120.0	40.0	-	0.0	-	0.0	-	-	-	0.0	-	24.6	-
120.0	45.0	-	0.0	-	0.0	-	-	-	0.0	-	5.2	-
120.0	55.0	-	0.0	-	0.0	-	-	-	0.0	-	20.0	-
120.0	60.0	-	0.0	-	0.0	-	-	-	0.0	-	6.0	-
123.0	36.0	-	0.0	-	0.0	-	-	-	0.0	-	0.0	-
123.0	42.0	-	0.0	-	0.0	-	-	-	0.0	-	0.0	-
123.0	45.0	-	0.0	-	0.0	-	-	-	0.0	-	0.0	-
123.0	60.0	-	0.0	-	0.0	-	-	-	0.0	-	0.0	-
127.0	34.0	-	0.0	-	0.0	-	-	-	0.0	-	1.5	-
127.0	40.0	-	0.0	-	0.0	-	-	-	0.0	-	3.2	-
127.0	50.0	-	0.0	-	0.0	-	-	-	0.0	-	0.0	-
130.0	25.5	-	0.0	-	0.0	-	-	-	0.0	-	3.3	-
130.0	26.0	-	0.0	-	0.0	-	-	-	0.0	-	3.3	-
130.0	28.0	-	0.0	-	0.0	-	-	-	0.0	-	32.2	-
130.0	30.0	-	0.0	-	0.0	-	-	-	0.0	-	2.0	-
130.0	35.0	-	0.0	-	0.0	-	-	-	0.0	-	42.7	-
130.0	40.0	-	0.0	-	0.0	-	-	-	0.0	-	6.0	-
133.0	21.0	-	0.0	-	0.0	-	-	-	0.0	-	8.9	-
133.0	23.0	-	0.0	-	0.0	-	-	-	0.0	-	3.3	-
133.0	25.0	-	0.0	-	0.0	-	-	-	0.0	-	32.2	-
133.0	30.0	-	0.0	-	0.0	-	-	-	0.0	-	2.0	-
133.0	35.0	-	0.0	-	0.0	-	-	-	0.0	-	5.7	-
137.0	21.0	-	0.0	-	0.0	-	-	-	0.0	-	20.7	-
137.0	22.0	-	0.0	-	0.0	-	-	-	0.0	-	0.0	-
137.0	23.0	-	0.0	-	0.0	-	-	-	0.0	-	19.3	-
137.0	30.0	-	0.0	-	0.0	-	-	-	0.0	-	6.1	-
137.0	35.0	-	0.0	-	0.0	-	-	-	0.0	-	13.3	-
137.0	40.0	-	0.0	-	0.0	-	-	-	0.0	-	2.8	-
											0.0	27.8
											0.0	3.0

TABLE 4. (cont.)

Soleidae												
STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
137.0	20.0	-	0.0	-	0.0	-	-	-	0.0	1.9	-	-
Tetraodontidae												
STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
133.0	21.0	-	0.0	-	0.0	-	-	-	0.0	2.3	-	-
140.0	28.8	-	0.0	-	-	-	-	-	1.0	-	-	-
140.0	29.0	-	0.0	-	-	-	-	-	2.0	-	-	-
Disintegrated fish larva												
STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
60.0	49.2	-	3.4	-	-	-	-	-	1.0	-	-	-
60.0	49.5	-	2.6	-	-	-	-	-	-	-	-	-
60.0	50.0	-	9.7	-	-	-	-	-	0.0	0.0	-	-
60.0	51.0	-	6.2	-	-	-	-	-	2.6	-	-	-
60.0	55.0	-	3.2	-	-	-	-	-	0.0	0.0	-	-
60.0	70.0	-	0.0	-	-	-	-	-	2.6	-	-	-
60.0	140.0	-	0.0	-	-	-	-	-	2.7	-	-	-
60.0	180.0	-	0.0	-	-	-	-	-	2.7	-	-	-
60.0	200.0	-	0.0	-	-	-	-	-	0.0	0.0	-	-
63.0	50.0	-	0.0	-	-	-	-	-	0.0	0.0	-	-
63.0	51.0	-	0.0	-	-	-	-	-	0.0	0.0	-	-
63.0	52.0	-	0.0	-	-	-	-	-	0.0	0.0	-	-
63.0	55.0	-	3.2	-	-	-	-	-	0.0	0.0	-	-
63.0	60.0	-	0.0	-	-	-	-	-	0.0	0.0	-	-
63.0	65.0	-	-	-	-	-	-	-	-	-	-	-
63.0	80.0	-	-	-	-	-	-	-	-	-	-	-
67.0	47.3	-	0.0	-	-	-	-	-	-	-	-	-
67.0	49.0	-	3.2	-	-	-	-	-	0.0	0.0	-	-
70.0	51.0	-	3.1	-	-	-	-	-	0.0	2.9	-	-
70.0	52.0	-	12.6	-	-	-	-	-	0.0	2.9	-	-
70.0	53.0	-	3.0	-	-	-	-	-	0.0	2.9	-	-
70.0	55.0	-	2.9	-	-	-	-	-	2.7	-	-	-
70.0	65.0	-	-	-	-	-	-	-	2.7	-	-	-
70.0	90.0	-	5.7	-	-	-	-	-	-	2.8	-	-
70.0	100.0	-	0.0	-	-	-	-	-	-	2.8	-	-
70.0	200.0	-	3.1	-	-	-	-	-	-	5.3	-	-
73.0	49.4	-	0.3	-	-	-	-	-	0.0	0.0	-	-
73.0	50.0	-	4.6	-	-	-	-	-	0.0	0.0	-	-
73.0	60.0	-	2.8	-	-	-	-	-	5.6	0.0	-	-
77.0	47.6	-	0.8	-	-	-	-	-	2.5	0.0	-	-
77.0	48.0	-	0.9	-	-	-	-	-	1.2	0.0	-	-
77.0	49.0	-	-	-	-	-	-	-	-	1.3	-	-

TABLE 4. (cont.)

## Disintegrated fish larva (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
77.0	51.0	5.9	-	-	9.0	-	-	2.9	-	-	-	-
77.0	55.0	8.6	-	-	0.0	-	-	3.1	-	-	-	-
77.0	57.0	0.0	-	-	7.4	-	-	0.0	-	-	-	-
77.0	65.0	-	-	-	2.5	-	-	-	-	-	-	-
80.0	55.0	21.8	-	-	12.8	-	-	0.0	0.0	-	-	-
80.0	65.0	0.0	-	-	0.0	-	-	3.0	0.0	-	-	-
80.0	70.0	0.0	-	-	10.2	-	-	0.0	0.0	-	-	-
80.0	90.0	0.0	-	-	-	-	-	2.8	-	-	-	-
80.0	160.0	5.8	-	-	-	-	-	-	-	-	-	-
80.0	200.0	0.0	-	-	-	-	-	-	-	-	-	-
82.0	47.0	0.0	-	-	0.0	-	-	2.5	-	-	-	-
83.0	39.0	0.0	-	-	-	-	-	3.9	-	-	-	-
83.0	40.0	1.9	-	-	0.8	-	-	0.0	0.0	-	-	-
83.0	43.0	20.7	-	-	0.0	-	-	6.3	-	-	-	-
83.0	51.0	0.0	-	-	5.2	-	-	0.0	0.0	-	-	-
83.0	55.0	8.7	-	-	0.0	-	-	0.0	0.0	-	-	-
83.0	60.0	0.0	-	-	0.0	-	-	3.5	-	-	-	-
83.0	70.0	0.0	-	-	6.3	-	-	0.0	0.0	-	-	-
83.0	80.0	3.5	-	-	0.0	-	-	0.0	0.0	-	-	-
83.0	90.0	6.1	-	-	0.0	-	-	0.0	0.0	-	-	-
87.0	32.5	12.6	-	-	0.7	-	-	0.0	0.0	-	-	-
87.0	33.0	17.9	-	-	5.3	-	-	0.0	0.0	-	-	-
87.0	34.0	10.7	-	-	0.0	-	-	0.0	0.0	-	-	-
87.0	40.0	8.6	-	-	0.0	-	-	0.0	0.0	-	-	-
87.0	45.0	3.1	-	-	3.4	-	-	0.0	0.0	-	-	-
87.0	55.0	3.0	-	-	0.0	-	-	0.0	0.0	-	-	-
87.0	65.0	0.0	-	-	5.4	-	-	0.0	0.0	-	-	-
87.0	70.0	0.0	-	-	3.2	-	-	0.0	0.0	-	-	-
87.0	100.0	-	-	-	3.1	-	-	0.0	0.0	-	-	-
90.0	28.0	16.7	-	-	0.0	-	-	0.0	0.0	-	-	-
90.0	32.0	37.8	-	-	3.0	-	-	0.0	0.0	-	-	-
90.0	37.0	6.0	-	-	3.1	-	-	2.1	-	-	-	-
90.0	45.0	0.0	-	-	6.3	-	-	0.0	0.0	-	-	-
90.0	53.0	16.1	-	-	3.3	-	-	0.0	0.0	-	-	-
90.0	60.0	6.0	-	-	0.0	-	-	0.0	0.0	-	-	-
90.0	65.0	0.0	-	-	5.9	-	-	5.3	-	-	-	-
90.0	70.0	6.2	-	-	0.0	-	-	0.0	0.0	-	-	-
90.0	90.0	0.0	-	-	12.8	-	-	0.0	0.0	-	-	-
90.0	100.0	8.3	-	-	3.1	-	-	2.7	-	-	-	-
90.0	120.0	2.7	-	-	-	-	-	0.0	0.0	-	-	-
90.0	160.0	2.6	-	-	-	-	-	0.0	0.0	-	-	-
90.0	180.0	2.7	-	-	-	-	-	0.0	0.0	-	-	-
93.0	26.5	0.0	-	-	-	-	-	4.0	-	-	-	-
93.0	27.0	0.0	-	-	-	-	-	0.0	-	-	-	-
93.0	28.0	0.0	-	-	-	-	-	6.4	-	-	-	-
93.0	30.0	8.6	-	-	-	-	-	3.1	-	-	-	-

TABLE 4. (cont.)

## Disintegrated fish larva (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
93.0	35.0	15.0	-	0.0	-	-	0.0	-	-	0.0	-	-
93.0	55.0	0.0	-	-	4.2	-	-	0.0	-	0.0	-	-
93.0	65.0	6.3	-	0.0	0.0	-	0.0	-	0.0	0.0	-	-
93.0	70.0	5.9	-	0.0	0.0	-	0.0	-	0.0	0.0	-	-
93.0	80.0	0.0	-	6.3	0.0	-	0.0	-	0.0	0.0	-	-
93.0	90.0	0.0	-	6.7	0.0	-	0.0	-	0.0	0.0	-	-
93.0	100.0	-	2.8	-	-	-	-	-	-	0.0	-	-
96.8	28.7	5.9	0.0	-	-	0.0	-	0.0	-	0.0	-	-
97.0	29.0	1.9	0.0	-	12.6	-	0.0	-	0.0	0.0	-	-
97.0	30.0	6.5	-	6.2	-	0.0	-	0.0	-	0.0	-	-
97.0	32.0	3.1	-	6.3	-	0.0	-	0.0	-	0.0	-	-
97.0	35.0	0.0	-	6.3	-	0.0	-	0.0	-	0.0	-	-
97.0	40.0	3.2	9.6	-	-	0.0	-	0.0	-	0.0	-	-
97.0	45.0	9.1	0.0	-	-	0.0	-	0.0	-	0.0	-	-
97.0	50.0	12.6	-	6.9	-	0.0	-	0.0	-	0.0	-	-
97.0	65.0	0.0	-	-	0.0	-	17.9	-	0.0	0.0	-	-
97.0	75.0	-	-	-	-	2.9	-	2.9	-	2.8	-	-
97.0	85.0	-	3.8	-	-	3.2	-	3.2	-	2.5	-	-
100.0	29.0	-	5.6	-	-	0.0	-	0.0	-	2.8	-	-
100.0	30.0	-	0.0	-	3.0	-	9.1	-	0.0	0.0	-	-
100.0	35.0	-	0.0	-	3.0	-	3.3	-	0.0	0.0	-	-
100.0	40.0	0.0	-	0.0	-	3.0	-	10.1	-	0.0	-	-
100.0	55.0	6.1	-	-	-	0.0	-	0.0	-	0.0	-	-
100.0	60.0	3.0	-	-	-	0.0	-	0.0	-	0.0	-	-
100.0	75.0	-	0.0	-	-	0.0	-	0.0	-	0.0	-	-
100.0	80.0	0.0	-	-	6.9	-	6.9	-	3.0	6.8	-	-
100.0	85.0	-	-	5.9	-	-	1.6	-	0.0	0.0	-	-
100.0	100.0	0.0	-	-	-	0.0	0.0	0.0	-	0.0	-	-
103.0	29.0	0.0	-	-	6.1	-	6.1	-	0.0	0.0	-	-
103.0	30.0	0.0	-	2.9	-	0.0	0.0	0.0	-	0.0	-	-
103.0	35.0	-	2.9	-	-	0.0	0.0	0.0	-	0.0	-	-
103.0	40.0	8.8	-	-	-	0.0	0.0	0.0	-	0.0	-	-
103.0	45.0	0.0	-	-	-	3.0	-	3.0	-	0.0	-	-
103.0	55.0	0.0	-	-	-	3.3	-	3.3	-	0.0	-	-
103.0	65.0	0.0	-	-	-	3.2	-	3.2	-	0.0	-	-
103.0	75.0	-	-	-	-	0.0	0.0	0.0	-	0.0	-	-
103.0	80.0	-	-	-	-	3.0	-	3.0	-	0.0	-	-
103.0	90.0	-	-	-	-	3.0	-	6.3	-	0.0	-	-
107.0	30.0	-	-	-	-	1.5	-	1.5	-	0.0	-	-
107.0	31.0	-	-	-	-	6.5	-	6.5	-	0.0	-	-
107.0	32.0	-	-	-	-	3.1	-	9.2	-	0.0	-	-
107.0	50.0	-	-	-	-	5.9	-	5.9	-	0.0	-	-
107.0	60.0	-	-	-	-	3.0	-	3.0	-	0.0	-	-
107.0	70.0	-	-	-	-	3.2	-	3.2	-	0.0	-	-
107.0	80.0	-	-	-	-	2.9	-	2.9	-	0.0	-	-
107.0	90.0	-	-	-	-	3.1	-	3.1	-	0.0	-	-

TABLE 4. (cont.)

## Disintegrated fish larva (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
110.0	32.0	-	0.0	-	0.0	-	-	-	0.0	5.2	-	-
110.0	35.0	-	0.0	-	0.0	-	-	-	17.1	0.0	-	-
110.0	40.0	-	0.0	-	0.0	-	-	-	2.8	0.0	-	-
110.0	55.0	-	2.7	-	5.9	-	-	-	0.0	0.0	-	-
110.0	65.0	-	0.0	-	2.9	-	-	-	0.0	0.0	-	-
110.0	70.0	-	5.6	-	0.0	-	-	-	0.0	-	-	-
110.0	85.0	-	-	-	3.1	-	-	-	-	-	-	-
110.0	90.0	-	-	-	3.1	-	-	-	-	-	-	-
110.0	100.0	-	-	-	5.3	-	-	-	-	-	-	-
110.0	120.0	-	-	-	5.2	-	-	-	-	-	-	-
110.0	140.0	-	-	-	2.6	-	-	-	-	-	-	-
110.0	160.0	-	-	-	0.0	-	-	-	-	-	-	-
113.0	28.5	-	-	-	5.6	-	-	-	-	-	-	-
113.0	29.5	-	-	-	0.0	-	-	-	-	-	-	-
113.0	30.0	-	-	-	0.0	-	-	-	-	-	-	-
113.0	35.0	-	-	-	0.0	-	-	-	-	-	-	-
113.0	40.0	-	-	-	0.0	-	-	-	-	-	-	-
113.0	50.0	-	-	-	6.1	-	-	-	-	-	-	-
113.0	60.0	-	-	-	0.0	-	-	-	-	-	-	-
113.0	65.0	-	-	-	0.0	-	-	-	-	-	-	-
113.0	70.0	-	-	-	0.0	-	-	-	-	-	-	-
113.0	80.0	-	-	-	2.8	-	-	-	-	-	-	-
115.0	35.0	-	-	-	2.8	-	-	-	-	-	-	-
117.0	26.0	-	-	-	2.8	-	-	-	-	-	-	-
117.0	35.0	-	-	-	0.0	-	-	-	-	-	-	-
117.0	40.0	-	-	-	0.0	-	-	-	-	-	-	-
117.0	45.0	-	-	-	0.0	-	-	-	-	-	-	-
117.0	55.0	-	-	-	0.0	-	-	-	-	-	-	-
117.0	60.0	-	-	-	0.0	-	-	-	-	-	-	-
117.0	75.0	-	-	-	0.0	-	-	-	-	-	-	-
117.0	85.0	-	-	-	2.9	-	-	-	-	-	-	-
117.0	90.0	-	-	-	0.0	-	-	-	-	-	-	-
118.0	39.0	-	-	-	8.8	-	-	-	-	-	-	-
119.0	33.0	-	-	-	0.0	-	-	-	-	-	-	-
120.0	22.5	-	-	-	0.0	-	-	-	-	-	-	-
120.0	23.0	-	-	-	0.0	-	-	-	-	-	-	-
120.0	24.0	-	-	-	0.0	-	-	-	-	-	-	-
120.0	25.0	-	-	-	0.0	-	-	-	-	-	-	-
120.0	30.0	-	-	-	8.7	-	-	-	-	-	-	-
120.0	35.0	-	-	-	9.1	-	-	-	-	-	-	-
120.0	40.0	-	-	-	2.7	-	-	-	-	-	-	-
120.0	45.0	-	-	-	1.9	-	-	-	-	-	-	-
120.0	50.0	-	-	-	18.5	-	-	-	-	-	-	-
120.0	55.0	-	-	-	11.9	-	-	-	-	-	-	-
120.0	60.0	-	-	-	3.0	-	-	-	-	-	-	-

TABLE 4. (cont.)

## Disintegrated fish larva (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
120.0	70.0	-	2.9	-	9.9	-	-	-	0.0	-	-	-
120.0	80.0	-	0.0	-	0.0	-	-	-	32.3	-	-	-
120.0	120.0	-	2.9	-	2.4	-	-	-	0.0	-	9.5	-
123.0	36.0	-	0.0	-	1.6	-	-	-	0.0	-	0.0	-
123.0	42.0	-	0.0	-	0.0	-	-	-	5.7	-	0.0	-
123.0	50.0	-	0.0	-	0.0	-	-	-	0.0	-	0.0	-
123.0	55.0	-	0.0	-	3.3	-	-	-	0.0	-	41.5	-
123.5	35.6	-	0.0	-	1.2	-	-	-	0.0	-	9.0	-
127.0	34.0	-	0.0	-	0.0	-	-	-	0.0	-	12.6	-
127.0	40.0	-	0.0	-	0.0	-	-	-	0.0	-	0.0	-
127.0	55.0	-	0.0	-	5.9	-	-	-	8.2	-	0.0	-
127.0	65.0	-	0.0	-	0.0	-	-	-	8.6	-	-	-
127.0	70.0	-	2.9	-	0.0	-	-	-	8.5	-	21.1	0.0
130.0	25.5	-	0.6	-	0.0	-	-	-	0.0	-	5.0	2.4
130.0	26.0	-	1.0	-	2.4	-	-	-	0.0	-	10.1	0.0
130.0	28.0	-	4.3	-	5.4	-	-	-	0.0	-	0.0	15.0
130.0	30.0	-	0.0	-	0.0	-	-	-	0.0	-	0.0	0.0
130.0	35.0	-	3.3	-	0.0	-	-	-	0.0	-	0.0	-
130.0	40.0	-	0.0	-	15.8	-	-	-	0.0	-	0.0	-
130.0	45.0	-	11.2	-	0.0	-	-	-	0.0	-	0.0	-
130.0	50.0	-	12.9	-	0.0	-	-	-	2.7	-	0.0	3.1
130.0	55.0	-	0.0	-	0.0	-	-	-	0.0	-	10.0	-
130.0	70.0	-	2.9	-	0.0	-	-	-	8.2	-	0.0	-
130.0	80.0	-	0.0	-	2.7	-	-	-	5.4	-	0.0	-
130.0	90.0	-	2.8	-	-	-	-	-	0.0	-	10.0	-
130.0	120.0	-	2.7	-	5.3	-	-	-	0.0	-	10.0	-
133.0	23.0	-	0.0	-	0.0	-	-	-	0.0	-	3.0	-
133.0	35.0	-	0.0	-	0.0	-	-	-	8.7	-	3.0	-
133.0	40.0	-	0.0	-	2.9	-	-	-	0.0	-	0.0	-
133.0	45.0	-	2.9	-	3.0	-	-	-	0.0	-	0.0	-
133.0	50.0	-	25.5	-	3.0	-	-	-	0.0	-	0.0	-
133.0	65.0	-	0.0	-	6.2	-	-	-	5.3	-	-	-
133.0	70.0	-	0.0	-	3.0	-	-	-	2.7	-	0.0	2.2
137.0	20.0	-	0.5	-	0.0	-	-	-	-	-	23.5	0.0
137.0	21.0	-	0.0	-	0.0	-	-	-	-	-	47.5	2.8
137.0	22.0	-	3.9	-	0.0	-	-	-	-	-	30.4	3.0
137.0	23.0	-	1.9	-	0.0	-	-	-	-	-	0.0	3.1
137.0	30.0	-	0.0	-	0.0	-	-	-	-	-	0.0	2.2
137.0	35.0	-	15.5	-	3.0	-	-	-	-	-	0.0	-
137.0	45.0	-	0.0	-	3.0	-	-	-	-	-	2.6	0.0
137.0	50.0	-	3.3	-	0.0	-	-	-	-	-	5.5	-
137.0	55.0	-	0.0	-	-	-	-	-	-	-	0.0	-
137.0	60.0	-	-	-	-	-	-	-	-	-	0.0	-
137.0	70.0	-	6.0	-	-	-	-	-	-	-	0.0	-
137.0	80.0	-	0.0	-	-	-	-	-	-	-	10.0	-
140.0	28.8	-	-	-	-	-	-	-	-	-	-	-

TABLE 4. (cont.)

## Disintegrated fish larva (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
140.0	29.0	-	2.3	-	-	-	-	-	0.0	-	-	-
140.0	30.0	-	0.0	-	-	-	-	-	2.8	-	-	-
140.0	35.0	-	12.1	-	-	-	-	-	-	5.3	-	-
140.0	45.0	-	3.1	-	-	-	-	-	-	0.0	-	-
140.0	50.0	-	0.0	-	-	-	-	-	-	2.3	-	-

## Unidentified fish larva

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
60.0	51.0	11.3	-	-	-	-	-	0.0	-	-	-	-
60.0	52.0	6.8	-	-	-	-	-	0.0	-	-	-	-
60.0	55.0	3.1	-	-	-	-	-	0.0	-	-	-	-
60.0	200.0	3.1	-	-	-	0.0	-	-	-	-	-	-
63.0	51.0	2.6	-	-	-	0.0	-	-	-	-	-	-
63.0	52.0	0.0	-	-	-	0.0	-	-	-	-	-	-
63.0	52.0	0.0	-	-	-	0.0	-	-	-	-	-	-
63.0	55.0	12.7	-	-	-	3.7	-	-	-	-	-	-
67.0	47.2	-	-	-	-	-	-	-	-	-	-	-
67.0	48.0	0.0	-	-	-	1.3	-	-	-	-	-	-
67.0	49.0	6.5	-	-	-	0.0	-	-	-	-	-	-
67.0	50.0	0.0	-	-	-	5.4	-	-	-	-	-	-
67.0	50.0	0.0	-	-	-	2.2	-	-	-	-	-	-
70.0	50.0	0.0	-	-	-	2.6	-	-	-	-	-	-
70.0	52.0	0.0	-	-	-	-	-	-	-	-	-	-
70.0	55.0	8.7	-	-	-	-	-	-	-	-	-	-
70.0	60.0	0.0	-	-	-	2.9	-	-	-	-	-	-
70.0	200.0	3.1	-	-	-	-	-	-	-	-	-	-
73.0	53.0	0.0	-	-	-	3.0	-	-	-	-	-	-
77.0	47.6	4.2	-	-	-	0.0	-	-	-	-	-	-
77.0	51.0	3.0	-	-	-	0.0	-	-	-	-	-	-
77.0	60.0	-	-	-	-	3.1	-	-	-	-	-	-
80.0	50.2	0.0	-	-	-	0.8	-	-	-	-	-	-
80.0	52.0	0.0	-	-	-	3.3	-	-	-	-	-	-
80.0	70.0	0.0	-	-	-	3.4	-	-	-	-	-	-
80.0	160.0	8.8	-	-	-	-	-	-	-	-	-	-
80.0	200.0	2.8	-	-	-	-	-	-	-	-	-	-
82.0	47.0	0.0	-	-	-	0.0	-	-	-	-	-	-
83.0	39.0	0.0	-	-	-	-	-	-	-	-	-	-
83.0	40.0	0.0	-	-	-	0.0	-	-	-	-	-	-
83.0	43.0	0.0	-	-	-	0.0	-	-	-	-	-	-
83.0	51.0	0.0	-	-	-	7.8	-	-	-	-	-	-
83.0	55.0	0.0	-	-	-	0.0	-	-	-	-	-	-
83.0	60.0	0.0	-	-	-	0.0	-	-	-	-	-	-
83.0	65.0	0.0	-	-	-	0.0	-	-	-	-	-	-
83.0	70.0	6.4	-	-	-	0.0	-	-	-	-	-	-
87.0	32.5	1.6	-	-	-	0.0	-	-	-	-	-	-
87.0	33.0	4.0	-	-	-	2.6	-	-	-	-	-	-

TABLE 4. (cont.)

## Unidentified fish larva (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
87.0	34.0	2.7	-	-	0.0	-	10.6	-	-	0.0	-	-
87.0	35.0	0.0	-	-	3.4	-	3.1	-	-	3.0	-	-
87.0	50.0	0.0	-	-	0.0	-	2.7	-	-	0.0	-	-
87.0	55.0	3.0	-	-	0.0	-	3.3	-	-	0.0	-	-
87.0	90.0	0.0	-	-	2.8	-	3.1	-	-	0.0	-	-
87.0	100.0	-	-	-	3.1	-	-	-	-	-	-	-
90.0	27.5	2.3	-	-	2.6	-	-	-	-	0.0	-	-
90.0	28.0	0.0	-	-	3.1	-	-	-	-	0.0	-	-
90.0	32.0	0.0	-	-	3.0	-	-	-	-	0.0	-	-
90.0	37.0	0.0	-	-	0.0	-	-	-	-	0.0	-	-
90.0	45.0	0.0	-	-	0.0	-	-	-	-	0.0	-	-
90.0	90.0	0.0	-	-	3.2	-	-	-	-	0.0	-	-
90.0	180.0	0.0	-	-	-	-	-	-	-	5.6	-	6.3
90.0	200.0	0.0	-	-	-	-	-	-	-	11.0	-	2.2
93.0	26.5	0.0	-	-	-	-	-	-	-	23.0	-	11.8
93.0	27.0	0.0	-	-	-	-	-	-	-	9.6	-	2.5
93.0	28.0	0.0	-	-	6.6	-	-	-	-	9.2	-	0.0
93.0	30.0	0.0	-	-	0.0	-	-	-	-	3.2	-	0.0
93.0	40.0	0.0	-	-	0.0	-	-	-	-	0.0	-	5.5
93.0	70.0	0.0	-	-	0.0	-	-	-	-	1.0	-	0.0
96.8	28.7	3.5	-	-	0.7	-	-	-	-	0.0	-	0.0
97.0	29.0	-	-	-	0.0	-	-	-	-	4.8	-	0.0
97.0	30.0	-	-	-	20.6	-	-	-	-	10.1	-	0.0
97.0	32.0	-	-	-	2.2	-	-	-	-	21.7	-	0.0
97.0	40.0	-	-	-	3.1	-	-	-	-	0.0	-	3.0
97.0	50.0	-	-	-	0.0	-	-	-	-	18.9	-	0.0
97.0	55.0	-	-	-	0.0	-	-	-	-	0.0	-	5.7
97.0	80.0	-	-	-	3.8	-	-	-	-	2.9	-	5.8
100.0	29.0	-	-	-	0.0	-	-	-	-	83.8	-	0.0
100.0	55.0	-	-	-	3.0	-	-	-	-	0.0	-	0.0
100.0	60.0	-	-	-	0.0	-	-	-	-	0.0	-	0.0
100.0	65.0	-	-	-	3.0	-	-	-	-	0.0	-	0.0
100.0	85.0	-	-	-	-	-	-	-	-	3.5	-	0.0
100.0	90.0	-	-	-	3.0	-	-	-	-	6.6	-	1.0
103.0	28.5	-	-	-	0.0	-	-	-	-	0.0	-	8.6
103.0	29.0	-	-	-	0.0	-	-	-	-	0.0	-	0.0
103.0	30.0	-	-	-	2.5	-	-	-	-	0.0	-	0.0
103.0	35.0	-	-	-	0.0	-	-	-	-	9.3	-	0.0
103.0	45.0	-	-	-	0.0	-	-	-	-	6.8	-	0.0
103.0	90.0	-	-	-	0.0	-	-	-	-	3.3	-	5.6
105.0	140.0	-	-	-	-	-	-	-	-	0.0	-	1.1
107.0	130.0	-	-	-	-	-	-	-	-	3.1	-	0.6
107.0	31.0	-	-	-	-	-	-	-	-	0.0	-	4.8
107.0	32.0	-	-	-	-	-	-	-	-	0.0	-	0.0
107.0	65.0	-	-	-	-	-	-	-	-	0.0	-	3.3
107.0	70.0	-	-	-	-	-	-	-	-	0.0	-	0.0

TABLE 4. (cont.)

## Unidentified fish larva (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
107.0	80.0	-	8.7	-	0.0	-	0.0	-	-	-	-	-
107.0	90.0	-	11.8	-	0.0	-	0.0	-	-	-	-	-
110.0	32.0	-	0.7	-	0.0	-	-	-	0.0	0.0	0.0	-
110.0	35.0	-	0.0	-	0.0	-	-	-	2.8	0.0	-	-
110.0	55.0	-	0.0	-	0.0	-	-	-	0.0	3.1	-	-
110.0	65.0	-	2.7	-	0.0	-	-	-	0.0	0.0	-	-
110.0	140.0	-	0.0	-	-	-	8.3	-	-	-	-	-
110.0	160.0	-	0.0	-	-	-	2.9	-	-	5.6	-	-
113.0	28.5	-	0.0	-	2.2	0.0	-	-	9.4	3.6	-	-
113.0	29.0	-	0.0	-	0.0	0.0	-	-	-	0.0	-	-
113.0	30.0	-	0.0	-	0.0	0.0	-	-	-	0.0	-	-
113.0	50.0	-	0.0	-	0.0	0.0	-	-	-	0.0	-	-
113.0	55.0	-	0.0	-	0.0	0.0	-	-	-	0.0	-	-
113.0	65.0	-	0.0	-	0.0	0.0	-	-	-	0.0	-	-
115.0	35.0	-	0.0	-	1.0	0.0	-	-	2.6	0.3	-	-
117.0	24.8	-	0.0	-	0.0	0.0	-	-	1.9	2.3	-	-
117.0	25.0	-	0.0	-	5.6	0.0	-	-	11.8	0.0	-	-
117.0	26.0	-	0.0	-	0.0	0.0	-	-	10.7	5.3	-	-
117.0	30.0	-	0.0	-	0.0	0.0	-	-	8.8	7.8	-	-
117.0	35.0	-	0.0	-	0.0	0.0	-	-	0.0	5.8	-	-
117.0	40.0	-	0.0	-	3.0	0.0	-	-	0.0	0.0	-	-
117.0	50.0	-	0.0	-	0.0	0.0	-	-	6.6	0.0	-	-
117.0	55.0	-	0.0	-	0.0	0.0	-	-	6.6	0.0	-	-
117.0	60.0	-	0.0	-	3.2	0.0	-	-	0.0	0.0	-	-
117.0	90.0	-	0.0	-	2.8	0.0	-	-	2.8	17.4	-	-
118.0	39.0	-	3.1	-	9.3	-	-	-	37.1	6.3	-	-
119.0	33.0	-	2.7	-	0.0	-	-	-	55.9	16.9	-	-
120.0	22.5	-	0.0	-	5.6	-	-	-	11.6	23.8	-	-
120.0	23.0	-	0.0	-	0.0	-	-	-	12.0	28.0	-	-
120.0	24.0	-	0.0	-	2.7	-	-	-	27.2	15.1	-	-
120.0	25.0	-	0.0	-	22.0	0.0	-	-	26.0	0.7	-	-
120.0	30.0	-	0.0	-	14.5	27.4	-	-	25.8	11.4	-	-
120.0	35.0	-	0.0	-	0.0	0.0	-	-	-	3.0	0.0	-
120.0	40.0	-	0.0	-	0.0	0.0	-	-	-	0.0	3.1	-
120.0	45.0	-	0.0	-	0.0	0.0	-	-	-	0.0	0.0	-
120.0	50.0	-	0.0	-	0.0	0.0	-	-	-	0.0	0.0	-
120.0	60.0	-	0.0	-	0.0	0.0	-	-	-	0.0	12.0	-
120.0	80.0	-	0.0	-	0.0	0.0	-	-	-	0.0	0.0	-
120.0	100.0	-	0.0	-	3.0	-	-	-	3.4	3.2	-	-
120.0	120.0	-	0.0	-	8.7	-	-	-	-	0.0	0.0	-
123.0	36.0	-	0.0	-	0.0	0.0	-	-	2.4	0.0	-	-
123.0	37.0	-	0.0	-	0.0	0.0	-	-	0.0	0.0	-	-
123.0	42.0	-	0.0	-	4.9	-	-	-	0.0	0.0	-	-
123.0	45.0	-	0.0	-	0.0	0.0	-	-	10.0	0.0	-	-
123.0	50.0	-	0.0	-	0.0	0.0	-	-	5.7	0.0	-	-
123.0	55.0	-	0.0	-	0.0	0.0	-	-	0.0	0.0	-	-

TABLE 4. (cont.)

## Unidentified fish larva (cont.)

STATION	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.
123.0	65.0	-	0.0	-	0.0	-	-	2.8	-	-	-	-
123.0	80.0	-	0.0	-	18.5	-	-	0.0	-	-	-	-
123.5	35.6	-	2.2	-	4.9	-	-	1.3	-	-	15.7	-
127.0	32.6	0.0	-	0.0	-	-	-	11.0	-	-	0.0	-
127.0	33.0	0.0	-	0.0	-	-	-	3.0	-	-	2.6	-
127.0	34.0	-	23.7	2.4	-	-	-	2.4	-	-	0.0	-
127.0	40.0	-	7.1	5.9	-	-	-	2.8	24.6	0.0	-	-
127.0	45.0	0.0	-	0.0	-	-	-	12.9	-	-	0.0	-
127.0	50.0	0.0	-	3.0	-	-	-	0.0	-	-	0.0	-
127.0	60.0	-	2.7	0.0	-	-	-	0.0	-	-	46.1	-
130.0	25.5	0.6	-	1.8	-	-	-	-	208.2	-	83.5	-
130.0	26.0	0.0	-	2.4	-	-	-	-	25.7	-	-	-
130.0	28.0	0.0	-	0.0	-	-	-	-	7.4	2.4	-	-
130.0	30.0	0.0	-	0.0	-	-	-	-	4.0	0.0	-	-
130.0	35.0	0.0	-	0.0	-	-	-	-	10.2	6.1	-	-
130.0	40.0	18.4	-	0.0	-	-	-	-	0.0	0.0	-	-
130.0	50.0	15.8	-	0.0	-	-	-	-	5.5	0.0	-	-
130.0	55.0	2.9	-	0.0	-	-	-	-	0.0	0.0	-	-
130.0	60.0	0.0	-	0.0	-	-	-	-	2.8	0.0	-	-
130.0	80.0	2.8	-	-	-	-	-	-	0.0	-	-	-
130.0	90.0	0.0	-	1.7	-	-	-	-	5.4	-	3.6	-
133.0	19.5	-	0.0	-	-	-	-	-	-	-	-	-
133.0	21.0	0.0	-	2.2	-	-	-	-	4.0	-	52.7	-
133.0	23.0	0.0	-	0.0	-	-	-	-	2.5	-	20.0	-
133.0	30.0	0.0	-	0.0	-	-	-	-	14.8	-	28.6	-
133.0	45.0	2.9	-	0.0	-	-	-	-	0.0	-	0.0	-
133.0	50.0	0.0	-	0.0	-	-	-	-	0.0	-	3.0	-
133.0	55.0	2.7	-	0.0	-	-	-	-	0.0	-	-	-
133.0	60.0	0.0	-	3.0	-	-	-	-	-	-	201.2	-
137.0	20.0	0.0	-	0.0	-	-	-	-	-	-	224.7	-
137.0	21.0	0.0	-	0.0	-	-	-	-	-	-	15.2	-
137.0	22.0	0.0	-	0.0	-	-	-	-	-	-	87.7	-
137.0	23.0	0.0	-	2.6	-	-	-	-	-	-	51.0	-
137.0	35.0	28.0	-	0.0	-	-	-	-	-	-	16.6	-
137.0	40.0	29.2	-	0.0	-	-	-	-	-	-	0.0	-
137.0	50.0	0.0	-	0.0	-	-	-	-	-	-	0.0	-
137.0	55.0	0.0	-	0.0	-	-	-	-	-	-	5.2	-
137.0	60.0	0.0	-	0.0	-	-	-	-	-	-	0.0	-
137.0	70.0	0.0	-	6.3	-	-	-	-	-	-	-	-
140.0	28.8	0.0	-	-	-	-	-	-	-	-	5.8	-
140.0	29.0	0.0	-	-	-	-	-	-	-	-	13.7	-
140.0	35.0	6.1	-	-	-	-	-	-	-	-	15.3	-
140.0	40.0	17.7	-	-	-	-	-	-	-	-	0.0	-
140.0	45.0	6.1	-	-	-	-	-	-	-	-	0.0	-

TABLE 5. Summary of pooled occurrences of all larval fish taxa taken on CalCOFI surveys from 1961 to 1969. Taxa are listed in the same order as Table 4.

NAME	1961	1962	1963	1964	1965	1966	1967	1968	1969
<i>Anguilliformes</i>									
<i>Etrumeus acuminatus</i>	7	8	20	8	24	17	5	3	13
<i>Opisthonema</i> spp.	4	7	36	37	35	26	7	1	9
<i>Sardinops sagax</i>	-	-	-	-	2	3	-	-	-
<i>Engraulis mordax</i>	53	58	99	88	104	143	31	10	79
<i>Argentina sialis</i>	408	454	567	707	618	987	150	188	880
<i>Microstoma microstoma</i>	18	49	33	37	49	93	21	18	98
<i>Nansenia candida</i>	12	19	11	31	17	48	9	19	73
<i>Nansenia crassa</i>	9	13	5	7	9	39	6	12	32
<i>BathyLAGUS</i> spp.	29	15	30	33	22	48	8	5	40
<i>BathyLAGUS milleri</i>	18	1	54	31	7	18	6	35	215
<i>BathyLAGUS ochotensis</i>	-	-	2	3	1	1	-	1	33
<i>BathyLAGUS pacificus</i>	57	66	98	196	127	260	28	106	359
<i>BathyLAGUS wesethi</i>	5	7	8	38	3	26	-	15	80
<i>Leuroglossus stilbius</i>	149	168	160	235	220	461	99	90	328
<i>Dolichopteryx</i> spp.	202	225	236	360	300	449	43	116	498
<i>Macropinna microstoma</i>	-	-	-	-	-	-	-	-	-
<i>Osmariidae</i>	1	-	-	-	-	-	-	-	1
<i>Stomiiformes</i>	-	-	-	-	-	-	-	-	-
<i>Gonostomatidae</i>	12	4	3	6	1	6	9	1	4
<i>Cyclothone</i> spp.	2	5	12	8	18	8	-	4	126
<i>Diplophos taenia</i>	214	277	241	247	265	593	80	65	346
<i>Ichthyococcus</i> spp.	5	5	7	-	3	11	1	1	7
<i>Vinciguerria lucetiae</i>	4	11	11	13	7	35	5	2	34
<i>Vinciguerria poweriae</i>	342	371	383	369	436	828	121	82	479
<i>Woodsia nonsuchae</i>	3	7	3	4	3	6	-	-	1
<i>Sternopychidae</i>	-	-	1	-	-	-	-	-	-
<i>Astronesthidae</i>	54	71	45	79	59	250	28	48	469
<i>Chauliodus macouni</i>	-	2	-	-	-	-	-	-	-
<i>Idiacanthus antrostomus</i>	28	28	31	68	57	171	9	46	189
<i>Aristostomias scintillans</i>	48	43	26	32	33	72	15	22	114
<i>Bathophilus</i> spp.	9	10	9	6	9	12	2	1	2
<i>Eustomias</i> spp.	5	10	4	3	4	5	2	-	-
<i>Photonectes</i> spp.	1	1	-	1	1	-	-	-	-
<i>Tactostoma macropus</i>	7	4	-	4	2	6	4	-	4
<i>Stomias atriventris</i>	58	76	98	81	100	326	24	46	214
<i>Evermannellidae</i>	1	3	1	1	1	-	-	-	-
<i>Paralepididae</i>	-	3	5	10	3	-	-	3	6
<i>Lestidiops ringens</i>	50	80	58	63	67	232	36	52	231
<i>Notolepis risso</i>	9	12	9	7	9	12	2	8	18
<i>Paralepis atlantica</i>	-	-	-	-	1	-	-	1	-
<i>Stemonosudis macrura</i>	4	6	-	2	6	5	-	-	-
<i>Sudis atrox</i>	2	4	-	2	4	-	1	-	-
<i>Aulopus</i> spp.	-	-	-	-	16	19	21	6	36
<i>Scopelelosaurus</i> spp.	16	10	8	50	21	33	114	13	93
<i>Scopelarchidae</i>	67	60	-	-	-	-	-	-	-

TABLE 5. (cont.)

NAME	1961	1962	1963	1964	1965	1966	1967	1968	1969
<i>Myctophidae</i>									
<i>Ceratoscopelus townsendi</i>	165	151	179	220	222	346	33	79	329
<i>Diaphus</i> spp.	149	157	128	146	156	302	37	23	153
<i>Lampanyctus urophoas</i>	77	56	46	101	80	187	46	34	110
<i>Lampanyctus</i> spp.	53	45	50	25	32	62	10	1	23
<i>Lampanyctus regalis</i>	148	139	199	155	183	401	67	65	550
<i>Lampanyctus ritteri</i>	13	12	2	20	9	46	12	11	19
<i>Notolynchus valdiviae</i>	154	204	120	189	234	523	43	72	155
<i>Notoscopelus resplendens</i>	29	13	22	16	21	22	7	1	10
<i>Parvilux ingens</i>	59	41	50	39	44	54	11	3	29
<i>Stenobrachius leucopsarus</i>	-	-	-	-	-	-	-	-	1
<i>Triphoturus mexicanus</i>	177	179	186	342	263	420	31	127	390
<i>Triphoturus nigrescens</i>	407	422	451	448	494	990	142	92	556
<i>Benthosema pterota</i>	4	-	-	-	1	-	-	-	-
<i>Centrobranchus</i> spp.	-	-	-	-	3	-	-	-	-
<i>Diogenichthys</i> spp.	2	10	-	2	2	-	1	-	2
<i>Diogenichthys atlanticus</i>	54	62	88	61	111	165	16	13	79
<i>Diogenichthys laternatus</i>	102	155	92	111	116	171	38	46	210
<i>Electrona rissoii</i>	94	127	161	163	249	361	63	32	210
<i>Gonichthys tenuiculus</i>	3	5	-	3	2	3	-	-	7
<i>Gonichthys</i> spp.	20	24	29	46	81	146	16	12	48
<i>Hygophum</i> spp.	4	3	29	6	11	4	-	-	13
<i>Hygophum atratum</i>	27	38	41	44	103	178	21	6	81
<i>Hygophum reinhardtii</i>	39	58	27	20	27	79	7	-	10
<i>Loweina rara</i>	8	4	5	4	8	6	1	-	11
<i>Myctophum nitidulum</i>	46	42	31	32	19	58	11	8	59
<i>Protomyctophum crockeri</i>	247	252	225	292	261	671	109	139	717
<i>Protomyctophum thompsoni</i>	-	-	-	-	-	-	-	-	9
<i>Symbolophorus californiensis</i>	82	140	78	116	111	291	38	61	157
<i>Tarletonbeania crenularis</i>	160	115	111	140	132	208	10	73	277
<i>Synodus</i> spp.	19	23	41	35	42	121	23	-	54
<i>Bregmaceros</i> spp.	-	-	-	-	-	-	-	-	-
<i>Microgadus proximus</i>	-	-	-	3	-	2	-	-	-
<i>Merluccius productus</i>	152	228	229	290	290	398	25	95	361
<i>Physiculus</i> spp.	-	1	1	1	3	2	1	-	2
<i>Macrouridae</i>	4	6	6	5	5	5	2	3	14
<i>Ophidiiformes</i>	16	16	35	49	37	69	10	16	45
<i>Brosomophycis marginata</i>	-	2	3	3	7	17	5	8	16
<i>Carapidae</i>	-	1	-	1	-	-	-	-	-
<i>Chilara taylori</i>	12	31	15	11	29	55	15	-	28
<i>Ophidion scriptae</i>	2	10	61	19	40	67	-	-	34
<i>Porichthys</i> spp.	1	-	1	1	-	1	1	-	2
<i>Ceratioidae</i>	15	26	17	7	18	43	-	-	30
<i>Gobiesocidae</i>	3	-	5	8	9	12	-	-	1
<i>Exocoetidae</i>	2	-	1	3	2	10	-	-	5
<i>Hemiramphidae</i>	-	-	-	2	1	-	-	-	-
<i>Cololabis saira</i>	11	6	13	22	9	31	11	3	32
<i>Atherinidae</i>	-	9	23	8	11	2	2	2	5
<i>Trachipteridae</i>	27	-	20	-	19	-	6	6	-
<i>Eutaeniophoridae</i>	-	-	-	-	-	-	-	-	5

TABLE 5. (cont.)

NAME	1961	1962	1963	1964	1965	1966	1967	1968	1969
<i>Melamphaes</i> spp.									
<i>Poromitra</i> spp.									
<i>Scopeloberyx robustus</i>	117	106	134	114	151	340	68	84	333
<i>Scopelogadus bispinosus</i>	13	18	28	28	32	51	6	14	27
<i>HacroAMPHOSUS gracilis</i>	4	2	2	—	7	—	—	—	2
<i>Syngnathus</i> spp.	18	34	10	31	13	60	4	5	17
<i>Agonidae</i>	3	6	6	3	7	6	7	—	11
<i>Anoplopoma fimbria</i>	3	6	16	24	22	20	6	3	10
<i>Cottidae</i>	—	—	—	1	—	5	5	4	9
<i>Scorpaenichthys marmoratus</i>	11	21	33	45	37	43	5	12	—
<i>Cyclopteridae</i>	3	3	7	13	20	15	5	24	
<i>Hexagrammidae</i>	8	2	12	14	16	14	4	4	17
<i>Ophiodon elongatus</i>	—	—	—	2	1	1	1	1	6
<i>Oxyplectus pictus</i>	6	3	7	27	13	7	5	20	
<i>Zaniolepis</i> spp.	2	9	12	11	7	26	7	3	19
<i>Scorpaenidae</i>	—	1	2	—	—	1	1	—	
<i>Scorpaena</i> spp.	11	11	17	16	25	62	8	3	12
<i>Sebastodes</i> spp.	311	273	289	492	387	698	81	207	705
<i>Sebastolobus</i> spp.	8	2	17	20	20	87	4	14	47
<i>Prionotus</i> spp.	10	9	40	15	30	25	—	—	1.9
<i>Acanthuridae</i>	—	—	—	—	—	—	—	—	
<i>Blenioidei</i>	1	1	14	6	4	—	3	—	4
<i>Hypsoblennius</i> spp.	11	14	68	69	73	77	19	6	61
<i>Clinidae</i>	12	21	31	44	64	51	9	10	51
<i>Gobiidae</i>	31	41	87	80	104	198	36	19	138
<i>Icosteus aenigmaticus</i>	1	1	1	1	—	3	—	1	
<i>Labridae</i>	—	2	9	—	7	—	2	3	
<i>Halichoeres</i> spp.	12	12	40	18	36	50	4	1	28
<i>Oxyjulis californica</i>	23	22	34	15	31	97	23	15	58
<i>Semicossyphus pulcher</i>	6	10	21	7	27	28	4	—	8
<i>Pomacentridae</i>	—	—	10	4	8	5	—	—	
<i>Chromis punctipinnis</i>	3	21	42	13	39	105	5	1	54
<i>Hypsypops rubicundus</i>	—	—	—	—	8	1	—	—	
<i>Hugil</i> spp.	—	—	—	—	1	1	1	1	4
<i>Apogonidae</i>	—	—	—	—	5	4	3	1	12
<i>Bowella brodiei</i>	16	7	—	—	4	—	—	—	3
<i>Brama</i> spp.	21	17	17	7	—	21	1	—	—
<i>Carangidae</i>	—	1	20	14	25	13	2	—	9
<i>Seriola lalandii</i>	5	12	15	7	14	30	5	4	4
<i>Trachurus symmetricus</i>	144	—	—	206	214	503	76	85	248
<i>Caristius macropus</i>	—	—	—	—	1	1	—	—	—
<i>Coryphaena hippurus</i>	—	7	2	1	10	5	1	—	1
<i>Chaetodipterus zonatus</i>	—	—	—	—	—	—	14	12	4
<i>Gerreidae</i>	—	—	—	—	—	—	—	—	4
<i>Haemulidae</i>	—	2	15	10	11	17	—	—	7
<i>Girella nigricans</i>	—	—	—	—	16	3	3	3	7
<i>Medialuna californiensis</i>	5	4	11	13	4	5	22	6	12
<i>Caulolatilus princeps</i>	4	4	3	2	2	5	7	3	2
<i>Mullidae</i>	—	—	—	—	—	—	—	—	—

TABLE 5. (cont.)

NAME	1961	1962	1963	1964	1965	1966	1967	1968	1969
<i>Sciaenidae</i>	28	42	85	135	147	32	38	38	195
<i>Serranidae</i>	10	6	68	38	59	23	2	72	-
<i>Sparidae</i>	-	-	1	-	-	-	-	-	-
<i>Polynemidae</i>	-	15	6	5	8	1	-	-	1
<i>Gempylidae</i>	7	-	3	2	4	7	8	2	2
<i>Scombridae</i>	3	-	-	-	-	-	-	-	-
<i>Auxis</i> spp.	-	15	6	5	8	4	-	-	-
<i>Sarda chiliensis</i>	7	3	10	8	9	29	1	-	30
<i>Scomber japonicus</i>	26	32	57	39	34	68	14	-	24
<i>Scomberomorus</i> spp.	1	-	1	1	5	3	-	-	-
<i>Trichiuridae</i>	10	23	27	17	27	74	10	-	23
<i>Sphyraena argentea</i>	6	6	22	10	25	31	7	4	15
<i>Icichthys lockingtoni</i>	38	39	52	78	53	131	18	48	202
<i>Nomeidae</i>	-	-	1	1	1	2	-	-	1
<i>Peprius simillimus</i>	2	19	19	18	45	52	22	11	45
<i>Tetragonurus cuvieri</i>	45	76	98	46	31	74	36	5	48
<i>Chiasmodontidae</i>	25	22	39	13	40	60	6	10	41
<i>Pleuronectiformes</i>	2	-	13	7	4	-	1	1	7
<i>Bothus</i> spp.	-	-	2	-	-	-	-	-	-
<i>Citharichthys</i> spp.	186	221	281	243	342	590	108	101	611
<i>Citharichthys stigmaeus</i>	50	97	65	73	65	171	19	42	269
<i>Hippoglossina stomata</i>	24	15	44	42	44	83	12	5	52
<i>Paralichthys californicus</i>	21	37	57	96	107	81	13	13	60
<i>Syacium ovale</i>	-	-	3	-	1	3	-	-	-
<i>Xystreurus liolepis</i>	1	9	15	18	8	30	4	-	22
<i>Glyptocephalus zachirus</i>	2	-	9	18	4	36	-	14	15
<i>Hypsopsetta guttulata</i>	1	-	4	5	10	3	-	-	6
<i>Lepidopsetta bilineata</i>	1	1	-	1	2	3	2	2	1
<i>Lyopsetta exilis</i>	32	31	33	46	33	72	4	20	65
<i>Microstomus pacificus</i>	2	-	11	13	16	52	13	17	56
<i>Parophrys vetulus</i>	14	32	41	41	41	80	6	21	80
<i>Platichthys stellatus</i>	-	-	-	-	-	3	-	-	-
<i>Pleuronichthys</i> spp.	4	3	10	12	1	-	10	3	15
<i>Pleuronichthys coenosus</i>	2	2	6	9	5	11	1	-	11
<i>Pleuronichthys decurrens</i>	1	4	-	1	4	11	-	2	1
<i>Pleuronichthys ritteri</i>	5	3	12	12	9	8	2	1	7
<i>Pleuronichthys verticalis</i>	10	47	56	74	88	81	24	18	66
<i>Psettichthys melanostictus</i>	1	1	5	12	9	10	-	4	14
<i>Syphurus</i> spp.	18	41	73	48	75	138	10	-	71
<i>Soleidae</i>	-	-	-	-	-	-	-	-	-
<i>Tetraodontidae</i>	-	-	-	-	-	-	-	-	-
Disintegrated fish larva	184	223	274	311	542	84	74	458	458
Unidentified fish larva	147	147	256	217	485	60	72	422	422

TABLE 6. List of stations which were occupied twice in one month during 1965.

Station	Month
---------	-------

113.0	30.0	2
123.0	42.0	2
127.0	40.0	2



## INDEX

This index lists taxa included in Table 4 with their page numbers.

	Page
<b>Anguilliformes . . . . .</b>	56
<b>Clupeiformes</b>	
<b>Clupeidae</b>	
<i>Etrumeus acuminatus</i> . . . . .	56
<i>Opisthonema</i> spp. . . . .	57
<i>Sardinops sagax</i> . . . . .	57
<b>Engraulidae</b>	
<i>Engraulis mordax</i> . . . . .	59
<b>Salmoniformes</b>	
<b>Argentinidae</b>	
<i>Argentina sialis</i> . . . . .	64
<i>Microstoma microstoma</i> . . . . .	65
<i>Nansenia candida</i> . . . . .	65
<i>Nansenia crassa</i> . . . . .	65
<b>Bathylagidae</b>	
<i>Bathylagus</i> spp. . . . .	66
<i>Bathylagus milleri</i> . . . . .	66
<i>Bathylagus ochotensis</i> . . . . .	66
<i>Bathylagus pacificus</i> . . . . .	69
<i>Bathylagus wesethi</i> . . . . .	69
<i>Leuroglossus stilbius</i> . . . . .	72
<b>Stomiiformes . . . . .</b>	76
<b>Gonostomatidae</b>	
<i>Cyclothona</i> spp. . . . .	77
<i>Diplophos taenia</i> . . . . .	80
<i>Ichthyococcus</i> spp. . . . .	80
<i>Vinciguerria lucetia</i> . . . . .	81
<i>Vinciguerria poweriae</i> . . . . .	85
<b>Sternopychidae</b>	
<b>Stomatoidea</b>	
<b>Chauliodontidae</b>	
<i>Chauliodus macouni</i> . . . . .	87
<b>Idiacanthidae</b>	
<i>Idiacanthus antrostomus</i> . . . . .	88
<b>Malacosteidae</b>	
<i>Aristostomias scintillans</i> . . . . .	89
<b>Melanostomiidae</b>	
<i>Bathophilus</i> spp. . . . .	89
<i>Eustomias</i> spp. . . . .	89
<i>Photonectes</i> spp. . . . .	89
<i>Tactostoma macropus</i> . . . . .	89
<b>Stomiidae</b>	
<i>Stomias atriventer</i> . . . . .	90
<b>Myctophiformes</b>	
<b>Alepisauroidei</b>	
<b>Evermannellidae</b> . . . . .	91

	Page
Paralepididae .....	92
<i>Lestidiops ringens</i> .....	92
<i>Notolepis risso</i> .....	93
<i>Paralepis atlantica</i> .....	93
<i>Stemonosudis macrura</i> .....	94
<i>Sudis atrox</i> .....	94
Chlorophthalmoidei	
Notosudidae	
<i>Scopelosaurus</i> spp. ....	94
Scopelarchidae .....	94
Myctophoidei	
Myctophidae .....	95
Lampanyctinae	
<i>Ceratoscopelus townsendi</i> .....	99
<i>Diaphus</i> spp. ....	101
<i>Lampadena urophaos</i> .....	103
<i>Lampanyctus</i> spp. ....	104
<i>Lampanyctus regalis</i> .....	107
<i>Lampanyctus ritteri</i> .....	107
<i>Notolychnus valdiviae</i> .....	110
<i>Notoscopelus resplendens</i> .....	111
<i>Stenobrachius leucopsarus</i> .....	112
<i>Triphoturus mexicanus</i> .....	115
<i>Triphoturus nigrescens</i> .....	121
Myctophinae	
<i>Centrobranchus</i> spp. ....	121
<i>Diogenichthys</i> spp. ....	121
<i>Diogenichthys atlanticus</i> .....	121
<i>Diogenichthys laternatus</i> .....	123
<i>Electrona rissoii</i> .....	126
<i>Gonichthys tenuiculus</i> .....	126
<i>Hygophum</i> spp. ....	127
<i>Hygophum atratum</i> .....	127
<i>Hygophum reinhardtii</i> .....	129
<i>Loweina rara</i> .....	129
<i>Myctophum nitidulum</i> .....	130
<i>Protomyctophum crockeri</i> .....	130
<i>Symbolophorus californiensis</i> .....	134
<i>Tarletonbeania crenularis</i> .....	136
Synodontoidei	
Synodontidae	
<i>Synodus</i> spp. ....	138
Gadiformes	
Merlucciidae	
<i>Merluccius productus</i> .....	139
Moridae	
<i>Physiculus</i> spp. ....	143
Macrouridae .....	143
Ophidiiformes .....	143
Bythitidae	
<i>Brosmophycis marginata</i> .....	144
Ophidiidae	

	Page
<i>Chilara taylori</i> .....	145
<i>Ophidion scrippsa</i> .....	145
Lophiiformes	
Ceratioidei .....	146
Gobiesociformes	
Gobiesocidae .....	147
Beloniformes	
Exocoetidae .....	147
Hemiramphidae .....	147
Scomberesocidae	
<i>Cololabis saira</i> .....	147
Atheriniformes	
Atherinidae .....	148
Lampriformes	
Trachipteridae .....	148
Beryciformes	
Melamphaidae	
<i>Melamphaes</i> spp. ....	148
<i>Poromitra</i> spp. ....	151
<i>Scopeloberyx robustus</i> ....	152
<i>Scopelogadus bispinosus</i> ....	152
Syngnathiformes	
Macroramphosidae	
<i>Macroramphosus gracilis</i> .....	152
Syngnathidae	
<i>Syngnathus</i> spp. ....	153
Scorpaeniformes	
Cottoidei	
Agonidae .....	153
Cottidae .....	154
<i>Scorpaenichthys marmoratus</i> .....	154
Cyclopteridae .....	155
Hexagrammidae .....	155
<i>Oxylebius pictus</i> .....	155
<i>Zaniolepis</i> spp. ....	156
Scorpaenoidei	
Scorpaenidae	
<i>Scorpaena</i> spp. ....	156
<i>Sebastes</i> spp. ....	157
<i>Sebastolobus</i> spp. ....	161
Triglidae	
<i>Prionotus</i> spp. ....	161
Perciformes	
Blennioidei	
Blenniidae	
<i>Hypsoblennius</i> spp. ....	162
Clinidae .....	163
Gobioidei	
Gobiidae .....	165
Labroidei	
Labridae .....	166
<i>Halichoeres</i> spp. ....	167

	Page
<i>Oxyjulis californica</i> .....	167
<i>Semicossyphus pulcher</i> .....	168
Pomacentridae .....	169
<i>Chromis punctipinnis</i> .....	169
<i>Hypsypops rubicundus</i> .....	170
Mugiloidei	
Mugilidae	
<i>Mugil</i> spp. .....	170
Percoidei	
Apogonidae	
<i>Howella brodiei</i> .....	170
Bramidae	
<i>Brama</i> spp. .....	170
Carangidae .....	171
<i>Seriola lalandi</i> .....	171
<i>Trachurus symmetricus</i> .....	172
Caristiidae	
<i>Caristius macropus</i> .....	175
Coryphaenidae	
<i>Coryphaena hippurus</i> .....	175
Gerreidae .....	175
Haemulidae .....	175
Kyphosidae	
<i>Girella nigricans</i> .....	176
<i>Medialuna californiensis</i> .....	176
Malacanthidae	
<i>Caulolatilus princeps</i> .....	176
Sciaenidae .....	176
Serranidae .....	178
Scombroidei	
Gempylidae .....	179
Scombridae .....	180
<i>Auxis</i> spp. .....	180
<i>Sarda chiliensis</i> .....	180
<i>Scomber japonicus</i> .....	180
<i>Scomberomorus</i> spp. .....	181
Trichiuridae .....	181
Sphyraenoidei	
Sphyraenidae	
<i>Sphyraena argentea</i> .....	182
Stromateoidei	
Centrolophidae	
<i>Icichthys lockingtoni</i> .....	183
Nomeidae .....	184
Stromateidae	
<i>Peprius simillimus</i> .....	184
Tetragonuridae	
<i>Tetragonurus cuvieri</i> .....	185
Trachinoidei	
Chiasmodontidae .....	185
Pleuronectiformes .....	186
Pleuronectoidei	
Paralichthyidae	

	Page
<i>Citharichthys</i> spp. ....	186
<i>Citharichthys stigmaeus</i> ....	190
<i>Hippoglossina stomata</i> ....	192
<i>Paralichthys californicus</i> ....	193
<i>Syacium ovale</i> ....	194
<i>Xystreurus liolepis</i> ....	194
<b>Pleuronectidae</b>	
<i>Glyptocephalus zachirus</i> ....	195
<i>Hypsopsetta guttulata</i> ....	195
<i>Lepidopsetta bilineata</i> ....	195
<i>Lyopsetta exilis</i> ....	195
<i>Microstomus pacificus</i> ....	196
<i>Parophrys vetulus</i> ....	196
<i>Pleuronichthys</i> spp. ....	198
<i>Pleuronichthys coenosus</i> ....	198
<i>Pleuronichthys decurrens</i> ....	198
<i>Pleuronichthys ritteri</i> ....	198
<i>Pleuronichthys verticalis</i> ....	199
<i>Psettichthys melanostictus</i> ....	200
<b>Soleoidei</b>	
<b>Cynoglossidae</b>	
<i>Sympodus</i> spp. ....	200
<b>Soleidae</b> ....	202
<b>Tetraodontiformes</b>	
<b>Tetraodontidae</b> ....	202
Disintegrated fish larva ....	202
Unidentified fish larva ....	207



## RECENT TECHNICAL MEMORANDUMS

Copies of this and other NOAA Technical Memorandums are available from the National Technical Information Service, 5285 Port Royal Road, Springfield, VA 22167. Paper copies vary in price. Microfiche copies cost \$4.50. Recent issues of NOAA Technical Memorandums from the NMFS Southwest Fisheries Center are listed below:

- NOAA-TM-NMFS-SWFC- 86 Ichthyoplankton and station data for California Cooperative Oceanic Fisheries Investigations survey cruises in 1958.  
E.M. SANDKNOP, R.L. CHARTER, H.G. MOSER, and J.D. RYAN  
(September 1987)
- 87 Ichthyoplankton and station data for California Cooperative Oceanic Fisheries Investigations survey cruises in 1959.  
E.G. STEVENS, R.L. CHARTER, H.G. MOSER, and M.S. BUSBY  
(September 1987)
- 88 Ichthyoplankton and station data for California Cooperative Oceanic Fisheries Investigations survey cruises in 1960.  
D.A. AMBROSE, R.L. CHARTER, H.G. MOSER, and C.R. SANTOS METHOT  
(September 1987)
- 89 Summary of distribution records of the spinner dolphin, *Stenella longirostris*, and the pantropical spotted dolphin, *S. attenuata*, from the western Pacific Ocean, Indian Ocean and Red Sea.  
J.W. GILPATRICK, JR., W.F. PERRIN, S. LEATHERWOOD, and L. SHIROMA  
(October 1987)
- 90 Summary of worldwide locality records of the striped dolphin, *Stenella coeruleoalba*.  
C.E. WILSON, W.F. PERRIN, J.W. GILPATRICK, JR., and S. LEATHERWOOD  
(December 1987)
- 91 Micropatch sampler data.  
R.W. OWEN and C.A. KIMBRELL  
(December 1987)
- 92 Ichthyoplankton and station data for California Cooperative Oceanic Fisheries Investigations survey cruises in 1961.  
E.M. SANDKNOP, R.L. CHARTER, H.G. MOSER, C.A. MEYER, and A.E. HAYS  
(January 1988)
- 93 Ichthyoplankton and station data for California Cooperative Oceanic Fisheries Investigations survey cruises in 1962.  
B.Y. SUMIDA, R.L. CHARTER, H.G. MOSER, and D.L. SNOW  
(January 1988)
- 94 Ichthyoplankton and station data for California Cooperative Oceanic Fisheries Investigations survey cruises in 1963.  
D.A. AMBROSE, R.L. CHARTER, H.G. MOSER, and B.S. EARHART  
(January 1988)
- 95 Ichthyoplankton and station data for California Cooperative Oceanic Fisheries Investigations survey cruises in 1964.  
E.M. SANDKNOP, R.L. CHARTER, H.G. MOSER, C.A. MEYER, and A.E. HAYS  
(January 1988)

